

Available online at [ScienceDirect](https://www.sciencedirect.com)

Resuscitation

journal homepage: www.elsevier.com/locate/resuscitation

Practice Guideline

European Resuscitation Council Guidelines 2025 Special Circumstances in Resuscitation



Carsten Lott^{a,*,1}, Vlasios Karageorgos^{b,c,1}, Cristian Abelairas-Gomez^{d,e}, Annette Alfonzo^f, Joost Bierens^{g,h}, Steve Cantellowⁱ, Guillaume Debaty^j, Sharon Einav^k, Matthias Fischer^l, Violeta González-Salvado^{m,n}, Robert Greif^o, Bibiana Metelmann^p, Camilla Metelmann^{p,q}, Tim Meyer^r, Peter Paal^s, David Peran^t, Andrea Scapigliati^{u,v}, Anastasia Spartinou^{b,w}, Karl Thies^x, Anatolij Truhlar^{y,z,aa}, Charles D. Deakin^{ab}

Abstract

These European Resuscitation Council (ERC) Guidelines 2025 on Special Circumstances in Resuscitation are based on the 2025 Consensus on Science with Treatment Recommendations (CoSTR) of the International Liaison Committee on Resuscitation (ILCOR), reviews by the expert writing group and relevant peer-reviewed literature. The guideline chapter provides guidance for laypeople and healthcare professionals on the modifications required to basic and advanced life support in adults for the prevention and treatment of cardiac arrest for in-hospital and out-of-hospital cardiac arrest. The ERC Guidelines 2025 Paediatric Life Support cover the special circumstances in children.

Keywords: Cardiac arrest, Special circumstances, Adults, Special causes, Special settings, Special patient populations

Introduction

Cardiac arrest, regardless of its aetiology, requires immediate intervention, involving rapid recognition, emergency request for assistance, and high-quality cardiopulmonary resuscitation (CPR), with minimal interruptions. Effective management involves the prompt identification and treatment of reversible causes, often summarised

in the mnemonic '4Hs and 4Ts.' However, standard life support protocols are not universally applicable, as special circumstances need modifications to these guidelines. The ERC categorise these special circumstances into three parts:

1. Special causes leading to specific interventions.
2. Special settings where unique site factors or aetiologies require tailored approaches.

Abbreviations: ABC, Airway-Breathing-Circulation, ACE-I, Angiotensin converting enzyme inhibitors, AED, Automated external defibrillator, AFE, Amniotic fluid embolism, ALS, Advanced life support, BLS, Basic life support, CAB, Circulation - Airway - Breathing, CAD, Coronary artery disease, CKD, Chronic kidney disease, COPD, Chronic obstructive pulmonary disease, CPR, Cardiopulmonary resuscitation, CoSTR, Consensus on Science with Treatment Recommendations, DVT, Deep venous thrombosis, ECG, Electrocardiogram, ECMO, Extracorporeal membrane oxygenation, ECPR, Extracorporeal cardiopulmonary resuscitation, ED, Emergency department, EMS, Emergency medical services, ERC, European Resuscitation Council, ET_{CO2}, End tidal carbon dioxide, HCP, Healthcare professionals, HD, Haemodialysis, ICU, Intensive Care Unit, IHCA, In-hospital cardiac arrest, ILCOR, International Liaison Committee on Resuscitation, IM, Intramuscular, IV, Intravenous, LVAD, Left Ventricular Assist Device, MRA, Mineralocorticoid receptor antagonist, NIV, Non-invasive ventilation, OHCA, Out-of-hospital cardiac arrest, OR, Operating room, PAD, Public access defibrillator, PCI, Percutaneous coronary intervention, PE, Pulmonary embolism, PEA, Pulseless electrical activity, POCUS, Point of care ultrasonography, PPE, Personal protective equipment, RCT, Randomised controlled trial, REBOA, Resuscitative endovascular balloon occlusion of the aorta, ROSC, Return of spontaneous circulation, SCD, Sudden cardiac death, STEMI, ST-elevation myocardial infarction, TCA, Traumatic cardiac arrest, TOE, Transoesophageal echocardiography, VA-ECMO, Veno-arterial extracorporeal membrane oxygenation, VF, Ventricular fibrillation, VT, Ventricular tachycardia, WHO, World Health Organization

* Corresponding author.

E-mail address: lott@uni-mainz.de (C. Lott).

¹ Both share the first authorship.

<https://doi.org/10.1016/j.resuscitation.2025.110753>

0300-9572/© 2025 European Resuscitation Council. Published by Elsevier B.V.

3. Special patient populations, characterised by pre-existing conditions or comorbidities that require different modified treatment strategies.

These ERC Guidelines 2025 on Special Circumstances in Resuscitation are based on the 2025 annual Consensus on Science with Treatment Recommendations (CoSTR) of the International Liaison Committee on Resuscitation (ILCOR).¹ Many topics addressed in these Special Circumstances Guidelines were not part of the ILCOR review. Thus, several recommendations are based on consensus of the ERC Guidelines 2025 Special Circumstances in Resuscitation writing group based on additional systematic or scoping reviews or selected original articles [Table 1](#). The broader methodology used for guideline development is presented in the Executive Summary.² These Guidelines were posted for public comments between 15 May and 30 May 2025. A total of 54 individuals submitted 66 comments, leading to 9 changes in the final version. Subsequently, the feedback was reviewed by the writing group, and the Guidelines was thereafter updated where relevant. The Guidelines were presented to, and approved by, the ERC Board and the ERC General Assembly in June 2025.

For the purpose of these Guidelines, the term CPR relates to the specific technical skills of cardiopulmonary resuscitation (e.g. performance metrics of chest compression and ventilation), whilst resusci-

tation is used as a generic term covering the broader range of skills and interventions. The term bystander is used to describe rescuers who happen to be at the scene to provide help, and the term first responder is used for those who have additional training and are alerted to attend the scene of a cardiac arrest. Healthcare Professionals (HCP) are defined as those who work in any healthcare sector (prehospital or in-hospital). Laypeople are persons not working in the healthcare sector. Basic Life Support (BLS) is defined as initiating the chain of survival, early high-quality chest compression, effective ventilation, and the early use of an AED. Any form of resuscitation education beyond BLS is described generically as advanced life support (neonatal, paediatric and adult life support). Where the term 'ALS' is used, this refers specifically to the ERC adult Advanced Life Support course.

The writing group of these ERC Guidelines 2025 on Special Circumstances in Resuscitation considered the recently introduced ERC approach to diversity, equality, equity, and inclusion and applied it whenever possible, recognising that this is a field for improvement in the development of evidence-informed guidelines. The ERC aims to advance resuscitation practices and improve patient care on a global scale (see [Fig. 1](#)). [Table 1](#) shows changes in the ERC Resuscitation Guidelines Special Circumstances from 2021 to 2025.

Table 1 – This table provides a synopsis of the major changes in the ERC Guidelines 2025 Special Circumstances in Resuscitation. Guidance from the 2021 ERC Guidelines for Resuscitation that is not listed here is still valid and applicable.

Summary of changes		
	ERC guidelines 2021	ERC guidelines 2025
Anaphylaxis	There is no evidence that supports the routine use of either steroids or antihistamines	- Antihistamine administration should not delay the administration of adrenaline. - The ERC recommends against the routine use of corticosteroids.
Resuscitation in hyper/hypokalaemia and other electrolyte disorders	Protect the heart: Give 10 ml calcium chloride 10 % IV by rapid bolus injection. Consider repeating dose if cardiac arrest is refractory or prolonged.	- No evidence for or against use of IV calcium in hyperkalaemia. - ECG – record pre and post IV calcium. - Recommendation against the routine use of sodium bicarbonate in the treatment of hyperkalaemia in non-cardiac arrest cases. - Hyperkalaemia algorithm revised – emphasis on initiation of potassium-lowering treatments. Guidance in cardiac arrest added.
Hyperthermia, malignant hyperthermia and toxin induced hyperthermia		- Figure for treatment of malignant hyperthermia added. - Section on toxin-induced hyperthermia with table added.
Accidental hypothermia and avalanche rescue	Scoring systems established	- Revised Swiss Staging for hypothermia, when core temperature cannot be measured. - Heart rate < 45/min as new high-risk criterion for hypothermia-induced cardiac arrest. - Criteria for qualification for extracorporeal life support do not apply for rewarming of arrested hypothermic patients, e.g. unwitnessed cardiac arrest, asystole as presenting arrhythmia. - New avalanche rescue algorithm. - BLS avalanche rescue algorithm for cases with insufficient personnel on site.

Table 1 (continued)

Summary of changes		
	ERC guidelines 2021	ERC guidelines 2025
Pulmonary embolism		• Specific drug and dosage schemes added.
Coronary thrombosis	• Individualised approach concerning reperfusion strategy in NSTEMI patients was recommended. Brief evaluation to rule out non-coronary causes and performing urgent coronary angiography in case of suspected myocardial ischaemia were suggested.	• Assess 12-lead electrocardiogram after ROSC for ischaemic features and consider repeating if findings are inconclusive, since defibrillation and the time elapsed from ROSC may influence the results • High-certainty evidence from new RCTs discourages routine emergent/early coronary angiography in stable patients without ST-elevation or equivalent on ECG. Immediate angiography is recommended in case of haemodynamic instability or suspected ongoing ischaemia. • Indications for antiplatelet and anticoagulation treatment in this setting.
Toxic agents		• New update on intoxication with opioids.
Traumatic cardiac arrest	• New algorithm established • Invasive procedures introduced	• Algorithm clarified. • Role of chest compressions specified. • REBOA de-emphasised
Cardiac arrest in the catheterisation laboratory	• Mentioned but not specifically included in the management algorithm. • Point of care ultrasonography (POCUS) may be considered to identify reversible causes of cardiac arrest in this setting. • General recommendations were given to consider mechanical CPR, extracorporeal CPR and circulatory support devices in the catheterisation lab.	• Guidance on resuscitation team performance in the catheterisation lab: adequate radiation protective equipment, pre-alerting the surgical team if necessary, clearly defining roles and allowing the interventional cardiologist to focus on corrective procedures rather than leading CPR. • Specifies management of extreme bradycardia/ asystole complicating invasive procedures in the catheterisation lab, consider external or transvenous temporary pacing (included in the management algorithm). • Updated evidence on the role of POCUS and specific indications on the use of transoesophageal echocardiography in this setting. • Updated evidence on the role of mechanical CPR, extracorporeal CPR and circulatory support devices in the catheterisation lab, with specific indications for cardiac arrest in this setting. • New limited evidence on the use of intracoronary adrenaline as an alternative administration route for cardiac arrest in the catheterisation lab, highlighting the need for further research.
Drowning	• First responders not covered	• The introduction of the role of first responders in drowning resuscitation. • Clarifications regarding some topics such as the role of bystanders during rescue, spinal stabilisation, and the emphasis on ventilation by bystanders and first responders.
Cardiac arrest in the operating room	• Less detailed	• Sudden decrease in systolic pressure < 50 mmHg despite interventions is criterion to initiate chest compressions. • Lower initial IV adrenaline dose and increment dosing for perioperative cardiac arrest. • Early ECPR for patients with delayed ROSC. • Open chest compressions as an option solely for trained healthcare professionals. • Emphasis on human factors development.
Local anaesthetic systemic toxicity	• No treatment algorithm	• New detailed algorithm.
Resuscitation in cardiac surgery and Left Ventricular Assist Device patients	• LVAD patients not covered	• Revised algorithm for cardiac arrest after cardiac surgery. • Addition of cardiac arrest algorithm in Left Ventricular Assist Device patients.

(continued on next page)

Table 1 (continued)

		Summary of changes
	ERC guidelines 2021	ERC guidelines 2025
Cardiac arrest in sports	• Short overview	• More robust and comprehensive data on incidence and survival. • Characterisation of causes by age group (± 35 years). • Awareness about risk in recreational and non-elite athletes. • Specific guidance for high-attendance or televised events.
Resuscitation during transport	• Limited to team approach in EMS	• New subchapter Emergency Medical Services includes initial approach of two-member ALS crews, HEMS and air ambulances and the use of arterial line in the prehospital setting. • Inflight cardiac arrest is now expanded by CPR in microgravity.
Asthma & COPD		• New shortened algorithm
Resuscitation in haemodialysis patients	• Focus on haemodialysis unit	• Now focusing on haemodialysis patients rather than on the. • Infographic of resuscitation whilst on dialysis.
Resuscitation in obese patients		• No differences – Standard BLS and ALS recommended.
Resuscitation in patients with pectus excavatum	• Not addressed	• Reduced chest compression depth to 3–4 cm. • Increased force is required to deliver effective chest compressions in case of Nuss bar correction. • Use of anteroposterior pad placement for defibrillation.
Resuscitation during Pregnancy	• Defines maternal cardiac arrest as occurring during pregnancy and up to 6 weeks post-partum. • Recommends manual left uterine displacement; left lateral tilt discussed with caveats. • Recommends delivery within 5 min of collapse if no ROSC. • Use of 4Hs and 4Ts.	• Addendum to the ALS algorithm introducing maternal reversible causes (“4P” – (pre-eclampsia/ eclampsia, puerperal sepsis, placental/ uterine issues, peripartum cardiomyopathy) • adds global mortality statistics and highlights that pregnancy may not be clinically obvious • Suggests manual displacement as preferred, due to practicalities. • De-emphasises 5 min; emphasises immediate preparation for resuscitative hysterotomy. • Practical recommendation to remove fetal monitors to prevent burns. • IV/IO access above diaphragm when possible.

Concise guidelines for clinical practice

General recommendations

- Initiate resuscitation following the standard ALS algorithm in cardiac arrest.
- Always address hypoxia, hypovolaemia, electrolyte disorders, hypothermia, cardiac tamponade, tension pneumothorax, thrombosis, and toxic agents.
- Where appropriate, prioritise treating reversible causes, even if chest compressions are briefly interrupted.

Special causes

Management and prevention of cardiac arrest due to anaphylaxis

- Prompt recognition of anaphylaxis is crucial.
- Recognise anaphylaxis by the presence of airway, breathing, or circulation problems with or without skin and mucosal changes.

- Remove or stop the trigger if immediately feasible.
- Immediately inject intramuscular adrenaline 0.5 mg at first suspicion of anaphylaxis and repeat if no improvement occurs within 5 min.
- Give an IV crystalloid fluid bolus early and monitor the response.

Hyper/hypokalaemia and other electrolyte disorders

Hyperkalaemia (Fig. 2)

- Shift potassium into cells
 - Give 10 units soluble insulin and 25 g glucose IV for treatment of moderate and severe hyperkalaemia. Follow with 10 % glucose infusion at 50 ml/hr for 5 h if pre-treatment blood glucose < 7 mmol/l.
 - Give nebulised salbutamol (10–20 mg) for moderate and severe hyperkalaemia, as an adjunct to Insulin-glucose therapy.
- Antagonise the effect of hyperkalaemia

SPECIAL CIRCUMSTANCES IN ADULT RESUSCITATION

KEY MESSAGES




Fig. 1 – Key messages of the ERC Guideline 2025 Special Circumstances in Resuscitation.

- o Use IV 10 mL 10 % calcium chloride for patients with severe hyperkalaemia with ECG changes.
- o Administer IV 10 mL 10 % calcium chloride and 50 mmol sodium bicarbonate, through separate lines or with flush in between, in the treatment of hyperkalaemic cardiac arrest in all settings of severe hyperkalaemia.
- Remove potassium from the body
 - o Give sodium zirconium cyclosilicate 10 g orally.
 - o Consider dialysis for patients with refractory severe hyperkalaemia.
- Consider ECPR in accordance with local protocols if initial resuscitation attempt is unsuccessful.

Hypokalaemia

- Treatment is guided by the severity of hypokalaemia and presence of symptoms and/or ECG abnormalities.
- Where appropriate replace potassium and correct magnesium deficit concurrently.
- Give IV 20 mmol potassium chloride over 2–3 min, followed by 10 mmol over 2 min in hypokalaemic cardiac arrest then monitor K⁺ level and adjust infusion rate accordingly.

Hyperthermia, malignant hyperthermia and toxin-induced hyperthermia

Hyperthermia

- Measure core temperature to guide treatment.

- Move patient to a cool environment.
- Simple external cooling may involve conductive, convective and evaporative measures (See ERC Guidelines 2025 First Aid).³
- With heat syncope and heat exhaustion quick removal to a cool place, simple external cooling and provision of fluids are sufficient
- With heat stroke prioritise active cooling methods that achieve the most rapid cooling rate such as application of ice and cold-water immersion (Fig. 3).

Malignant hyperthermia



QR code. 1 – Malignant hyperthermia algorithm.

Treatment of malignant hyperthermia, modified from (Hopkins et al., 2021)⁹¹ and (Kollman-Camaiera et al., 2017).¹⁰⁹

- Stop triggering agents immediately. This includes turning off and removing vaporiser and changing the ventilator circuit.
- Give IV 2.5 mg/kg dantrolene as soon as possible.
- Start active cooling.
- Give 100 % oxygen and aim for normocapnia using hyperventilation.
- Change the ventilator. If the ventilator cannot be changed, change charcoal filters




Fig. 2 – Treatment algorithm of hyperkalaemia in adults.

ABCDE – airway breathing circulation disability exposure, ECG – electrocardiography, IV – intravenous, VT – ventricular tachycardia, K – potassium.

- Contact a malignant hyperthermia centre for advice and follow-up.

Toxin-induced hyperthermia

- Minimise exposure and absorption of the toxin.
- Use active cooling techniques. Antipyretics have no benefit as central thermoregulatory mechanisms are affected by toxins.

Accidental hypothermia and avalanche rescue

Accidental hypothermia

- Check vital signs for up to one minute in an unconscious hypothermic patient (Fig. 5).
- Measure core temperature with a low reading thermometer to diagnose accidental hypothermia.
- Use the Swiss Staging System if core temperature cannot be measured.




Fig. 3 – Treatment algorithm of hypokalaemia in adults.

ABCDE – airway breathing circulation disability exposure, ECG – electrocardiography, IV – intravenous.




Fig. 4 – Emergency treatment of hyperthermia.

CPR – cardiopulmonary resuscitation, ALS – advanced life support. Footnote: *Core temperature should be measured, e.g. tympanic, rectal, or oesophageal. ** Dry mouth, thirsty, hypotensive. ***Return to activity considerations following a diagnosis of exertional heat illness and exertional head stroke have been published recently elsewhere. ^{96,99}

- Transfer hypothermic patients with risk factors for imminent cardiac arrest and those in cardiac arrest directly to an extracorporeal life support (ECPR) centre for rewarming.
- Delay CPR or use intermittent CPR in hypothermic cardiac arrest patients with a core temperature below 28 °C when immediate or continuous CPR is not feasible.
- Delay further defibrillation attempts if ventricular fibrillation (VF) persists after three shocks until core temperature is > 30 °C.
- Below 30 °C adrenaline will accumulate and may have more detrimental than beneficial effects. Give IV 1 mg adrenaline once to facilitate ROSC unless planning imminent initiation of ECPR. Increase administration intervals for adrenaline to 6–10 min if the core temperature is 30–35 °C.
- Consider use of a mechanical CPR device if transport is prolonged, or when there are difficulties with the terrain.
- Base in-hospital prognostication of successful rewarming on the Hypothermia Outcome Prediction after Extracorporeal Life Support (HOPE) score.
- Rewarm hypothermic arrested patients with veno-arterial extracorporeal membrane oxygenation (VA-ECMO).

- Initiate non-extracorporeal life support rewarming if an ECPR centre cannot be reached within a reasonable time (e.g. 6 h).

Avalanche rescue

- Base initiation of CPR in cardiac arrest on core temperature, burial time, and airway patency (Fig. 6).
- Consider proceeding according to the AvaLife algorithm, in multi-burial avalanche accidents with BLS providers only and insufficient numbers of rescuers.

Thrombosis

Pulmonary embolism

- Consider pulmonary embolism in all patients with sudden onset of progressive dyspnoea and absence of known heart or pulmonary disease.
- Obtain 12-lead ECG (exclude acute coronary syndrome, look for right ventricular strain).




Fig. 5 – Emergency treatment of accidental hypothermia.

^a Decapitation; truncal transection; whole body decomposed or whole body frozen solid (chest wall not compressible). ^b Systolic blood pressure < 90 mmHg is a reasonable prehospital estimate of cardiocirculatory instability but for in-hospital decisions, the minimum sufficient circulation for a deeply hypothermic patient (e.g., <28 °C) has not been defined. ^c Swiss staging of accidental hypothermia. ^d Direct transport to an ECLS centre is recommended in an arrested hypothermic patient. In remote areas, transport decisions should balance the risk of increased transport time with the potential benefit of treatment in an extra corporal life support centre. Transfer should be commenced if such a centre can be reached within 6 h. ^e Warm environment, chemical, electrical, or forced air heating packs or blankets, and warm IV fluids (38–42 °C). In case of cardiac instability refractory to medical management, consider rewarming with extra corporal life support. ^f If the decision is made to stop at an intermediate hospital to measure serum potassium, a hospital enroute to an extra corporal life support centre should be chosen. The HOPE score ^{156,158,162} should not be used in children, consider expert consultation instead. ECLS extracorporeal life support, ROSC return of spontaneous circulation, SBP systolic blood pressure.

- Identify haemodynamic instability and high-risk pulmonary embolism.
- Perform bedside echocardiography.
- Initiate anticoagulation therapy (heparin 80 IU/kg IV) during diagnostic process, unless signs of bleeding or absolute contraindications.
- Confirm diagnosis with computed tomographic pulmonary angiography.
- Consider surgical embolectomy or catheter-directed treatment as alternative to rescue fibrinolytic therapy in rapidly deteriorating patients.

Cardiac arrest due to pulmonary embolism.

- Low ETCO₂ values (<1.7 kPa/13 mmHg) while performing high quality chest compressions may support a diagnosis of pulmonary embolism, although it is a non-specific sign.
- Use fibrinolytic drugs for cardiac arrest when pulmonary embolism is the suspected cause of cardiac arrest.
- Use fibrinolytic drugs or surgical embolectomy or percutaneous mechanical thrombectomy for cardiac arrest when pulmonary embolism is the known cause of cardiac arrest.
- Consider ECPR as a rescue therapy for selected patients with cardiac arrest when conventional CPR is failing in settings in which it is implemented.




Fig. 6 – Decision-making algorithm for advanced management of critically buried avalanche victims in cardiac arrest.¹⁸⁸

CPR – cardiopulmonary resuscitation, ROSC – return of spontaneous circulation, ECLS – extracorporeal life support, HOPE – Hypothermia Outcome Prediction after ECLS rewarming for hypothermic arrested patients, ECG – electrocardiography. ^a Decapitation, truncal transection, whole body decomposed. ^b An obstructed or blocked airway requires that both the nose and mouth be completely filled with compact snow or debris. ^c With a deeply hypothermic patient (<28 °C), if rescue is too dangerous consider delayed CPR and if transport is difficult consider intermittent CPR. ^d If core temperature measurement is not available, hypothermic CA may be considered, at the rescuer's discretion, despite a burial duration of < 60 min in a victim with a patent airway and no vital signs when there is a possibility of very rapid cooling (e.g. burial during ascent, low body mass index or small persons, minimally dressed, sweating before burial). ^e In-hospital prognostication of successful rewarming in an avalanche victim should include estimation of the survival probability using the HOPE score. If any doubt exists whether the avalanche victim may have asphyxiated despite critical burial, the HOPE score should be calculated using the NON-ASPXYXIA option. This will reduce the risk of under-treatment. If the HOPE score cannot be determined, the combination of potassium < 7 mmol L⁻¹ and a temperature < 30 °C may be used instead to help indicate ECLS rewarming.

- Set-up a multidisciplinary team for making decisions on management of high-risk pulmonary embolism depending on local resources.

Coronary thrombosis

- Enhance health education to recognise symptoms and minimise delays in seeking medical care
- Promote BLS training for likely rescuers of high-risk groups.
- Strengthen regional networks to ensure timely percutaneous coronary intervention (PCI).
- Transfer the patient to a centre with PCI capability and activate existing STEMI networks in case of ST-elevation or suspected ongoing ischaemia.
- In patients with sustained ROSC and ST-elevation on ECG:
 - o Perform immediate coronary angiography (and PCI if required) within 120 min of diagnosis.
 - o Consider fibrinolysis in pre-hospital and non-PCI-capable settings if greater delay is expected.
- In patients with sustained ROSC and no ST-elevation on ECG:

- o Consider immediate coronary angiography (and PCI if required) if the patient is haemodynamically unstable or shows signs of ongoing ischaemia.
- o In stable patients without ischaemic signs, emergent cardiac catheterisation laboratory evaluation should not be systematic and can be delayed if there is no estimated high probability of acute coronary occlusion.
- o Assess for non-coronary-causes if the clinical context suggests an alternative aetiology of the arrest.
- o Unless on-going resuscitation is considered futile, transfer patients without sustained ROSC with ongoing CPR, to a PCI centre for consideration for angiography or ECPR depending on available resources and team expertise.

Toxic agents

- Ensure your personal safety (Fig. 7), as direct skin contact (e.g., mouth-to-mouth ventilation) might transmit toxic agents.

- Assess all patients in cardiac arrest for potential intoxication.
- Reduce absorption, consider using specific treatment measures as antidotes, decontamination and enhanced elimination.
- Administer antidotes, where available, as soon as possible.
- Be prepared to continue resuscitation for a prolonged period of time, as the toxin concentration may fall as it is metabolised or excreted during extended resuscitation measures.
- Consult regional or national poison centres for information on treatment of the poisoned patient.

Traumatic cardiac arrest (TCA)

- Traumatic cardiac arrest is different from cardiac arrest due to medical causes; this is reflected in the treatment algorithm (Fig. 8).
- The response to traumatic cardiac arrest is time-critical and success depends on a well-established chain of survival, including focused pre-hospital and specialised trauma centre care.
- Early and aggressive management of reversible causes (e.g. haemorrhage control, airway management, chest decompression) is essential for survival.
- Ultrasound aids in identifying the cause of cardiac arrest and guides resuscitative interventions.

Special settings

Cardiac arrest in the catheterisation laboratory

- Promote adequate training of the staff in technical skills and ALS, and consider periodic emergency drills.
- Ensure emergency equipment is readily available and functional.
- Plan elective procedures carefully to minimise potential complications and promote the use of safety checklists.
- Consider echocardiography in case of haemodynamic instability or suspected complication.
- Resuscitate according to the ALS algorithm, BUT modify:
 - Apply 3 consecutive shocks in case of shockable rhythm.
 - Consider external or transvenous pacing for extreme bradycardia.
- Consider, in selected cases depending on clinical context, team expertise and availability:

- Mechanical CPR, if manual compression is not feasible or safe for the provider.
- Extracorporeal CPR in selected patients with refractory cardiac arrest, especially if it allows for critical procedures to correct reversible causes.
- Circulatory support devices, for selected patients in cardiogenic shock after achieving ROSC.

Drowning

- Rescuers and first responders should prioritise their safety and use the safest rescue technique.
- Bystanders should call for professional help and use rescue techniques with which they feel confident, based on their competencies.
- First responders should use rescue material and flotation devices they are trained to use.
- Spine immobilisation in water should not delay removing the victim from the water when resuscitation is required.
- Start with 5 ventilations using 100 % inspired oxygen when available, continue with standard CPR protocol.
- Airway and ventilation equipment can be used when the provider is trained appropriately.
- Increase ventilation pressure gradually when a high inspiration pressure is needed, in order to avoid stomach inflation.
- Consider scaling up to ECPR if initial resuscitation is unsuccessful, in accordance with local protocols.
- Follow the recommendations for hypothermia.

Cardiac arrest in the operating room (OR)

- Prevent and mitigate the risk of cardiac arrest through pre-operative screening and identification of high-risk patients, clear communication with surgeons about potential critical procedures, advanced monitoring and continuous presence of an anaesthesiologist during patient instability.
- Start chest compressions if the systolic blood pressure suddenly decreases below 50 mmHg, in association with fall in ETCO₂, despite appropriate interventions.
- Inform the surgeon and the operating room team of the cardiac arrest.
- Initiate high-quality chest compressions and adjust the height of the operating table to improve work efficiency.
- Ensure the airway is secure, review the ETCO₂ tracing, and deliver effective ventilation, administering 100 % oxygen. Exclude unrecognised oesophageal intubation.
- Use ultrasound to guide resuscitation addressing the reversible causes.
- Exclude tension pneumothorax.
- Consider early ECPR as therapy for selected patients when conventional CPR is failing.
- Trained healthcare professionals may consider open chest cardiac compression in specific cases as an alternative, if ECPR is unavailable.
- Human factors are crucial to improve survival of intraoperative cardiac arrest – ensure familiarity with equipment, assign strategies and roles during surgical team time-outs and include perioperative cardiac arrest in multidisciplinary and interprofessional team training, in-situ simulation, and ALS courses.




Fig. 7 – Management of toxic exposure.




Fig. 8 – Traumatic cardiac arrest algorithm.

ROSC – return of spontaneous circulation, ALS – advanced life support. This algorithm should be used to guide the management of traumatic cardiac arrest (TCA) and peri-arrest trauma patients with impending circulatory collapse. In contrast to the sequential ALS algorithms, it is a framework for context-sensitive decisions on clinically prioritised interventions. The aim is to treat the most urgent, reversible cause based on individualised treatment (e.g. patients with a cardiac tamponade need urgent thoracotomy; patients with pelvic fracture in haemorrhagic shock but with a stable airway may benefit more from immediate application of a pelvic binder and rapid transfer without airway intervention).

Local anaesthetic systemic toxicity

- Stop the local anaesthetic if possible.
- Hyperventilate the patient to increase plasma pH if metabolic acidosis is present.
- Give a lower adrenaline dose ($\leq 1 \mu\text{g kg}^{-1}$ instead of 1 mg IV bolus)
- Give an initial IV bolus of 20 % lipid emulsion at 1.5 mL kg^{-1} over 1 min, followed by an infusion at $0.25 \text{ mL kg}^{-1} \text{ min}^{-1}$ but do not exceed a maximum cumulative dose of 12 mL kg^{-1} IV 20 % lipid emulsion.
- If ROSC has not been achieved at 5 min, double the rate of lipid infusion and give a maximum of two additional lipid boluses at 5-minute intervals until ROSC has been achieved.

- Consider prolonged resuscitation ($>1 \text{ h}$) and ECPR
- Treat seizures by administering benzodiazepines

Cardiac surgery

- Confirm cardiac arrest by clinical signs and pulseless pressure waveforms.
- Consider ultrasound to identify reversible causes.
- Provide up to 3 consecutive shocks in VF/pVT
- Use epicardial pacing at maximum output in asystole or extreme bradycardia
- Perform re-sternotomy up to 10 days post-surgery within 5 min regardless of the patient's location.

- Provide open cardiac compressions once the chest is reopened.
- Reduce IV adrenaline dose (0.05–0.1 mg)
- Consider ECPR for prolonged resuscitation or minimally invasive cases where reopening may be delayed

Left ventricular assist device (LVAD) patients

- Immediately activate specialised teams for unconscious LVAD patients.
- Start CPR while simultaneously attempting to restore device function if multiple rescuers are available.
- Consider delaying CPR for up to 2 min to attempt device restoration if a single rescuer is present.
- Troubleshoot device issues as a priority, following relevant protocols.

Cardiac arrest in sports

- Screening as primary prevention plays an important role, but remains controversial.
- All sports and exercise facilities should undertake a risk assessment which considers the likelihood and consequence of cardiac arrest and put in place mitigation strategies to reduce the risk.
- Gain immediate and safe access to the field of play.
- Awareness programs in sport events have proven to be feasible to raise awareness amongst target groups not yet involved with cardiac arrest.

Emergency medical services

- Healthcare professionals should provide resuscitation at the scene rather than undertake ambulance transport with ongoing resuscitation, unless there is an appropriate indication to justify transport (bridging to in-hospital treatment).
- Consider mechanical CPR for transport with ongoing resuscitation.
- Consider obtaining invasive arterial blood pressure to guide resuscitation and post-resuscitation care already in the prehospital setting if it is feasible.
- EMS systems should use registry and data provided from equipment data (e. g. defibrillators) for debriefing and continuous quality improvement.

Inflight cardiac arrest and microgravity resuscitation

Inflight cardiac arrest

- Healthcare professional help should be sought (in-flight announcement).
- The rescuer should kneel in the leg-space in front of the aisle seats to perform chest compressions if the patient cannot be transferred within a few seconds to an area with adequate floor space (galley).
- Overhead-CPR is a possible option in limited space environments.
- Airway management should be based on the equipment available and the expertise of the rescuer.
- If the planned route leads over an area where no airport can be reached for a longer period of time with high possibility of ROSC during an ongoing resuscitation, consider an early diversion.




Fig. 9 – CPR after drowning for rescuers with a duty to respond.

CPR – cardiopulmonary resuscitation, EMS – emergency medical services, AED – automatic external defibrillator.

- Consider risks of diversion if ROSC is unlikely and give appropriate recommendations to the flight crew.
- If CPR is terminated (no ROSC) there is no medical need for flight diversion – follow airline policy.

Microgravity resuscitation

- Airway management, defibrillation and IV/IO access are similar to terrestrial ALS, but only once the patient is secured.
- Consider mechanical CPR
- Consult telemedicine support during cardiac arrest in low earth orbit if feasible and manpower allows.
- The crewmember with the highest medical qualification should decide on termination of resuscitation, consulting telemedicine support.




Fig. 10 – Cardiac arrest during dialysis.

Abbreviations: CPR – cardiopulmonary resuscitation; EMS – emergency medical services; K⁺ – potassium; HD – haemodialysis; ROSC – return of spontaneous circulation; POCT – point of care testing.

Cruise ship

- Use all healthcare resources immediately (personal, equipment).
- Activate helicopter emergency medical service if close to the coastline.
- Consider early telemedicine support.
- Have all equipment needed for ALS available on board.
- In case of insufficient number of health care professionals to treat CA, call for further medical staff via an on-board announcement.

Special patient groups

Asthma and chronic obstructive pulmonary disease

- Treat life-threatening hypoxia with 100 % oxygen.
- Check for evidence of (tension) pneumothorax.
- Provide endotracheal intubation (because of high inflation pressures).
- Consider manual decompression and disconnection from ventilator to manage dynamic hyperinflation.
- Consider ECPR in accordance with local protocols if initial resuscitation efforts are unsuccessful.

Cardiac arrest in haemodialysis patients

- Assign a trained dialysis nurse or technician to operate the dialysis machine.

- Stop dialysis and return the patient's blood volume with a fluid bolus.
- Disconnect from dialysis machine (unless defibrillation-proof) and beware of wet surfaces.
- Leave dialysis access open and use for drug administration.
- Dialysis may be required in the early post resuscitation period.

Resuscitation in obese patients

- Obese patients should receive standard resuscitation treatment – no deviation from standard BLS and ALS is needed.

Resuscitation in patients with pectus excavatum

- Consider reduced chest compression depth of 3–4 cm.
- In the case of a Nuss bar correction, substantially increased force is required to deliver effective chest compressions.
- Consider early implementation of ECPR if chest compressions are ineffective.
- Use anteroposterior pad placement for defibrillation using standard energies.

Cardiac arrest in pregnancy

- Consider pregnancy in any collapsed woman of childbearing age.
- Pregnant and peripartum women can deteriorate to cardiac arrest anywhere and medical facilities and services must be prepared for such events.

- Obstetric-specific early warning systems enable early recognition of deteriorating pregnant patients.
- Relieve aortocaval compression as early as possible and maintain it throughout resuscitation. Manual left uterine displacement in maternal cardiac arrest is suggested due to practicalities.
- Systematically address the 4Hs and 4Ts and seek pregnancy-specific causes of cardiac arrest, the 4Ps: Pre-eclampsia and eclampsia, Puerperal sepsis, Placental and uterine complications and Peripartum cardiomyopathy.
- Resuscitative hysterotomy is a time-sensitive intervention. Preparation to perform the procedure should be undertaken early.
- Resuscitative hysterotomy should be performed as soon as possible at the site of cardiac arrest by a skilled team.
- Post-resuscitation care in pregnant and postpartum women requires a multidisciplinary approach.

The evidence informing the guidelines

Special causes

Management and prevention of cardiac arrest due to anaphylaxis

There is no universally accepted definition of anaphylaxis. It is an acute systemic hypersensitivity reaction, usually rapid in onset and may be fatal if not appropriately managed. Diagnosis of anaphylaxis is clinical.

The estimated incidence of anaphylaxis is 1.5–7.9 per 100,000 persons per year in Europe.⁴ The most common triggers of anaphylactic reactions are food, medications and hymenoptera venom.^{5–7}

This guidance on anaphylaxis is based on the most recent ILCOR CoSTR,⁸ guidelines and updates from the World Allergy Organization Anaphylaxis Committee,⁹ European Academy of Allergy and Clinical Immunology,¹⁰ North American Practice Parameter,¹¹ Australasian Society of Clinical Immunology and Allergy,¹² the findings from the UK National Audit Project (NAP 7) of perioperative anaphylaxis.¹³ Most of these recommendations are based on observational data, good practice statements and expert consensus.

Recognition of anaphylaxis

According to the World Allergy Organization Anaphylaxis Committee, anaphylaxis is likely when at least one of the following criteria is fulfilled⁹:

1. Acute onset within minutes to several hours, with simultaneous involvement of the skin, mucosal tissue, or both (e.g. generalised hives, pruritus or flushing, swollen lips, tongue or uvula) and one of the following:
 - a. Respiratory compromise (e.g., dyspnoea, wheeze/bronchospasm, stridor, reduced peak expiratory flow, hypoxemia)
 - b. Reduced blood pressure or associated symptoms of end-organ dysfunction (e.g., hypotonia, syncope, incontinence)
 - c. Severe gastrointestinal symptoms (e.g., severe crampy abdominal pain, repetitive vomiting), especially after exposure to non-food allergens

2. Acute onset of hypotension or bronchospasm or laryngeal involvement after exposure to a known or highly probable allergen for that patient (minutes to several hours), even in the absence of typical skin involvement.

Anaphylaxis can occur without any skin involvement or with skin involvement remote to the site of exposure, and an anaphylactic reaction may initially present only with respiratory or cardiovascular involvement. During general anaesthesia or mechanical ventilation in emergency or intensive care settings, anaphylaxis may clinically manifest through suddenly increased ventilation pressures and prolonged expiration, combined with a decrease in blood pressure, with or without skin and mucosal changes.

Early recognition of symptoms suggesting anaphylactic reaction, should elicit immediate reaction.

The initial approach if suspecting an anaphylactic reaction

Stop or remove the trigger if possible – stop or remove any potential or probable trigger. Do not delay treatment if removing the trigger is not feasible.

Ensure the patient is lying – do not sit or stand the patient up – fatal anaphylaxis has been associated with upright posture or patient sitting/standing up.^{14–16} Position patients according to their symptoms and specifically consider:

- o Supine position with raised legs, to improve venous return.
- o Sitting position with legs stretched if airway or breathing symptoms are present without circulation problems.

Give adrenaline

Administer IM adrenaline as first-line drug for the management of anaphylaxis.¹⁷ Initial IM adrenaline both aims provide haemodynamic support as well as acting to inhibit mast cell degranulation. The preferred site is the lateral mid-thigh,¹⁸ standard dose for adults is 0.5 mg.¹⁰ This should be repeated after 3–5 min, according to clinical response.¹⁹ Auto-injectors or intranasal adrenaline applicators can be used for early self-administration or injection by a rescuer. If symptoms fail to resolve or the patient is in a monitored area e.g. intensive care unit (ICU), operating room (OR), emergency department (ED), IV administration of adrenaline, starting at a dose of 0.1 $\mu\text{g kg}^{-1} \text{ min}^{-1}$ titrated according to response can be administered by appropriately trained personnel when patients are fully monitored.²⁰

In a patient with upper airway obstruction or bronchospasm, consider adding nebulised adrenaline.^{10,21} However, systemic absorption of inhaled adrenaline is minimal and should not delay concurrent IM adrenaline administration.²²

Secure vascular access – give intravenous fluids

Anaphylaxis may result in severe hypovolaemia because of peripheral vasodilation and increased vascular permeability (distributive shock).^{23,24} Fluid resuscitation is essential, in addition to the vasoconstrictive effect of adrenaline. Immediately after the first adrenaline dose, ensure adequate vascular access (e.g. wide-bore IV catheter, intraosseous access). Administer crystalloid solution in bolus doses of 10–20 mL kg^{-1} and repeat according to the patient's response. Large volumes of fluids might be required to restore haemodynamic stability.

Give oxygen

Provide 15 L min⁻¹ 100 % oxygen to every patient with an anaphylactic reaction and titrate to an oxygen saturation targeting 94–98 %.²⁵

Other medications to support the circulation

In the case of no response to adrenaline, consider administration of a second vasoactive drug, such as noradrenaline or vasopressin.^{10,21}

Role of steroids and antihistamines in the management of anaphylaxis

H1- antihistamines antagonise the pro-inflammatory action of histamine, which can relieve the cutaneous symptoms of histamine release. However, their onset of action is slow, and they should not delay the administration of adrenaline.

Caution: Rapid IV administration of first generation H1-antihistamines might cause or exaggerate hypotension.²⁶

Corticosteroid administration to prevent protracted symptoms or biphasic reactions has been based on low-certainty evidence.^{27,28} Recent studies indicate increased risk for negative outcomes (second dose of adrenaline, hospitalisation or ICU admission) in patients treated with corticosteroids in the pre-hospital setting.^{29,30} Based on experts' opinion, the ERC does not recommend the routine use of corticosteroids in the management of anaphylactic reactions.

Considerations for cardiac arrest in anaphylaxis

In a case-series of perioperative cardiac arrest, 94 % of patients with anaphylactic cardiac arrest had initial pulseless electrical activity.¹³

Expert consensus is to follow the standard ALS protocol, including IV adrenaline administration and correction of relevant reversible causes, in particular hypovolaemia and hypoxia. Consider ECPR in cases of refractory cardiac arrest, specifically in hospitals where this can be implemented.⁸

Hyper/hypokalaemia and other electrolyte disorders

Electrolyte abnormalities are a recognised cause of arrhythmias and cardiac arrest, potassium disorders (hyperkalaemia and hypokalaemia) are most commonly seen in clinical practice and a U-shaped association between serum K⁺ and mortality has been shown.³¹

Hyperkalaemia has also been subject to a recent ILCOR review (2025).³²

Hyperkalaemia – Non-cardiac arrest

Hyperkalaemia occurs in 1–10 % of hospitalised patients.^{33–35} In patients with chronic kidney disease (CKD), hyperkalaemia results in a significantly higher risk of in-hospital major adverse cardiovascular events and arrhythmias compared to patients without hyperkalaemia.³⁶ Dialysis patients are also more prone to severe hyperkalaemia than non-dialysis patients (45.8 % vs 10.3 %) during hospitalisation.³⁵ In-hospital mortality is significantly higher in patients with hyperkalaemia (18.1 %) compared to those with hypokalaemia (5 %) or normokalaemia (3.9 %).³³ In-hospital mortality is 3.9-fold higher in patients with a serum K⁺ > 6.5 mmol L⁻¹ when compared with normokalaemic patients.³⁷

Renal and cardiac disease often co-exist and these patients are at high risk of hyperkalaemia exacerbated by drug therapy (e.g ACE-inhibitors, angiotensin II antagonists and mineralocorticoid receptor antagonists). However, down-titration or stopping these drugs in response to hyperkalaemia is associated with worse patient

outcomes.^{38–40} There is now a role for potassium binders (sodium zirconium cyclosilicate and patiomer) to provide cardio-renal protective therapy.^{41,42}

Hyperkalaemia is evident with a serum potassium (K⁺) concentration greater than 5.5 mmol/L, although hyperkalaemia is a continuum, and its severity guides treatment. Hyperkalaemia is classified as 'mild' (K⁺ 5.5–5.9 mmol L⁻¹), 'moderate' (K⁺ 6.0–6.4 mmol L⁻¹), 'severe' (K⁺ ≥ 6.5 mmol L⁻¹), or 'extreme' (K⁺ ≥ 9.0 mmol L⁻¹).

The risk of hyperkalaemia increases with multiple simultaneous risk factors (e.g. concomitant use of ACE-I and/or mineralocorticoid receptor antagonist (MRA) in the presence of CKD).

Diagnosis of hyperkalaemia

Consider hyperkalaemia in all at risk patients (e.g. renal failure, heart failure, diabetes mellitus, chronic liver disease) with an arrhythmia or cardiac arrest. Limb weakness, flaccid paralysis or paraesthesia may be indicators of severe hyperkalaemia.

Confirm hyperkalaemia using point-of-care testing, as formal laboratory samples will take time.^{43–45}

ECG changes may reflect the severity and rate of rise of serum K⁺,^{46,47} but it may be normal even in severe hyperkalaemia. When the diagnosis of hyperkalaemia can be established based on the ECG, treatment can be initiated before the laboratory result is available.⁴⁶ These ECG signs of hyperkalaemia may develop progressively and include (Fig. 11a):

- Tall, peaked (tent) T waves (i.e. T wave larger than R wave in more than one lead);
- First degree heart block (prolonged PR interval > 0.2 s);
- Flattened or absent P waves;
- Widened QRS (>0.12 s);
- Sine wave;
- Ventricular tachycardia;
- Bradycardia;
- Cardiac arrest (PEA, VF/VT, asystole).

In patients with severe hyperkalaemia, arrhythmias or cardiac arrest have been shown to occur in 15 % of patients within 6 h of the presenting ECG.⁴⁶ Therefore, delays in treatment may have serious consequences.

Emergency treatment of hyperkalaemia

Treatment is guided by the severity of hyperkalaemia and the presence of ECG changes. Avoid delay in initiating potassium-lowering treatments (i.e. insulin-glucose, salbutamol and sodium zirconium cyclosilicate). Follow a systematic approach as outlined in the hyperkalaemia treatment algorithm (Fig. 2). The treatment of mild hyperkalaemia is not within the scope of this guideline. Systematic reviews of the pharmacological treatment of hyperkalaemia have been conducted both by Cochrane and ILCOR.^{32,48,49}

Insulin and glucose. This is the most effective and reliable K⁺-lowering therapy which works by shifting K⁺ into cells. The conventional dose of soluble insulin (10 units) has been shown to reduce serum K⁺ by 0.7–1.4 mmol L⁻¹.^{32,50} Studies have suggested a possible dose-dependent effect of insulin^{51–55} and a potential correlation between the severity of hyperkalaemia and the degree of K⁺-lowering with 10 units insulin.⁵⁶ These findings are particularly relevant in the resuscitation setting.

Hypoglycaemia remains a major iatrogenic risk with a reported incidence of up to 28 %.^{53,57–62} Reducing the dose of insulin (5 units) did not significantly lower the risk of hypoglycaemia (blood




Fig. 11 – Effect of IV calcium on ECG in hyperkalaemia.

ECG on admission (a) and following pre-emptive administration of 20 ml 10 % calcium gluconate IV (b) in a patient who presented with a flaccid paralysis. Response to IV calcium shown with narrowing of QRS complex. Blood gas analyser was unavailable and formal lab result later

glucose $< 4 \text{ mmol L}^{-1}$).^{51,53,60,61} A scoping review ($n = > 15,000$) found that the most consistent risk factor for hypoglycaemia is a low pre-treatment blood glucose value.⁶³ A threshold of $< 7 \text{ mmol L}^{-1}$ blood glucose has been consistently identifies patients at increased risk of iatrogenic hypoglycaemia.^{55,64–67} In one study, administration of 50 g glucose over 4 h resulted in hypoglycaemia in 6.1 % of patients.⁵⁹ Based on these collective findings and on evidence that this strategy is effective,⁶⁸ a modified protocol which is

recommended in the ERC algorithm (Fig. 2) has been developed in order to reduce iatrogenic hypoglycaemia.⁵⁰

Salbutamol. Salbutamol is a beta-2 adrenoceptor agonist and promotes the intracellular shift of K^+ . A meta-analysis including seven studies administering 10–20 mg inhaled salbutamol demonstrated a reduction in serum K^+ by 0.9 mmol L^{-1} within 120 min.³² Early studies suggested that salbutamol may be less effective in

patients receiving non-selective beta-blockers and in up to 40 % of patients with end-stage kidney disease.^{69,70} On this basis, monotherapy has not been advised. Studies combining insulin-glucose and beta-2 agonists were more effective than either treatment alone.^{69,70} The ILCOR review noted a K^+ -reduction of 1.2 mmol L^{-1} with combined therapy and supported the use of both drugs.³² Thus, the ERC recommends simultaneous administration of insulin-glucose and beta-2-agonists for reducing potassium values. Insulin-glucose and salbutamol are effective for 4–6 h, after which rebound hyperkalaemia may occur.

Intravenous calcium salts (calcium chloride or gluconate). Intravenous calcium has been used for decades for the treatment of hyperkalaemia, particularly in the presence of ECG changes, despite lacking clinical evidence. A recent randomised trial suggested harm from the routine use of IV calcium during OHCA; however patients with hyperkalaemia were excluded.⁷¹ In resuscitation practice, the use of IV calcium has been restricted to the treatment of electrolyte disorders (i.e. hyperkalaemia, hypocalcaemia, hypermagnesaemia) and calcium channel-blocker overdose. In clinical practice, the improvement in ECG changes after IV calcium may be seen even before K^+ -lowering drugs have taken effect (Fig. 11).

The rationale for the use of IV calcium in hyperkalaemia and hypermagnesaemia is 'antagonism' of its effect on cardiac and skeletal muscle (i.e. membrane stabilisation), but efficacy may be influenced by the dose and rate of administration. An alternative mechanism has been proposed in an animal study where IV calcium was shown to restore 'conduction' resulting in improvement of hyperkalaemia-induced ECG changes.⁷² In pre-eclampsia, IV calcium ameliorates limb weakness, respiratory depression and cardiac effects induced by iatrogenic hypermagnesaemia.⁷³

The recent ILCOR systematic review included only one non-cardiac arrest study of IV calcium in adults and found no evidence to support a clinical effect of IV calcium in hyperkalaemia, although considered the study to have a critical risk of bias.³²

The Cochrane reviews included no studies evaluating the effect of IV calcium, but suggested that withholding IV calcium in patients with severe hyperkalaemia or ECG changes or both is not advised given potentially life-threatening consequences.⁴⁸ There is currently insufficient evidence for or against the use of IV calcium in the treatment of hyperkalaemia, therefore the ERC continues to recommend the administration of IV calcium to patients most at risk of arrhythmias (i.e. severe hyperkalaemia with ECG changes present).

Sodium bicarbonate. The ILCOR review included five studies in adults using variable doses of bicarbonate (50–390 mmol) and found a K^+ -lowering of only 0.1 mmol L^{-1} within 60 min.³² The Cochrane review also concluded that there is no evidence to support the use of sodium bicarbonate in hyperkalaemia.^{48,49} Therefore, the ERC recommends against routine use of sodium bicarbonate in the treatment of hyperkalaemia in non-cardiac arrest cases.

Potassium binders. Novel potassium binders were not included in the ILCOR review, but sodium zirconium cyclosilicate (SZC) and Patiromer are well tolerated and have a role in both acute and chronic hyperkalaemia. SZC has a more rapid onset of action (within

1 h) than Patiromer.⁷⁴ Within 48 h, SZC lowers serum K^+ by 1.1 mmol/L and shows greater efficacy with higher potassium levels (1.5 mmol/L K^+ -reduction in patients with a serum $K^+ > 6.0 \text{ mmol/L}$).

Indications for dialysis

Dialysis is the most definitive treatment for hyperkalaemia, and the main indications with hyperkalaemia are:

- Severe life-threatening hyperkalaemia with or without ECG changes or arrhythmia;
- Hyperkalaemia resistant to medical treatment;
- End-stage kidney disease;
- Oliguric acute kidney injury (urine output $< 400 \text{ mL/day}$);
- Marked tissue breakdown (e.g. rhabdomyolysis).

Following dialysis, rebound hyperkalaemia may also occur.

Pre-hospital setting and hyperkalaemia

The management and identification of patients with hyperkalaemia in the pre-hospital setting can be challenging, because of the limited diagnostic options.

Resuscitation of hyperkalaemic cardiac arrest

Hyperkalaemia is the most common metabolic disturbance associated with cardiac arrest and is potentially reversible, with a reported IHCA incidence of 1 %⁷⁵ to 12 %, ⁷⁶ with PEA as the most common initial arrest rhythm.⁷⁷ The ILCOR review included only one study of hyperkalaemia in cardiac arrest (high risk of bias due to resuscitation time). This review describes decreased rates of ROSC with calcium, sodium bicarbonate or both.³²

A study of hyperkalaemic IHCA ($n = 109$; mean serum $K^+ 7.8 \text{ mmol L}^{-1}$) used IV calcium and sodium bicarbonate ⁷⁶ in a very heterogeneous population and achieved 36.7 % ROSC, but only 12.8 % survived $> 24 \text{ h}$ and 3.7 % survived to hospital discharge. Given the poor outcome in patients with extreme hyperkalaemia, the authors proposed that dialysis may be an option.

Dialysis is the definitive treatment for hyperkalaemia, but is rarely used in cardiac arrest. A review of case reports of dialysis during CPR demonstrate successful outcomes in patients with a serum K^+ ranging between $8.2\text{--}10.2 \text{ mmol L}^{-1}$.⁵⁰ All dialysis modalities (with and without ECMO) have been used during CPR, demonstrating that it is technically feasible.^{78–81} The mean duration of dialysis to achieve ROSC was 45.4 min (range 15–95 min) and the mean reduction in serum K^+ was 3.2 mmol L^{-1} which would be difficult to achieve with medications alone, particularly during cardiac arrest.⁵⁰ The severity of hyperkalaemia appears to be a good indicator of the likelihood of achieving and sustaining ROSC. A protocol for initiation of dialysis during CPR has been reported.⁵⁰

A study assessing the outcome of OHCA following anti-hyperkalaemic medical treatment ($n = 465$; mean $K^+ 6.9 \text{ mmol L}^{-1}$, IQR $5.7\text{--}8.4 \text{ mmol L}^{-1}$) found no effect on ROSC, despite achieving K^+ -lowering (mean $K^+ 5.4 \text{ mmol L}^{-1}$, IQR $4.4\text{--}6.8 \text{ mmol L}^{-1}$).⁸² There is currently insufficient evidence for or against the use of IV calcium in the treatment of hyperkalaemia, therefore the ERC continues to recommend the administration of IV calcium to patients with hyperkalaemic cardiac arrest in all settings.

Hypokalaemia

Hypokalaemia is defined as a serum $K^+ < 3.5 \text{ mmol L}^{-1}$ [mild ($K^+ 3.0\text{--}3.4 \text{ mmol L}^{-1}$), moderate ($K^+ 2.5\text{--}2.9 \text{ mmol L}^{-1}$) or severe ($K^+ < 2.5 \text{ mmol L}^{-1}$)].

<2.5 mmol L⁻¹ or symptomatic].⁸³ It is usually caused by excessive K⁺ loss, transmembrane shift of K⁺ into cells, or reduced K⁺ intake.⁸⁴ Hypokalaemia is associated with higher in-hospital mortality and an increased risk of ventricular arrhythmias. Risks are increased in patients with pre-existing heart disease,^{85,86} those treated with digoxin⁸⁷ or those undergoing primary percutaneous coronary intervention for ST-elevation myocardial infarction.⁸⁸ In acute heart failure, hypokalaemia may be associated with increased short and long-term all-cause mortality after hospital discharge.⁸⁹

Treatment of hypokalaemia

Treatment is guided by the severity of hypokalaemia and presence of symptoms and/or ECG abnormalities, as illustrated in Fig. 3. Slow replacement of potassium is preferable, but in an emergency, rapid IV replacement, preferable via central route is required. The target

level is 4 mmol L⁻¹ K⁺.⁹⁰ Correction of any concomitant hypomagnesaemia is essential. Concomitant magnesium deficiency is common in patients with hypokalaemia. Repletion of magnesium will facilitate more rapid correction of hypokalaemia.⁹¹ Seek expert advice for potassium replacement in patients with severe renal impairment (see Fig. 3).

Calcium and magnesium disorders

The recognition and treatment of calcium and magnesium disorders is summarised in Table 2.

Hyperthermia and malignant hyperthermia

Hyperthermia results from a body temperature above normal (core temperature 36.5–37.5 °C) because of failed thermoregulation. Extremes of age and multimorbidity are specific risk factors.^{92,93}

Table 2 – Calcium and magnesium disorders with associated clinical presentation, ECG manifestations and recommended treatment.

Disorder	Causes	Presentation	ECG	Treatment
Hypercalcaemia Calcium > 2.6 mmol L ⁻¹	Primary or tertiary hyperparathyroidism Malignancy Sarcoidosis Medications	Confusion Weakness Abdominal pain Hypotension Arrhythmias Cardiac arrest	Short QT interval Prolonged QRS interval Flat T waves AV block Cardiac arrest	Guided by underlying cause Fluid replacement IV Furosemide 1 mg kg ⁻¹ IV Hydrocortisone 200–300 mg IV Pamidronate 30–90 mg IV
Hypocalcaemia Calcium < 2.1 mmol L ⁻¹	Chronic kidney disease Acute pancreatitis Calcium channel blocker overdose Toxic shock syndrome Rhabdomyolysis Tumour lysis syndrome Massive blood transfusion	Paraesthesia Tetany Seizures AV-block Cardiac arrest	Prolonged QT interval T wave inversion Heart block Cardiac arrest	Bolus: 10–20 mL 10 % Calcium gluconate over 5–10 min Infusion: 100 mL 10 % Calcium gluconate in 1000 mL 0.9 % Saline or 5 % Glucose at 50 mL h ⁻¹ IVI (monitor Ca ²⁺ level and adjust rate) 50 % Magnesium sulphate 4–8 mmol IV (if necessary)
Hypermagnesaemia Magnesium > 1.1 mmol L ⁻¹	Renal failure Iatrogenic	Confusion Flaccid paralysis Respiratory depression Hypotension AV-block Cardiac arrest	Prolonged PR and QT intervals T wave peaking AV block Cardiac arrest	Consider treatment when magnesium > 1.75 mmol L ⁻¹ : 10 % Calcium gluconate (10–30 mL) or 10 % Calcium chloride (5–10 mL) IV repeated if necessary Saline diuresis – 150 mL 0.9 % saline/ hr IV and Furosemide 1 mg/ kg IV Haemodialysis – when renal function impaired or severe symptoms, but risk of causing hypocalcaemia. Ventilatory support if necessary
Hypomagnesaemia Magnesium < 0.6 mmol L ⁻¹	GI loss Polyuria Starvation Alcoholism Malabsorption	Tremor Ataxia Nystagmus Seizures Arrhythmias – torsade de pointes Cardiac arrest	Prolonged PR and QT intervals ST-segment depression T-wave inversion Flattened P waves Increased QRS duration Torsades de pointes	Severe or symptomatic: 50 % Magnesium sulphate 2 g (8 mmol) IV over 15 min Torsades de pointes: 50 % magnesium sulphate 2 g (8 mmol) IV over 1–2 min Seizures: 50 % magnesium sulphate 2 g (8 mmol) IV over 10 min

Malignant hyperthermia is a rare pharmacogenetic disorder of skeletal muscle calcium homeostasis characterised by muscle contracture and life-threatening hypermetabolic crisis following exposure of genetically predisposed individuals to halogenated anaesthetics, succinylcholine, or neuroleptics.^{94,95} Rarely, malignant hyperthermia can be triggered non-pharmacologically. The most common clinical signs and symptoms include hypercapnia (34 %), sinus tachycardia (25 %), hyperthermia (20 %), masseter spasm (11 %), generalised muscle rigidity (3 %), acidosis, hyperkalaemia and death if left untreated.⁹⁶ This section is based on two systematic reviews, five non-systematic reviews, and two scoping reviews the latter being most recently performed on June 15, 2024.^{94–101}

Hyperthermia

Environmental-associated hyperthermia (core temperature > 38 °C) can be avoided by acclimatisation, adequate hydration and avoidance of physical activity in hot weather.^{92,98,102} The major risk factor is dehydration, which risks progression to heat syncope, heat exhaustion, heat stroke, and finally multiple organ dysfunction and cardiac arrest. Treatment of hyperthermia see Fig. 4 and Table 3. Core temperature should be measured centrally (e.g. tympanic, rectal, oesophageal) to guide treatment.¹⁰³

Malignant hyperthermia

Prevention of malignant hyperthermia is key; it is a genetic illness that if untreated results in death.^{97,108,109} Drugs such as 3,4-methylenedioxymethamphetamine (MDMA, ‘ecstasy’) and amphetamines may also cause a condition similar to MH and the use of dantrolene may also be therapeutic in these cases.¹¹⁰ Always consider malignant hyperthermia, with unexplained, unexpected increases in ETCO_2 , heart rate or temperature. Further information on malignant hyperthermia management (See QR code. 1).^{94,111}

Briefly, stop exposure, provide high flow oxygen, give dantrolene until $\text{ETCO}_2 < 45$ mmHg (6 kPa) with normal minute ventilation and core temperature < 38.5 °C, cool the patient. If cardiac arrest occurs, follow the universal ALS algorithm and continue to cool the patient. Apply the same cooling techniques as for post-resuscitation care targeted temperature management (see Fig. 5).

It is essential to contact an expert malignant hyperthermia centre for ongoing advice.^{97–112} Dantrolene should be stored centrally where anaesthesia is provided, and algorithms for MH management should be readily available (see QR code 1).

Toxin-Induced hyperthermia

Intoxication may present with hyperthermia as an early sign. Risk is higher in people taking psychotropic drugs (for treatment or recreation), sympathomimetics, anticholinergics, salicylates, weight-loss drugs or following ingestion of wild mushrooms.

Carry out an initial ABCDE approach after applying appropriate personal protective equipment (PPE) as in all cases of unknown toxic agents. Specific symptoms and the patient’s history may help to identify the suspected substance, even before toxicology results are available. For most poison-induced hyperthermia cases, prompt symptomatic treatment is key. Active cooling and targeted strategies for specific symptoms: benzodiazepines to reduce agitation and tremor, dantrolene for muscle contraction and rigidity, and measures to reduce toxin concentrations (See Table 4).

Accidental hypothermia and avalanche rescue

Accidental hypothermia is defined as a decrease in core temperature < 35 °C, during which vital signs and consciousness fade and finally disappear.¹⁵⁰ However, vital signs may be minimal but still be present with temperatures < 24 °C.

In hypothermic patients with spontaneous circulation, key interventions are insulation, and hospital triage followed by transfer and rewarming. In hypothermic patients with cardiac arrest, key interventions are continuous CPR and extracorporeal life support rewarming. These measures may result in good neurological outcome even with prolonged no-flow or low-flow (i.e. CPR) times, if hypothermia (e.g. < 30 °C) ensued before cardiac arrest. A scoping review up to February 13th 2025 found three systematic reviews for this recommendation,^{151–153} but also included other relevant articles.^{150,154–161}

Prevention from cardiac arrest

Primary accidental hypothermia is induced by exposure to cold, while secondary accidental hypothermia is triggered by illness and other external causes. Primary hypothermia is common in outdoor activities (mainly athletes and lost persons) and urban environments (e.g. homeless and intoxicated persons), while the incidence of secondary hypothermia is increasing among old and multimorbid persons in the indoor environment.^{154,157}

Measuring the core temperature with a low reading thermometer is the gold standard for diagnosis of hypothermia¹⁵⁷:

- tympanic in spontaneously breathing patients and
- oesophageal in patients having a tracheal tube or supraglottic airway in situ.¹⁵⁹

If the core temperature cannot be measured, temperature should be assessed according to the level of consciousness with the Revised Swiss Staging System.¹⁵⁶ (Table 5 and Fig. 12).

- In the Revised Swiss System, “Alert” corresponds to a GCS score of 15; “Verbal” corresponds to a GCS score of 9–14, including confused patients; “Painful” and “Unconscious” correspond to a GCS score < 9. While shivering is not used as a stage-defining sign in the Revised Swiss System, its presence usually means that the temperature is > 30 °C, a temperature at which hypothermic cardiac arrest is unlikely to occur.
- No apparent respiration, no apparent palpable carotid or femoral pulse, no measurable blood pressure – check for signs of life (pulse and, especially, respiration) for up to 1 min.
- The transition of colours between stages represents the overlap of patients within groups. The estimated risk of cardiac arrest is based on accidental hypothermia being the only cause of the clinical findings (Fig. 12). If other conditions impair consciousness, such as asphyxia, intoxication, high altitude cerebral oedema or trauma, the Revised Swiss System may falsely predict a higher risk of cardiac arrest due to hypothermia. Caution should be taken if a patient remains “alert” or “verbal” showing signs of haemodynamic or respiratory instability such as bradycardia, bradypnea, or hypotension because this may suggest transition to a stage with higher risk of cardiac arrest.

Hypothermic patients should be protected from the cold environment through minimal exposure and insulation and be transferred as fast as possible to the next appropriate hospital for rewarming. While not harmful, active rewarming is impractical during brief transport

Table 3 – Treatment of hyperthermia.

Degree of hyperthermia	Symptoms	Treatment
Mild – Heat syncope	Transient loss of consciousness and fast return to normal neurologic baseline.	Remove patient to a cool environment, passive cooling, resting and administration of oral isotonic or hypertonic fluids (the latter only if $\text{Na}^+ \leq 130 \text{ mmol L}^{-1}$).
Moderate – Heat exhaustion	Intense thirst, weakness, discomfort, anxiety, dizziness, syncope. Caused by mild to moderate hyperthermia ($>40^\circ\text{C}$) due to exposure to high environmental heat or excessive exercise.	Additionally, lie patient flat and administer isotonic IV fluids. Simple external cooling measures are usually not required but may involve conductive (e.g. cold floor, ice sheets; commercial ice packs to hands, feet and cheeks), convective (cold water immersion, cold shower) and evaporative measures (spraying cold water, fanning bare skin).
Severe – Heat stroke	Triad of severe hyperthermia (core temperature $> 40^\circ\text{C}$), neurological symptoms and recent passive environmental exposure (classic or passive heat stroke) or excessive exercise (exertional heat stroke). Symptoms include central nervous system dysregulation (e.g. altered mental state, seizure, coma), tachycardia, tachypnoea and arterial hypotension. ⁹² Mortality is approximately 10 %, and when combined with hypotension approaches 33 %. ⁹⁸ The outcome worsens if the core temperature is sustained at $> 40.5^\circ\text{C}$.	Rapidly cool the patient to $< 39^\circ\text{C}$, preferably $< 38.5\text{--}38.0^\circ\text{C}$ as quickly as possible. ⁹⁸ Prioritise active cooling methods over passive cooling – a cooling rate $\geq 0.155^\circ\text{C min}^{-1}$ is recommended ¹⁰⁴ . It is important to account for the time between symptom onset and the provision of care when selecting a cooling method, reaching the target temperature within 30 min of onset of heat stroke should be the goal ¹⁰⁴ . For exertional heatstroke, a cooling rate faster than $0.10^\circ\text{C min}^{-1}$ is safe and desirable. Ice or cold-water immersion (from neck down) or full body conductive cooling should be used, cooling rates of $0.2\text{--}0.35^\circ\text{C min}^{-1}$ can be achieved. ^{98,639} Cold water immersion should be continued until the symptoms have resolved or for a reasonable amount of time, e.g. 15 min, because benefit outweighs risk (weak recommendation, very low certainty evidence). ¹⁰⁵ if cold water immersion is not available, a combination of simple cooling techniques may be used, including conductive, convective and evaporative measures, although there are no comparative studies to guide the best option. ¹⁰⁵ A systematic review concluded that water immersion ($1\text{--}26^\circ\text{C}$ water) lowers body temperature faster and more effectively compared to passive cooling (low to very low certainty of evidence). Misting and fanning cooling techniques are marginally faster than passive cooling, and cold showers (20.8°C) cool faster than passive cooling. ¹⁰⁵ IV isotonic or hypertonic fluids should be administered (if $\text{Na}^+ \leq 130 \text{ mmol L}^{-1}$, for example up to $3 \times 100 \text{ mL } 3\% \text{ NaCl}$ at 10 min intervals). ¹⁰⁶ Additional electrolyte replacement with isotonic fluids should be considered and substantial amounts of fluids may be required. Follow the ABCDE approach in any patient with deteriorating vital signs. Critically ill patients will require aggressive and extended treatment in an intensive care unit. ^{92,107} There may be a requirement for advanced cooling techniques including external or internal devices used for targeted temperature management. There are no specific medications lowering core temperature.

Treatment is progressive in accordance with the degree of hyperthermia.

periods (e.g., less than 1 h).¹⁵³ Hypothermic patients with prehospital cardiac instability (i.e. heart rate $< 45/\text{min}$, systolic blood pressure $< 90 \text{ mmHg}$, ventricular arrhythmia, core temperature $< 30^\circ\text{C}$) should be rewarmed in-hospital using minimally invasive techniques with timely consultation with an ECMO centre being essential.^{156,161}

Management of cardiac arrest

The lowest published temperature from which successful resuscitation and rewarming has been achieved is currently 11.8°C ¹⁵⁵ for accidental hypothermia and 4.0°C for induced hypothermia.¹⁶² A systematic review reported only five patients (28–75 years of age) who had arrested at a body core temperature $> 28^\circ\text{C}$, suggesting

that cardiac arrest caused by primary hypothermia at $\geq 28^\circ\text{C}$ is possible, but unlikely.¹⁵¹ Some patients may still have minimal vital signs at a core temperature $< 24^\circ\text{C}$.¹⁶³

Check for signs of life for one minute – not only by clinical examination but also using ECG and ultrasound.¹⁶⁴

In hypothermic cardiac arrest, outcome relevant information should be collected to estimate the survival probability from hypothermic cardiac arrest with the HOPE (Hypothermia Outcome Prediction after ECLS rewarming for hypothermic arrested patients) –score.^{158,160,165}

The prognosis of patients with primary hypothermic cardiac arrest can be excellent,¹⁵⁷ while in cases of secondary hypothermia outcome is more influenced by co-morbidity.¹⁶⁶ Hypothermic patients

Table 4 – Toxins inducing hyperthermia.

Category	Substance	Mechanism	Symptoms	Diagnostic testing	Specific Management
Antipsychotics ^{113–117}	Risperidone, Aripiprazole, Haloperidol, Olanzapine, Quetiapine, Clozapine, Blonanserin	Dopamine antagonism/acute withdrawal of dopamine agonists → neuroleptic malignant syndrome (NMS)	Hyperthermia, muscle rigidity, tremor, autonomic dysfunction, altered mental status	Blood testing for antipsychotics	Benzodiazepines against agitation. Bromocriptine for hypodopaminergic state. Dantrolene as muscle relaxant
Antidepressants ^{118–122}	Lithium, MAOI, SSRI, SNRI, TCA	Serotonin syndrome	Hyperthermia, flushing, shivering, akathisia, agitation, mydriasis, autonomic dysfunction	Blood testing for antidepressants, serum lithium level	Benzodiazepines against agitation. Consider Chlorpromazine. Consider sodium bicarbonate for TCA
Recreative/Party Drugs ^{123–137}	MDMA (Ecstasy), LSD	Central catecholamine release and reuptake inhibition in the CNS, hypermetabolic condition with skeletal muscle stimulation, tachycardia and vasoconstriction	Hyperthermia, euphoria, hallucinations, agitation, shivering, mydriasis, nausea	Urine screening (MDMA, LSD)	Benzodiazepines for agitation. Dantrolene for hyperthermia. Consider carvedilol for MDMA
	Methamphetamine		Hyperthermia, hallucinations, tremor, agitation, mydriasis Autonomic dysfunction	Urine or blood screening for amphetamines	Ammonium chloride for excretion (urine acidifier). Activated charcoal to reduce absorption
	Cocaine	Nonselective dopamine, serotonin, noradrenaline reuptake inhibitor. CNS stimulation, sympathetic system activation. Direct sodium channel blockade. Coronary vasoconstriction	Hyperthermia, hallucinations, tremors, agitation, mydriasis, autonomic dysfunction, arrhythmias, chest pain	Urine and blood screening	Consider sodium bicarbonate for broad complex tachycardia or cardiac arrest, avoid β -blockers
Anticholinergics ^{138–142}	Atropine, scopolamine, plant alkaloids (belladonna, brugmansia, amanita)	Blockade of muscarinic receptors. Anticholinergic hyperthermic syndrome	Hyperthermia, tachycardia, sweating Inhibitor, dry skin & mucus membranes, flushing, mydriasis, altered mental status	Urine and serum screening	Benzodiazepines and central acting cholinesterase inhibitors against agitation. Activated charcoal within 1st hour (time window may be extended as anticholinergics reduce gastrointestinal motility)
Sympathomimetics ^{143–145}	Ephedrine, pseudoephedrine	Sympathetic system activation Increased metabolic rate and peripheral vasoconstriction	Hyperthermia, tachycardia, hypertension, arrhythmias, muscle twitching, nausea, mydriasis, urinary retention	Urine screening, serum toxicology testing for pseudoephedrine/ephedrine	Benzodiazepines against agitation
Salicylates ^{146,147}	Aspirin, methyl salicylate	Uncoupling of oxidative phosphorylation leading to increased heat production.	Mild hyperthermia, nausea, tachypnea, confusion, tinnitus	Serum salicylate levels	Urine alkalization with NaHCO_3 . Prevent hypoglycemia. Consider dialysis for drug excretion and cooling
Miscellaneous ^{148,149}	Dinitrophenol (DNP)	Uncoupling of oxidative phosphorylation leading to increased heat production	Hyperthermia, sweating, flushing, agitation, tachycardia & tachypnea, fatigue	Serum toxicology testing for DNP	Consider dantrolene

CNS denotes central nervous system, LSD denotes lysergic acid diethylamide, MAOI – Monoaminoxidase inhibitor, MDMA – 3,4-Methylenedioxymethamphetamine, SNRI – Serotonin and norepinephrine reuptake inhibitor, SSRI – Selective serotonin reuptake inhibitor, TCA – Tricyclic antidepressant.

in witnessed and unwitnessed cardiac arrest have good chances of neurological recovery if hypothermia developed before hypoxia and cardiac arrest, and if the chain of survival functioned well.^{151,152,166} Hypothermia diminishes the oxygen demand of the body (6–7 % per 1 °C cooling) and thereby protects the most oxygen dependent organs of the body (brain and heart) against hypoxic damage.¹⁶⁷ A systematic review of witnessed hypothermic cardiac arrest patients ($n = 214$) reported a survival to hospital-discharge rate of 73 %, with 89 % of survivors having a favourable neurologic outcome. A systematic review of unwitnessed hypothermic cardiac arrest patients ($n = 221$) reported a survival rate of 27 %, with 83 % of survivors having a favourable neurological outcome. Of note, the first rhythm was asystole in 48 % of these survivors.¹⁵²

Hypothermic cardiac arrest patients should receive continuous CPR until circulation has been re-established. Chest compression and ventilation rate should follow the standard ALS algorithm as

for normothermic patients. Efforts should also focus on maintaining normoxia during resuscitation.

Hypothermic cardiac arrest is often refractory to defibrillation and adrenaline. Defibrillation attempts have been successful in patients with a core temperature > 24 °C, however, ROSC tends to be unstable with lower temperature.¹⁶⁸ The hypothermic heart may be unresponsive to cardioactive drugs, attempted electrical pacing and defibrillation. If ventricular fibrillation or pulseless ventricular tachycardia persist after three shocks, it is reasonable to delay further attempts until the core temperature is > 30 °C.

Medication metabolism is slowed, leading to potentially toxic plasma concentrations, therefore, medications should be administered cautiously. The evidence for the efficacy of medication in severe hypothermia is limited and based mainly on animal studies. In severe hypothermic cardiac arrest, the effectiveness of amiodarone is reduced.¹⁶⁹ Adrenaline may be effective in increasing coronary perfusion pressure, but not survival.^{170,171} Adrenaline

Table 5 – Swiss staging of accidental hypothermia.¹⁵⁷

Stage	Clinical Findings	Core temperature (°C) (if available)
Hypothermia I (mild)	Conscious*	35–32 °C
Hypothermia II (moderate)	Impaired consciousness*	<32–28 °C
Hypothermia III (severe)	Unconscious*; vital signs present	<28 °C
Hypothermia IV (severe)	Apparent death; Vital signs absent	Variable**

* Consciousness may be impaired by comorbid illness (e.g. trauma, central nervous system pathology, toxic ingestion) or medications (e.g. sedatives, muscle relaxants, opioids) independent of core temperature.

** The risk of cardiac arrest increases < 30 °C, older and sicker patients. Alternative causes should be considered. Some patients still have vital signs < 24 °C.

	Hypothermia I	Hypothermia II	Hypothermia III	Hypothermia IV
Clinical findings	"Alert" from AVPU	"Verbal" from AVPU	"Painful" or "Unconscious" from AVPU Vital signs present	"Unconscious" from AVPU AND No detectable vital signs
Risk of cardiac arrest	Low	Moderate	High	Hypothermic cardiac arrest

Fig. 12 – Revised Swiss System for staging of accidental hypothermia.¹⁵⁶

AVPU – alert verbal pain unresponsive.

may induce myocardial injury and impaired neurologic recovery; it is reasonable to withhold adrenaline and other CPR drugs until the patient has been warmed to a core temperature ≥ 30 °C. If access to extracorporeal life support is delayed, since shorter CPR duration might be associated with better outcome, it may be reasonable to administer 1 mg of adrenaline even at lower core temperature in an attempt to achieve ROSC.¹⁷² Once 30 °C has been reached, the intervals between drug doses should be doubled when compared with those during normothermia (i.e. adrenaline every 6–10 min). Once normothermia is achieved (≥ 35 °C), use standard drug protocols.

If possible, transfer arrested hypothermic patients (or those at risk of arrest), directly to a hospital able to provide extracorporeal life support. If prolonged transport is required or the terrain is difficult, mechanical CPR is suggested. In hypothermic arrested patients with a body temperature < 28 °C delayed CPR may be used when CPR is too dangerous, and intermittent CPR can be used when continuous CPR is not possible, for example because of technically difficult rescue (Fig. 13).¹⁷³

Establish extracorporeal life support only in cardiac arrest or deteriorating patients (e.g. decreasing blood pressure, increasing acidosis). Primary ECLS may be considered in patients with ETCO_2 < 10 mmHg (1.3 kPa) or a systolic blood pressure ≤ 60 mmHg.¹⁷⁴ An unwitnessed cardiac arrest with asystole as first rhythm is not a contraindication for ECLS rewarming.¹⁵² Rewarming should preferably be performed with VA-ECMO rather than cardiopulmonary

bypass.^{175,176} If ECLS is not available within 6 h, non-extracorporeal life support rewarming may be used.^{177,178}

In-hospital prognostication of successful rewarming should be based on the HOPE score.^{158,160} The 5A score is a screening tool for predicting in-hospital mortality among elderly patients with accidental hypothermia with or without cardiac arrest which may also guide treatment options.^{179,180} In hypothermic arrested patients, criteria for ECLS rewarming should not be based on guidelines for normothermic cardiac arrest.¹⁸¹ This may result in potentially life-saving treatment being withheld in patients with a potential for good neurological outcome.¹⁷⁶

Emergency medical services and hospitals should install structured protocols to improve prehospital triage, transport and treatment, as well as in-hospital management of hypothermic patients.

Avalanche rescue

The chances of surviving avalanche burial is steadily improving, due to collaborative efforts to improve avalanche search and rescue, and subsequent medical interventions.¹⁸² Most avalanche victims die from asphyxia, fewer from trauma or hypothermia. In cases of unwitnessed cardiac arrest presenting in asystole, avalanche victims have a poor chance of survival, even if guidelines are followed.^{183–185}

Several factors improve the likelihood of good outcome from avalanche burial. These include non-critical burial (i.e. head and chest out of the snow), superficial burial, burial for a short time, burial dur-




Fig. 13 – Decision making in hypothermic cardiac arrest.

CPR – cardiopulmonary resuscitation. Delayed and intermittent CPR in hypothermic patients when continuous CPR is not possible during dangerous or difficult rescue.¹⁷⁰

ing daylight hours, the presence of an air pocket (i.e. open airways in addition to any space in the snow in front of nose and mouth), retrieval before cardiac arrest, witnessed cardiac arrest and ROSC in the first few minutes of CPR.^{182–184,186,187} The International Commission for Mountain Emergency Medicine performed a scoping review and published guidelines for the management of victims buried in avalanches (Fig. 14).¹⁸⁸

The AvaLife algorithm should be used in multiple burial scenarios with too few rescuers on site. In this triage situation, AvaLife enables identification of the i) buried subjects who should be excavated first (cut-off < 1.5 m of burial depth) and ii) who should be resuscitated and for how long.¹⁸⁹ AvaLife has been specifically developed for BLS provider.

The quality of CPR can be compromised by several factors, including the confined space of the burial site, chest compressions on snow, hypobaric hypoxia on mountains leading to faster exhaustion of rescuers, and long and difficult extrication and transport of victims. Mechanical chest compression devices can be helpful in technically difficult and prolonged rescues.^{190–192}

Thrombosis

Pulmonary embolism

Cardiac arrest from acute pulmonary embolism (PE) is the most serious clinical presentation of venous thromboembolism, in most cases originating from a deep venous thrombosis (DVT).¹⁹³ The reported incidence of cardiac arrest caused by PE is 2–7 % of all OHCA, ^{194,195} and 5–6 % of all in-hospital cardiac arrests,^{75,196} but it is likely to be underestimated. Overall survival is low, even if invasive treatment or ECPR is used.^{195,197,198} Specific treatments for cardiac arrest resulting from PE include administration of fibrinolytics, surgical embolectomy and percutaneous mechanical thrombectomy.

The 2020 ILCOR systematic review explored the influence of specific treatments (e.g. fibrinolytics, or any other) yielding favour-

able outcomes.¹⁹⁹ The 2019 ILCOR summary statement reviewed the use ECPR for cardiac arrest in adults,²⁰⁰ while the 2022 ILCOR summary statement updated evidence on sensitivity and specificity of POCUS for specific pathophysiological states, including PE, and reviewed four small additional observational studies on cardiac arrest from PE which were published since the previous review. No additional evidence was found to change the management.²⁰¹ Additional evidence was identified from the European Society of Cardiology (ESC) guideline on pulmonary embolism.¹⁹³

These ESC Guidelines define ‘confirmed pulmonary embolism’ as a probability of pulmonary embolism high enough to indicate the need for specific treatment.¹⁹³ Clinical history, assessment, capnography, and echocardiography (if available) can assist in the diagnosis of acute pulmonary embolism during CPR. Cardiac arrest commonly presents as PEA.¹⁹⁷ Constant low ET_{CO}₂ readings (below 1.7 kPa (13 mmHg)) while performing high-quality chest compressions may support a diagnosis of pulmonary embolism, although it is a non-specific sign.^{202,203} If a 12-lead ECG can be obtained before cardiac arrest, changes indicative of right ventricular strain may be helpful for decision making.

Common symptoms preceding cardiac arrest are described in Table 6.

Acute pulmonary embolism can cause right ventricular pressure overload and dysfunction, POCUS might be helpful for detection, (Table 6). but no individual echocardiographic parameter provides fast and reliable information on right ventricle size or function. Signs of right ventricular overload or dysfunction may also be caused by other cardiac or pulmonary diseases.²⁰⁴

Initial treatment (Fig. 15)

All patients with a sudden onset of progressive dyspnoea, especially in patients without pre-existing cardiac or pulmonary disease are suspicious for pulmonary embolism. Hypoxaemia is usually reversed with administration of oxygen, while, in some patients, correction of




Fig. 14 – Initial management of critically buried avalanche victim.¹⁸⁸

CPR – cardiopulmonary resuscitation, ECG – electrocardiography, ECLS – extracorporeal life support. Figure legend. ^A Assess for lethal injuries: decapitation; truncal transection; whole body decomposed. If present, do not start CPR. ^B Signs of life include any of the following: A, V or P from AVPU (alert, responsive to verbal stimuli, responsive to pain, unresponsive) or Glasgow Coma Scale > 3, any visible movement, respirations, or a palpable carotid or femoral pulse (for experienced ALS providers). ^C Standard compression / ventilation rates. Drug dose and defibrillation depending on core temperature or, if not available, burial duration. If ventricular fibrillation persists after three shocks, delay further attempts until the core temperature is > 30 °C.

hypoxaemia will not be possible without simultaneous pulmonary reperfusion.¹⁹³

High-risk pulmonary embolism is suspected with shock or persistent arterial hypotension and is immediately life-threatening. Leave hypotensive patients lying flat if breathing allows, to prevent further progression of hypotension and cardiac arrest. Acute right ventricular failure is the leading cause of death in patients with high-risk pulmonary embolism. Aggressive volume expansion is of no benefit and may even worsen right ventricular function. However, cautious volume loading (≤ 500 mL over 15–30 min) may increase cardiac output, and the use of vasopressors and/or inotropes is frequently beneficial.¹⁹³

IV anticoagulation should be initiated while awaiting the results of diagnostic tests. Unfractionated heparin is recommended for patients with shock and hypotension, and for whom primary reperfusion is considered. Thrombolytic treatment of acute pulmonary embolism restores pulmonary perfusion more rapidly than anticoagulation with heparin alone.²⁰⁶

Modifications to ALS

Fibrinolysis. If pulmonary embolism is the suspected cause of cardiac arrest, fibrinolytic drugs should be administered, based on evidence from an ILCOR CoSTR.²⁰¹ There is insufficient evidence

Table 6 – Non-specific patient characteristics observed with higher incidence in OHCA caused by pulmonary embolism compared with other causes^{193,198,205}.

Symptoms preceding cardiac arrest

Sudden onset of dyspnoea

Pleuritic or substernal chest pain

Cough

Haemoptysis

Syncope (otherwise unexplained and/or repeated collapses)

Signs of DVT (unilateral low extremity swelling)

Signs of right ventricular strain on 12-lead ECG (if obtained before cardiac arrest)

- Inversion of T waves in leads V1–V4
- QR pattern in V1
- S1 Q3 T3 pattern (i.e. a prominent S wave in lead I, a Q wave and inverted T wave in lead III)
- Incomplete or complete right bundle-branch block

Assessment during resuscitation

Female biological sex

Lower age

EMS witnessed cardiac arrest

PEA as first observed rhythm

Low ETCO₂ readings during high quality CPR (<1.7 kPa/13 mmHg)

Presence of right heart dilation with poor filling of the left heart, D sign (straightening of interventricular septum) and/or intracardiac thrombi (intra-arrest POCUS)

to recommend optimal drug and dosing strategy for fibrinolysis during CPR. ROSC and survival were observed after recombinant tissue type plasminogen activator (alteplase, bolus 50 mg IV with or without additional 50 mg after 30 min, or 0.6–1.0 mg/kg IV – max. 100 mg).^{197,207,208} When thrombolytic drugs have been administered, continue CPR for at least 60–90 min.^{195,209,210} Based on expert opinion, thrombolysis or surgical embolectomy should be considered for pregnant women with high-risk pulmonary embolism and PE causing cardiac arrest.¹⁹³

Surgical embolectomy or percutaneous mechanical thrombectomy.

Successful surgical embolectomy and percutaneous mechanical thrombectomy have been reported in cardiac arrest patients and are recommended if PE is the known cause of cardiac arrest, and the specialised skills are available.^{211–213} Treatment decisions should be made by a highly experienced interdisciplinary team, involving a thoracic surgeon or interventional cardiologist.^{193,214}

Extracorporeal CPR. Some observational studies suggest the use of ECPR if cardiac arrest is associated with pulmonary embolism.^{198,215} ECPR maintains circulation and gas exchange. Time to ECPR correlates with neurological outcome.²¹⁶ Consider ECPR as a rescue therapy for selected patients with cardiac arrest when conventional CPR is failing in settings in which it can be implemented.²⁰¹ It is recommended that adequate ECPR flow is established within 60 min of the onset of cardiac arrest.²¹⁷ Favourable neurological outcomes in patients with pulmonary embolism undergoing ECPR is inferior to outcomes related to other aetiology.^{198,218} Despite the extremely high mortality rate of patients with cardiac arrest refractory to standard ALS, ECPR is also an option to increase the pool of organ donors.¹⁹⁸

Coronary thrombosis

Coronary artery disease (CAD) remains the leading cause of OHCA in adults, either due to ventricular arrhythmias triggered by acute myocardial ischaemia, or those arising from the fibrotic arrhythmogenic substrate in patients with previous myocardial infarction.^{219,220} While CAD prevalence increases with age, it is also the most common cause of cardiac death in adults aged 35 to 50 years.^{221,222}

A recent systematic review and meta-analysis reported significant CAD in 75 % of OHCA cases,²²³ ranging from 88 % in patients with initial shockable rhythm and ST-elevation on ECG, to 54 % in non-shockable cases without ST-elevation. Culprit lesions were identified in nearly 60 %, more often in patients with shockable rhythm and ST-elevation, while acute coronary occlusion was found in around 40 %, with a higher prevalence in patients with ST-elevation. Among those with refractory cardiac arrest, 75 % had significant CAD and 70 % had a culprit lesion, often involving the left main coronary artery.

Prevention strategies for CAD as a cause of OHCA should comprise promoting healthy lifestyles in asymptomatic individuals, as well as addressing modifiable risk factors of atherosclerosis in patients with known disease, most effectively through cardiac rehabilitation.^{224,225}

Health education should target time reduction from symptom onset (i.e. chest pain) to seeking medical help, in order to enable early diagnosis and treatment. Behaviour change techniques (action planning, information about health consequences, signs and symptoms and instruction on what to do) might be helpful for this particular purpose.²²⁶

1. Prevent & be prepared



Promote cardiovascular health to reduce the risk of acute coronary events



Enhance health education to recognise symptoms & BLS training for likely rescuers



Strengthen regional STEMI networks to ensure timely PCI

2. Suspect & react



Evaluate 12-lead ECG after ROSC; repeat if inconclusive



Assess clinical data suggesting acute coronary syndrome



Activate the STEMI network & transfer the patient to PCI centre if ST-elevation or ongoing ischemia

3. Resuscitate & establish reperfusion strategy

Patients with sustained ROSC

ST-elevation

Is PCI possible within 120 min?

Yes

Immediate coronary angiography ± PCI

No

Consider fibrinolysis if not contraindicated & transfer to PCI centre

No ST-elevation

Is the patient unstable or shows signs of ongoing ischemia?

Yes

Immediate coronary angiography ± PCI

No

Consider ruling-out non-coronary causes

Consider delayed coronary angiography ± PCI

No Sustained ROSC

Does the OHCA setting suggest likely futility?

Yes

Consider stopping CPR

No

Consider transfer to PCI centre with ongoing CPR

↓
Consider mechanical CPR

↓
Consider ECPR

Fig. 15 – Coronary thrombosis algorithm.

BLS – basic life support, STEMI – ST elevation myocardial infarct, PCI – percutaneous coronary intervention, ECG – electrocardiography, OHCA – out-of-hospital cardiac arrest, ROSC – return of spontaneous circulation.

An updated 2022 ILCOR CoSTR concluded that BLS training consistently improved BLS skills, knowledge, and increased confidence to perform CPR. Thus, BLS training for likely bystanders of high-risk populations is recommended and should be actively promoted by healthcare professionals.²⁰¹

Along with initiatives raising public awareness, healthcare systems should establish regional STEMI networks to ensure equal and timely access to percutaneous coronary intervention (PCI).²²⁵ A recent systematic review and meta-analysis found that such networks may reduce STEMI case-fatality by 35 % and long-term mortality by 27 %.²²⁷ Integration of the emergency transport system was a critical factor for success.

Detect features suggesting coronary thrombosis and activate STEMI network

A 12-lead electrocardiogram (ECG) should be obtained and evaluated after ROSC to identify possible ischaemic features. Additional ECGs may help decision-making, as defibrillation and the time elapsed from ROSC to ECG acquisition may affect findings, with a higher percentage of false-positive for STEMI if the ECG is performed ≤ 8 min from ROSC.²²⁸ ST elevation remains the most sensitive and specific sign of coronary artery occlusion; however, its absence does not completely exclude the condition.²²⁹ Other ECG patterns—such as bundle branch block or diffuse ST-depression with concurrent ST-segment elevation in aVR and/or V1—may suggest coronary occlusion and should be considered if the clinical context is compatible.²²⁵ Additionally, non-cardiac conditions, such as subarachnoid haemorrhage, can also cause ST-changes,²³⁰ highlighting the importance of clinical correlation.

A comprehensive approach integrating ECG findings and clinical information suggesting a possible coronary cause—such as a history of CAD, chest pain before arrest, or an initial shockable rhythm—²³¹ is recommended. Once clinical suspicion is established, the STEMI network should be promptly activated to ensure early transfer to a centre with PCI capability.

Resuscitate according to ALS algorithm & establish reperfusion strategy

Patients with ROSC and ST-elevation. OHCA resuscitation should follow the standard ALS protocol (see ALS chapter). Conscious patients with persistent ST-elevation (or equivalent) after ROSC should undergo immediate coronary angiography and PCI within 120 min of diagnosis, as per standard STEMI management. If significant delays are expected, pre-hospital fibrinolysis may be considered unless CPR was prolonged or traumatic or there are other contraindications, followed by immediate transfer to PCI centre.²²⁵ Comatose patients with ST-elevation after ROSC have significantly worse prognosis, and no RCTs have directly compared early versus delayed angiography in this group. However, since early angiography is the standard of care for STEMI and no evidence suggests otherwise, both the ILCOR²³² and the European Society of Cardiology²²⁵ recommend the same approach for these patients.

Pharmacological therapy should align with STEMI protocols for non-arrest patients.²²⁵ In patients targeted for a primary PCI strategy, aspirin is recommended and early administration of a P2Y12 inhibitor (ticagrelor or prasugrel over clopidogrel, or IV cangrelor if oral intake is not possible) may be considered. Unfractionated heparin (IV) is the preferred anticoagulant, with enoxaparin (SC) or bivalirudin (IV) as alternatives.

Patients with ROSC and no ST-elevation. While previous observational studies suggested a potential benefit of early coronary angiography in resuscitated patients without ST-elevation on ECG,^{233,234} recent RCTs have not confirmed this. A 2021 ILCOR systematic review and meta-analysis²³⁵ compared routine early (2–6 h) versus delayed (within 24 h) coronary angiography in unconscious adult patients with ROSC after OHCA. This included data from the Direct or Subacute Coronary angiography in Out-of-hospital Cardiac Arrest-DISCO trial ($n = 79$),²³⁶ Coronary Angiography After Cardiac Arrest Trial-COACT ($n = 552$)²³⁷ and Randomized Pilot Clinical Trial of Early Coronary Angiography-PEARL ($n = 99$)²³⁸ – the latter being underpowered. No survival or neurological benefit was found for early versus delayed angiography in patients without ST-elevation. In 2022, an updated ILCOR CoSTR²³² added the results of the Angiography after Out-of-Hospital Cardiac Arrest without ST-Segment Elevation-TOMAHAWK ($n = 554$) trial,²³⁹ which also found no advantage of a routine early approach. Accordingly, the Task Force deemed both early and delayed angiography reasonable in patients without ST-elevation after ROSC (weak recommendation, low-certainty evidence).²³² Two latter RCTs, Emergency vs Delayed Coronary Angiogram in Survivors of Out-of-Hospital Cardiac Arrest-EMERGE ($n = 279$)²⁴⁰ and Coronary angiography in Out-of-hospital cardiac arrest Patients without ST-segment Elevation-COUPÉ ($n = 66$)²⁴¹ have reinforced these findings, despite variations in study protocols and outcome definitions, as well as limited statistical power.

Importantly, patients with shock, electrical instability, or signs of ongoing ischaemia were excluded or allowed to cross over to the emergency arm. Thus, current evidence does not support routine early over delayed angiography in stable patients without ST-elevation on the post-ROSC ECG. However, individual decisions should consider the patient's haemodynamic status and signs of ongoing ischaemia. Also, ruling out non-coronary causes is recommended if the clinical context suggests a potential alternative aetiology for the arrest.

Pharmacological treatment should generally follow the recommendations for non-ST-elevation acute coronary syndrome if suspected.²²⁵ Aspirin is recommended, but routine early use of P2Y12 inhibitors is discouraged if early angiography is planned. Unfractionated heparin (IV) and fondaparinux (SC) are the preferred anticoagulants for the early and delayed angiography strategies, respectively. However, if there are concerns about other non-coronary causes for the arrest, withholding antiplatelet and anticoagulant therapies until further evaluation is reasonable.

Patients without sustained ROSC. Mechanical CPR and, ultimately, ECPR might be the only therapeutic option for refractory OHCA. The first has not proved consistent superiority over manual CPR, but it may facilitate delivering high-quality chest compression during transportation or while performing coronary angiography.²⁴² Studies addressing ECPR in this setting have reported conflicting results. An updated ILCOR systematic review compared ECPR with manual or mechanical CPR for refractory OHCA, assessing survival and neurological outcomes.^{8,243} This included two single-centre RCTs: the ARREST ($n = 30$) trial was terminated early because of superiority in the ECPR arm for primary outcome (survival to hospital discharge). Interpretation of included studies was difficult due to different study designs and inconsistent findings.^{244,245} A multicentre

RCT found no difference in 30-day survival with favourable neurological outcomes.²⁴⁶ Given these inconclusive findings but considering the potential risk–benefit balance in this poor-prognosis scenario, ECPR may be considered as a rescue therapy for selected OHCA patients when conventional CPR fails to achieve ROSC, provided it can be implemented.²⁴³

Toxic agents

Intoxication is one of the eight reversible causes of cardiac arrest. Thus, all patients with cardiac arrest should be screened for signs of intoxication, especially in suspicious cases, unexpected cardiac arrests or in cases with more than one casualty. If intoxication is likely, resuscitation teams should avoid contamination by first donning appropriate personal protective equipment (PPE). Direct skin contact or mouth to mouth ventilation might transmit toxic agents and should therefore be avoided (See Fig. 7). The therapeutic strategy of intoxication consists of the three pillars:

- Decontamination
- Enhanced elimination
- Administration of antidotes

Helpful online databases on poison centres and toxins can be found in the QR codes 2 and 3). The patient's temperature should be measured because hypo- or hyperthermia may occur during drug overdose, as mentioned earlier. The 2021 ERC guidelines provide a detailed overview about intoxication.⁹⁹ These updated recommendations are based on recent scientific evidence, systematic reviews, and expert consensus. A separate subchapter addresses local anaesthetic systemic toxicity.



QR code. 2 – World directory of poisons centres – provided by the World Health Organization.



QR code. 3 – PubChem – world's largest collection of freely accessible chemical information – provided by the National Library of Medicine.

Intoxication with opioids

An ILCOR systematic review found heterogenous results on the benefit of opioid-specific advanced life support-level therapies for cardiac arrest.²⁴⁷ Existing evidence is not sufficient to recommend administration of an opioid-antagonist (e.g. naloxone) for cardiac arrest caused by opioid poisoning.

Intoxication with cardiac glycoside

A narrative review on treatment of patients with haemodynamic instability caused by cardiac glycoside poisoning found an improved haemodynamic status and survival after administration of digoxin immune-Fab fragments. Favourable outcomes were seen in patients receiving magnesium, cardioversion, or cardiac pacing.²⁴⁸

ECPR

ECPR seems to be associated with increased survival in intoxicated patients in refractory cardiogenic shock or cardiac arrest.²⁴⁹

Traumatic cardiac arrest

Traumatic cardiac arrest is associated with a very high mortality and good neurological outcome is reported in less than 50 % of those who survive. The response to traumatic cardiac arrest is time-critical and success depends on a well-established chain of survival, including advanced prehospital and specialised trauma centre care. Immediate resuscitative efforts in traumatic cardiac arrest focus on simultaneous treatment of reversible causes, which takes priority over chest compressions. Evidence is based on nine systematic reviews and evidence updates and consensus from the expert group.^{250–258}

In Europe, traumatic cardiac arrest accounts for 4 % of all cardiac arrests occurring in the pre-hospital setting.²⁵⁹ Registry data for survival range from 0 %²⁶⁰ to 37 %.²⁶¹ A recent systematic review²⁵⁵ of traumatic cardiac arrest identified 36 studies, and a total of 51,722 patients, reporting an overall mortality rate of 96.2 %, with 43.5 % of survivors achieving a favourable neurological outcome. The presence of a physician at the prehospital scene was associated with improved outcomes (6.1 % vs 2.4 % survival and 57.0 % vs 38.0 % favourable neurological outcome, with or without physician respectively). Another systematic review confirmed these findings.²⁵³

Factors relevant for prognostication are listed in Table 7.

Withholding resuscitation

The American College of Surgeons and the National Association of EMS Physicians recommend withholding resuscitation in situations where death is inevitable or established and in trauma patients presenting with apnoea, pulselessness and without organised ECG activity.²⁷¹ However, neurologically intact survivors initially presenting in this state have been reported.²⁷² The ERC based on expert consensus recommends the following approach:

Consider withholding resuscitation in traumatic cardiac arrest in any of the following conditions:

- no signs of life within the preceding 15 min.
 - massive trauma incompatible with survival (e.g. decapitation, extensive cardiac destruction, massive head injury with loss of brain tissue).
- We suggest termination of resuscitative efforts if there is:
- no ROSC within 20 min after reversible causes have been addressed.
 - no detectable ultrasonographic cardiac activity in PEA 20 min after reversible causes have been addressed.

Preventable deaths

For a diagnosis of TCA, there must be plausible signs of traumatic injury or, at least, a mechanism of injury consistent with trauma. In the absence of an identifiable mechanism or visible signs of injury, the standard ALS algorithm should be followed instead. The use of ultrasound supports findings in this context, aiding in differentiating *per-arrest* from TCA, identifying reversible causes, and guiding resuscitative efforts accordingly.²⁷³

A considerable proportion of trauma-related deaths can be attributed to management errors. In an urban German population, the

Table 7 – Prognostication in traumatic cardiac arrest.^{254,262–270.}

–	+
Age	Emergency surgery
Female	Major trauma centre care
Increased ISS	Shockable ECG rhythm
Head injury	Reactive pupils
Shock on admission	Respiratory activity
Need for blood transfusion	Spontaneous eye or extremity movements
CPR in the emergency department	Organised ECG rhythm
Children	Shorter CPR duration
	Reduced prehospital time
	Penetrating chest injury
	Witnessed arrest
	Cardiac motion in ultrasound

prevalence of potentially preventable prehospital deaths from trauma was reported as 15.1 %, ²⁷⁴ whereas in an urban U.S. population, this figure was higher at 29 %. ²⁷⁵ In Australia, the rate of potentially preventable trauma deaths was found to be 20 %. ²⁷⁶ Throughout these studies, exsanguination was identified as a leading cause of preventable death. Taking these data into account, the ERC recommends participation in an accredited trauma management training

program for those involved in the care of trauma patients. Ideally such a training format reflects the standard team approach provided in the European systems. ²⁷⁷

The reversible causes associated with TCA are uncontrolled haemorrhage (48 %); tension pneumothorax (13 %); asphyxia (13 %); and cardiac tamponade (10 %). The presenting rhythms found in traumatic cardiac arrest are usually PEA or asystole, depending on the time interval between circulatory arrest and the first ECG recording; PEA (66 %); asystole (30 %); VF (4 %). ²⁷⁸ The key pathophysiological mechanisms leading to traumatic cardiac arrest are listed in Table 8:

As these mechanisms progress beyond a critical threshold, irreversible circulatory failure ensues:

- Progressive bradycardia and hypotension (caused by severe hypoxia and acidosis)
- PEA with minimal cardiac activity on ultrasound (Pseudo-PEA) refers to a state where the heart exhibits weak contractile activity on ultrasound but fails to generate an effective pulse or circulation. ²⁶⁶
- PEA without with no cardiac activity detectable on ultrasound. The mortality of this condition approaches 100 %.
- Asystole → The final stage of cardiac arrest, with no electrical or mechanical activity.

Effectiveness of chest compressions

In cases of cardiac arrest due to hypovolaemia, cardiac tamponade, or tension pneumothorax, the effectiveness of chest compressions is uncertain. Their impact likely depends on ventricular filling, which is compromised in these conditions, and in *peri*-arrest states (pseudo-PEA), chest compressions may even reduce the remaining cardiac output by further decreasing venous return. ^{279–281} Therefore, chest compressions must not delay immediate treatment of reversible causes.

Following resuscitative thoracotomy, open cardiac compressions might be beneficial. Patients sustaining traumatic cardiac arrest within the first six hours of hospital admission may benefit from open

Table 8 – Major causes of traumatic cardiac arrest and their pathophysiology.

Cause	Mechanism	Impact on Circulation
Hypovolaemic shock	Severe blood loss → inadequate preload & cardiac output	Decrease in right ventricular filling → hypotension, decrease in cardiac output → tissue hypoperfusion including coronary perfusion, → tissue hypoxia → metabolic acidosis, shock, ECG typically PEA, followed by asystole
Tension pneumothorax	Increased intrathoracic pressure → impaired venous return	
Cardiac tamponade	Pericardial blood accumulation → restricted ventricular filling	
Hypoxia/asphyxia	Airway compromise, lung injury, or Brain Impact Apnoea	Progressive hypoxaemia → bradycardia → PEA → asystole
Metabolic acidosis (contributing factor)	Ischaemia from prolonged hypoxia & hypoperfusion	Cardiac dysfunction, reduced contractility
Neurogenic shock (contributing factor)	Loss of sympathetic tone	Aggravates other shock states, worsening hypotension

cardiac compression instead of closed chest compressions; in one retrospective trial, open cardiac compressions in traumatic cardiac arrest were associated with a higher long-term survival both in penetrating and blunt trauma. However, the study does not address the management of reversible causes.²⁸²

Providers without the training to manage reversible causes should follow the standard ALS algorithm and prioritise rapid transport under ALS or BLS, depending on system capability. Those with the necessary expertise should make case-by-case decisions based on available resources, likely reversible causes, and regional guidelines.

The role of inotropes and vasopressors

Although adrenaline may increase the rates of ROSC with medical cardiac arrest,²⁸³ the role of inotropes and vasopressors in traumatic cardiac arrest (TCA) is unclear.²⁸⁴ They may serve as a temporary bridge until reversible causes are addressed. Their use may also be considered in late-stage shock with cardiac dysfunction or mixed shock states such as trauma with a neurogenic component. The primary aim in traumatic cardiac arrest is to identify and treat any reversible cause which should take preference over inotrope/vasopressor therapy.

Pre-hospital care

The TCA algorithm (Fig. 8) applies not only to patients in traumatic cardiac arrest but also to *peri-arrest* trauma patients who present with signs of impending circulatory collapse. In both scenarios, the response must be immediate, targeted, and adapted to the capabilities of the prehospital system. The choice between a “stay and treat” strategy versus “scoop and run” or “treat-as-you-go” depends less on geography and more on the available skill set, equipment, and system infrastructure. In highly trained systems with extended capabilities, life-saving interventions may be feasible on scene. In others, immediate transfer may be the best chance for survival, provided that basic haemorrhage control and oxygenation are ensured during transport. Regardless of the system configuration, prolonged prehospital time must be avoided.²⁷³ Early pre-alerting of the Major Trauma Centre (MTC) is essential. In cases of ongoing haemorrhage or severe physiological compromise, the patient should be declared “Code Red” (or equivalent), enabling the hospital to activate the massive transfusion protocol, mobilise the trauma team, and ensure immediate access to surgical care upon arrival.

The recently published practice guidelines of the Royal College of Surgeons of Edinburgh on traumatic cardiac arrest provide a comprehensive overview of prehospital management.²⁸⁵

In-hospital care

Successful treatment of traumatic cardiac arrest requires a highly coordinated, team-based approach, where all life-saving interventions are carried out in parallel rather than sequentially, based on the clinical priorities dictated by the patient’s presentation. The focus is on rapid identification and treatment of all potentially reversible causes to maximise the chances of survival. A prospective observational study in patients with exsanguinating injuries and systolic blood pressure < 90 mmHg showed lower 30-day mortality (17.5 % vs. 72.0 %, $p < 0.001$) when prioritising circulation (CAB approach) over tracheal intubation (ABC approach), thus enabling for resuscitative bleeding control before ventilation.²⁸⁶ Fig. 8 illustrates the traumatic cardiac (peri-)arrest algorithm, which is based on the universal ALS

algorithm but integrates trauma-specific modifications to accommodate the unique challenges of traumatic cardiac arrest management. A multidisciplinary team approach, including trauma surgeons, emergency physicians, anaesthetists, radiologists, intensivists and nursing staff, is essential to ensure that each critical intervention is initiated without delay, according to the patient’s immediate needs.

Treatment of the reversible causes in traumatic cardiac arrest Hypovolaemia. The treatment of severe hypovolaemia has several components. The main principle is to achieve immediate haemostasis. Temporary haemorrhage control can be lifesaving.²⁷³ Compressible external haemorrhage can be treated with elevation (of a bleeding limb), direct or indirect pressure, pressure dressings, tourniquets and topical haemostatic agents.^{273,287} Non-compressible haemorrhage is more difficult to address and splints (pelvic splint), blood products, IV fluids and tranexamic acid can be used while transferring the patient for surgical haemorrhage control.²⁸⁸

Immediate aortic occlusion is recommended as a last resort measure in patients with exsanguinating and uncontrollable infra-diaphragmatic torso haemorrhage. This can be achieved through resuscitative thoracotomy and cross-clamping (or manual compression) of the descending aorta or resuscitative endovascular balloon occlusion of the aorta (REBOA). There is no evidence for one technique being superior to the other.²⁸⁹ Both are highly specialised interventions, appropriate expertise and equipment. A recent UK REBOA study²⁹⁰ in traumatic shock revealed a possible increase in mortality. The ERC does not recommend the use of REBOA outside of clinical trials.

In hypovolaemic traumatic cardiac arrest, immediate restoration of the circulating blood volume with blood products is mandatory. Prehospital transfusion of fresh plasma and packed red blood cells may improve survival, especially in cases with prolonged prehospital time.^{291,292} Recent evidence suggests additional benefit from whole blood transfusion.^{258,293}

Hypoxia. In traumatic cardiac arrest, hypoxaemia can be caused by airway obstruction, traumatic asphyxia or impact brain apnoea.²⁹⁴ Impact brain apnoea is an underestimated cause of morbidity and mortality in trauma, but not necessarily associated with an unsurvivable brain injury.²⁹⁵ Data from the National Trauma Audit and Research Network database in UK reported a 15 % survival rate with 90 % favourable outcome among traumatic cardiac arrest patients with brain impact apnoea.²⁹⁶ Brain impact apnoea may aggravate the course of traumatic brain injury and can lead to asphyxiation if left untreated. Effective airway management and ventilation with oxygen can prevent and reverse hypoxic cardiac arrest. In a recent national Dutch study, 52 % of traumatic cardiac arrest were attributed to traumatic brain injury-related apnoea.²⁹⁷

Controlled ventilation in circulatory compromised patients is associated with major risks because of the increase in intrathoracic pressure^{298,299} that may lead to further decreased cardiac output caused by:

- Impeded venous return to the heart, particularly in severe hypovolaemia.
- Reduced diastolic filling, particularly in cardiac tamponade
- Conversion of pneumothorax into a tension pneumothorax
- Increase in blood loss from venous bleeding sites

Minimising intrathoracic pressure using low tidal volumes and minimal PEEP may help optimise cardiac preload. Ventilation should

be monitored with capnography and adjusted to achieve normocapnia.^{265,300,301}

Tension pneumothorax. Tension pneumothorax is a reversible cause of cardiac arrest and must be excluded in traumatic cardiac arrest. It may lead to cardiac arrest by obstructing venous return through mediastinal shift and impairing effective gas exchange. Positive pressure ventilation can convert a simple pneumothorax into a tension pneumothorax.³⁰² The prevalence of tension pneumothorax is approximately 0.5 % in all major trauma patients treated in the pre-hospital setting and 13 % of those develop traumatic cardiac arrest.²⁷⁸ Diagnosis of tension pneumothorax in a patient with cardiac arrest or haemodynamic instability must be based on clinical examination or POCUS. The symptoms include haemodynamic compromise (hypotension or cardiac arrest) in conjunction with signs suggestive of a pneumothorax (preceding respiratory distress, hypoxia, absent unilateral breath sounds on auscultation, subcutaneous emphysema and mediastinal shift with tracheal deviation and jugular venous distention).³⁰² The presentation of a tension pneumothorax is not always classical, but when it is suspected in the presence of cardiac arrest or severe hypotension, chest decompression by open thoracostomy should be carried out immediately. To decompress the chest in traumatic cardiac arrest, perform bilateral thoracostomies in the 4th intercostal space at the midaxillary line, allowing extension to a clamshell thoracotomy if required. Needle thoracocentesis should be attempted if the provider is not competent to perform a thoracotomy, or if the necessary surgical instruments are not immediately available.^{303–306}

Cardiac tamponade and resuscitative thoracotomy. Cardiac tamponade is a frequent cause of cardiac arrest in penetrating chest trauma. Those with the clinical ability, competence, and equipment to perform it, can restore circulation with immediate resuscitative thoracotomy via a clamshell or left anterolateral incision.^{307,308} A systematic review on needle pericardiocentesis in traumatic cardiac tamponade concluded that there remains a limited role for this method in non-trauma centres where definitive surgical management is not immediately available.³⁰⁹

While resuscitative thoracotomy is often performed for the relief of cardiac tamponade, it is not exclusively indicated for this condition. Resuscitative thoracotomy is also applicable for direct haemorrhage control, aortic cross-clamping, and open cardiac compressions.²⁸²

A recent systematic review and meta-analysis examined resuscitative thoracotomy in civilian thoracic trauma, analysing 49 studies across both pre-hospital and emergency department settings.²⁵¹ The study found that timing is critical, as performing pre-hospital resuscitative thoracotomy more than five minutes after arrival at the scene was associated with increased neurological complications, while a delay of more than ten minutes from the initial encounter to resuscitative thoracotomy was linked to higher mortality rates. Patients with an Injury Severity Score of 25 or higher and those without signs of life also had poorer outcomes. Overall, mortality was higher in pre-hospital resuscitative thoracotomy (93.5 %) compared to ED-resuscitative thoracotomy (81.8 %) ($P = 0.02$). Among emergency department-resuscitative thoracotomy cases, blunt trauma patients had significantly higher mortality (92.8 %) compared with penetrating trauma patients (78.7 %) ($P < 0.001$). In summary, emergency resuscitative thoracotomy is a high-risk intervention, with better survival rates in the hospital, and timely intervention within 5–

10 min is crucial for improving outcomes. Blunt trauma patients have poorer survival prospects compared with those with penetrating injuries.

The prerequisites for a successful resuscitative thoracotomy can be summarized as the 'four E rule':

- **Expertise:** teams that perform resuscitative thoracotomy must be led by a highly trained and competent healthcare practitioner. These teams must operate under a robust governance framework.
- **Equipment:** adequate equipment to carry out resuscitative thoracotomy and to deal with the intrathoracic findings is mandatory.
- **Environment:** ideally resuscitative thoracotomy should be carried out in an operating room. Resuscitative thoracotomy should not be carried out if there is inadequate physical access to the patient, or if the receiving hospital is not easy to reach.
- **Elapsed time:** the time from loss of vital signs to commencing a resuscitative thoracotomy should not be longer than 15 min

If any of the four criteria is not met, resuscitative thoracotomy is futile and exposes the team to unnecessary risks.³¹⁰ A retrospective cohort study found that prehospital resuscitative thoracotomy for traumatic cardiac arrest was associated with 3.8 % of patients surviving with favourable neurological outcome, all presenting with cardiac tamponade following penetrating trauma, and none of the patients with exsanguination or other pathology surviving with favourable neurological outcome. There were no survivors beyond 15 min of traumatic cardiac arrest for cardiac tamponade and 5 min after exsanguination-induced cardiac arrest.³¹¹

Post resuscitation care after traumatic cardiac arrest

As with all aetiologies of cardiac arrest, optimising cardiac output is essential for restoring oxygenation and meeting the metabolic demands of post-arrest physiology. This includes fluid resuscitation and vasopressor support as needed to maintain adequate tissue perfusion and oxygen delivery. Early whole-body CT scanning can help identify major injuries and guide initial management following ROSC by detecting life-threatening conditions such as ongoing haemorrhage, pneumothorax, or cardiac tamponade, which may require immediate intervention.

The principle of damage control resuscitation is advocated for uncontrolled haemorrhage in trauma. Damage control resuscitation integrates permissive hypotension, haemostatic resuscitation, and damage control surgery. Evidence supports a conservative IV fluid approach, maintaining blood pressure at a level no more than is necessary to sustain a radial pulse, until surgical haemostasis is achieved. However, caution is needed in traumatic brain injury, where raised intracranial pressure may necessitate a higher blood pressure to achieve sufficient cerebral perfusion pressure. The duration of hypotensive resuscitation should not exceed 60 min, as the risk of irreversible organ damage then outweighs its benefits.^{312–314}

Temperature management is crucial, as hypothermia exacerbates coagulopathy, acidosis, and haemodynamic instability (the 'lethal triad'). To prevent further deterioration, active warming strategies, including fluid warming, external warming devices, and maintaining an optimal resuscitation environment, should be implemented early.

Coagulation management is a key component of resuscitation, as trauma-induced coagulopathy can worsen haemorrhage and organ dysfunction. Strategies include early administration of blood products

(e.g., plasma, platelets, fibrinogen or whole blood), tranexamic acid, and point-of-care coagulation monitoring to guide targeted haemostatic therapy.

Special settings

Cardiac arrest in the catheterisation laboratory

The occurrence and outcomes of cardiac arrest in the cardiac catheterisation laboratory ('cath lab') vary depending on the setting, patient profile and type of intervention performed. Cases range from routine elective procedures to the emergent treatment of critically ill patients with cardiac ischaemia, shock, or life-threatening arrhythmias. Additionally, structural heart interventions, often targeting elderly and high-risk patients, are becoming increasingly prevalent and complex. Causes of cardiac arrest among others in this setting include severe ischaemia, tamponade, arrhythmias, vascular perforation/dissection, or anaphylaxis.

A recent ILCOR scoping review³¹⁵ estimated the overall incidence of cardiac arrest in the catheterisation lab as 0.2 % to 0.5 % across all procedures.^{316,317} For PCI, the incidence is slightly higher (0.8–2 %),^{318–320} and for ST-elevation myocardial infarction higher still (STEMI – 1.9 % to 5.5 %), primarily with an initial rhythm of ventricular fibrillation.^{321–325}

Overall initial survival rates range from 67 % to 77 % at the event time^{316,317,326} and between 38 % to 56 % at hospital discharge.^{317,326} For PCI and STEMI, survival to discharge is higher (82–100 %).^{321,322,325}

Prevent and be prepared

Adequate Education is essential to ensuring the best cardiac arrest outcomes in the catheterisation lab. Staff should be trained in advanced life support and prepared to quickly recognise and manage procedure-specific risks and complications, including pacing, pericardiocentesis, initiation of ventricular assist devices, or ECPR. Emergency equipment should be functional and readily available (See Fig. 16).

Studies support periodic simulation emergency drills to enhance technical and non-technical skills in the catheterisation lab.³²⁷ Training in non-technical skills has the potential to minimize errors in this setting.³²⁸ Careful risk–benefit assessment and planning for elective procedures are advised to minimise complications. The use of safety checklists has been reported as effective in reducing human errors and has resulted in fewer complications.³²⁹

Detect and react

Patient's monitored vital parameters including ECG must be checked regularly to detect early complications. Echocardiography should be readily accessible in case of haemodynamic instability or suspected complication (e.g. cardiac tamponade). Transoesophageal echocardiography (TOE), often used for monitoring structural heart interventions in the catheterisation lab, may facilitate early detection of complications with higher quality images and precision.³³⁰

Cardiac arrest in the catheterisation lab should prompt an immediate call for help and activation of the resuscitation team.³¹⁶ While catheterisation lab staff must promptly initiate CPR, additional support is needed to sustain CPR while addressing reversible causes. Depending on the suspected underlying complication or cause of arrest, the cardiac surgery team should also be alerted. Importantly,

as highlighted by the Joint British Societies' guideline on management of cardiac arrest in the catheterisation lab,³³¹ all emergency team members should don lead aprons before entering, since fluoroscopy may often be necessary. Ensure that an adequate supply of lead aprons, protection for eyes and thyroid is readily available in advance.

Resuscitate and treat possible causes

Cardiac arrest in the catheterisation lab should be generally managed according to the ERC Guidelines 2025 ALS,¹⁶⁴ with some modifications. In the presence of monitored VF/pVT, immediate defibrillation with up to three stacked shocks if a shockable rhythm persists is recommended before starting chest compressions. This approach aligns with the recommendations for witnessed arrests in monitored patients with a shockable rhythm when a defibrillator is readily available. In case of PEA/ asystole, CPR should be initiated and adrenaline injected. However, extreme bradycardia may be treated with external or transvenous temporary pacing, especially if it arises from atrio-ventricular block complicating certain procedures, such as transaortic valve implant, tricuspid valve interventions or catheter ablation. During cardiac arrest, invasive procedures to correct reversible causes (such as PCI, pericardiocentesis, or mechanical support device initiation) might be needed during on-going CPR or immediately after ROSC.³³¹

Point of care ultrasound (POCUS)

ILCOR reviewed the diagnostic accuracy of POCUS in detecting reversible causes of cardiac arrest in different scenarios, concluding that POCUS evaluation may be considered during cardiac arrest in the catheterisation lab if performed by experienced personnel without interrupting CPR, especially when a specific reversible cause is clinically suspected.³³² In patients already undergoing TOE at the time of cardiac arrest, maintaining the TOE probe in place could be considered if the airway is secured, since TOE may offer additional feedback and assist in guiding ECPR or mechanical support device placement if performed by a skilled operator, without interfering with chest compression. However, the use of these tools should not compromise adherence to the ALS protocol.

Mechanical CPR

An ILCOR systematic review evaluated the impact of mechanical chest compression devices compared with manual compression on cardiac arrest outcomes,²⁴² and found evidence against the routine use of automated mechanical chest compressions to replace manual compressions for in-hospital cardiac arrest. However, mechanical CPR was considered a reasonable alternative when high-quality manual compressions may compromise the safety of the provider or interfere with critical procedures, such as those performed in the catheterisation lab. Additionally, the ILCOR scoping review on cardiac arrest in the catheterisation lab³¹⁵ described outcomes of cardiac arrest in the catheterisation lab following the use of mechanical CPR, mainly during PCI or cannulation for mechanical circulatory support or extracorporeal CPR. The mixed cohorts included in the remaining studies and the inconsistent outcome reporting made interpretation difficult, so the overall results in this setting are uncertain. Until new evidence is produced, mechanical CPR may be used in the catheterisation lab following the ILCOR CoSTR considerations discussed above.²⁴² Importantly, if used,

1. Prevent & be prepared



Promote adequate training of the staff in technical skills & ALS



Ensure emergency equipment is functional and readily available



Plan elective procedures carefully and use safety checklists

2. Detect and react



Check patient's status and monitored vital signs regularly




Consider echocardiography if haemodynamic instability or complication



In case of cardiac arrest, call for help & activate the resuscitation team

3. Resuscitate and treat possible causes



Continue CPR according to the ALS algorithm

- Check & correct reversible causes, including **ultrasound & angiography**
- Consider **mechanical CPR** to facilitate chest compression
- Consider **ECPR** in refractory cardiac arrest
- Consider **circulatory support devices** in case of shock after ROSC

Fig. 16 – Treatment of cardiac arrest in catheterisation laboratory.

ALS – advanced life support, VF – ventricular fibrillation, pVT – pulseless ventricular tachycardia, ROSC – return of spontaneous circulation, PEA – pulseless electrical activity, POCUS – point of care ultrasound.

pauses during device placement should be minimized, and correct positioning ensured to avoid visceral injuries.

Extracorporeal CPR

The use of veno-arterial extracorporeal membrane oxygenation (VA-ECMO) to support refractory cardiac arrest patients (ECPR) was recently updated by ILCOR.²⁴³ In the in-hospital setting, ECPR was considered as a rescue therapy for selected patients with refractory cardiac arrest if implementation is feasible. A recent ILCOR scoping review³¹⁵ covered the use of ECPR in the catheterisation lab. Most evidence derived from the Extracorporeal Life Support Organization registry with 39 % survival to hospital discharge.³³³ Despite its potential advantages in sustaining resuscitation while addressing reversible causes, evidence is limited due to study heterogeneity and high risk of bias. Nevertheless, considering the risk–benefit ratio, ECPR may be considered for selected cardiac arrest patients in the catheterisation lab based on clinical circumstances, available resources and experience. ECPR facilitates performing diagnostic or therapeutic procedures, including coronary angiography/angioplasty, pulmonary embolectomy or cardiac surgery.

Mechanical circulatory support

A variety of temporary devices can be used in the catheterisation lab to assist cardiac pumping function, offering different levels of haemodynamic support for severe heart failure or during high-risk procedures. Common examples include the intra-aortic balloon pump, Impella®, TandemHeart® or VA-ECMO.

The use of mechanical circulatory support devices for cardiogenic shock after ROSC in any setting was addressed by ILCOR.³³⁴ This showed no survival benefit at different follow-up points when comparing early routine device use with standard care. However, most evidence derived from cardiogenic shock patients rather than those specifically resuscitated from cardiac arrest. In summary, mechanical circulatory support in patients with cardiogenic shock after ROSC in the catheterisation lab may be considered in selected cases. Individualised decisions should consider the clinical picture, device availability and team expertise. When used, close monitoring for complications is recommended to ensure prompt management.

Intracoronary adrenaline

The recent ILCOR scoping review on cardiac arrest in the catheterisation lab included a comparison of intracoronary adrenaline with either peripheral intravenous or central venous administration.³³⁴ Adrenaline via central intravenous and intracoronary routes was associated with higher rates of ROSC, survival to hospital discharge, and survival with good neurological outcomes compared with peripheral intravenous administration. However, intracoronary administration was associated with an increased risk of stent thrombosis. Despite these promising findings, current evidence is insufficient to recommend routine intracoronary administration of adrenaline during cardiac arrest in the catheterisation lab.

Drowning

Once the World Health Organization recognised accidental drowning as a serious public health problem—especially in low-resource settings—annual global deaths from drowning decreased from 370,000 in 2000 to 300,000 in 2021³³⁵. This decline might be attributed to legislation and regulations (e.g., pool fencing and mandatory

lifejacket use),³³⁶ as well as effective preventive measures (e.g., day-care centres for children and water safety training).³³⁷ Fatal drownings are associated with several risk factors: 1) sex: males account for up to six times more drownings than females; 2) age: although in Europe drowning is the fourth leading cause of death in children aged 5–14 years, those older than 50 years of age have the highest incidence; 3) location: inland waterways represent a significant risk location; 4) circumstances such as alcohol consumption and migrant status increase the likelihood of drowning. The WHO statistics do not include fatalities from drowning caused by suicide (the most frequent cause of drowning death in several European countries), transport accidents, or natural disasters such as floods.^{338–340} Most fatal drownings never reach the health system because rescue or retrieval often occurs too late for medical intervention.³⁴¹

Recommendations are based on a scoping review in 2021 and a systematic review in 2023.^{342,343} There are no RCTs. Most recommendations are good practice statements or expert opinions based on indirect evidence and are in line with the position statements from the Medical Commission of the International Lifesaving Federation.³⁴⁴

Drowning is defined as ‘the process of experiencing respiratory impairment from submersion or immersion in liquid’.³⁴⁵ Immersion is a situation in which the person stays in a liquid with the airway above the liquid surface, while submersion refers to the state in which the airway remains submerged. When respiratory impairment progresses during submersion, the heart slows down and finally stops due to prolonged hypoxia, and only resuscitation can prevent the drowning mechanism to become fatal.^{346,347} Submersion time is the most robust and independent prognostic factor.³⁴⁸ The Adult BLS CoSTR 2020 recommends submersion time as the sole indicator guiding prognosis and decisions regarding clinical management and search and rescue operations.³⁴⁹ Less than 5–10 min of submersion is associated with a better prognosis.^{348–350}

The extent of lung damage depends on the amount and degree of pollution of the aspirated water.³⁴⁶ Salinity is of no proven predictive value.^{348,349,351}

The autonomic conflict, caused by simultaneous stimulation of the sympathetic and parasympathetic autonomic nervous system may explain the genesis of cardiac arrhythmias in circumstances with cold water immersion and breath-holding.³⁵² In drowning, low water temperature may exert a protective effect by reducing brain metabolism.^{346,348,349} However, this phenomenon is exceedingly rare and is most likely in small individuals who cool quickly in water below 6 °C.³⁵³ Submersion longer than 60 min with good neurological outcomes has been reported,³⁵⁴ particularly in children immersed in cold water, preceding the submersion.^{354,355} These reports suggest that if water temperature is above 6 °C, survival or successful resuscitation becomes extremely unlikely after about 30 min of submersion, whereas in water at or below 6 °C, this threshold may extend to around 90 min.³⁵⁴

Water rescue

Water rescue is needed to prevent drowning, to interrupt drowning, and to provide immediate life-saving intervention. Many untrained people have died while attempting a rescue.³⁵⁶ Rescuers should never deliberately place themselves in danger. For this reason, it is strongly recommended that bystanders, especially those who are

unable to swim, do not enter deep water. Instead, they should rely on indirect rescue methods, such as using public rescue equipment, throwing any available flotation device or reaching out with a long object, while seeking professional assistance. First responders with rescue training have the competencies to select and apply the appropriate rescue technique, rescue material and flotation devices where possible.³⁵⁷ Rescue material and flotation devices also reduce the rescue time.³⁵⁷

Physical fatigue is a limiting factor when CPR is required, especially after water rescue.^{358,359} If possible, a person not involved in the rescue should perform CPR.³⁵⁷ Spinal injuries are uncommon after drowning.³⁶⁰ Spinal stabilisation should not delay assessment of vital signs or resuscitation. If resuscitation is not needed and clear signs of a cervical spine injury are present, it is recommended that at least three people carry out spinal motion restriction during extrication from water, preferably with at least one of them specifically trained in the procedure.³⁶¹ If the required people are unavailable, the extrication should not be postponed any further. In every situation of a spinal injury in water, emergency medical services should be alerted immediately.

Cardiac arrest caused by drowning

Evidence showed better prognosis when drowned persons are ventilated when initiating CPR.^{362–364} For this reason, in-water ventilation and on-board resuscitation are gaining acceptance within the lifesaving community, as these techniques are feasible with training.^{342,362,365–367} In practice, some circumstances do not allow this. For example, due to lack of training, absence of flotation equipment or the conditions of the water. Then, it is more practical to wait until the victim is onshore.

Onshore, first responders and EMS should follow the ABC approach.^{362,365} CPR should start with 5 ventilations. If the person remains unconscious without normal breathing, continue with standard CPR (Fig. 9). Starting CPR with compression is only advised if the rescuer is unwilling to provide ventilations.³⁶⁸

For simplicity, untrained bystanders should start with chest compressions whilst trained first aid providers should consider ventilations, since the time to ventilation is critical in restoring cardiac arrest in drowned victims. In isolated respiratory arrest, ventilations can prevent cardiac arrest (see Fig. 10).

In drowning, an initial shockable rhythm is present in less than 10 % of cases because of the cardiac response to hypoxia. Although victims removed from the water might be in cardiac arrest due to cardiac cause, for example because of arrhythmias caused by autonomic conflict³⁵² or the presence of cardiovascular disease,³⁶⁹ CPR should not be delayed by AED application.

Oxygen should be provided early because drowning is a respiratory event. However, a recent study showed that it is difficult to provide evidence of the impact of oxygen on prognosis.³⁷⁰ Airway management should be based on the competencies and training of the first responder or EMS.³⁶⁵ Tracheal intubation compared with supraglottic airway devices was associated with higher ROSC, but not associated with survival or favourable neurological outcome at one month.³⁷¹

Impact of hypothermia

Most drowned victims are hypothermic (core body temperature < 35 °C), which affects diagnosis, treatment and prognosis.^{346,348,372,373} In general, guidelines for accidental hypothermia can also be applied to hypothermic drowned victims, including the

one-minute vital sign check, accurately measuring core temperature, the alternative treatment regarding defibrillation, use of medication when body temperature is below 30 °C, and access to extracorporeal life support rewarming.³⁷² Hypothermia reflects a longer submersion time, resulting in poorer neurological outcome and survival. However, when drowning occurs in cold water, generally considered below 6 °C, and notably after a period of immersion, associated hypothermia may exert a neuroprotective effect.³⁷⁴

The Adult BLS CoSTR 2020 suggests not using water temperature when making prognostic decisions³⁴⁹ and no specific guidelines exist to indicate precisely when to move from rescue to recovery—a stance that also acknowledges the exclusion of rare favourable outcomes in icy water. This is partly due to the wide variety of drowning scenarios regarding submersion time, water temperature, available resources, or environmental hazards. Some agencies or countries have local or national protocols. In the absence of definitive guidance, it is crucial that those on scene reach a consensus on the point at which further medical intervention is deemed futile.³⁷⁴

Cardiac arrest in the operating room (OR)

Cardiac arrest in the operating room is a rare event with an incidence of about 3/10,000 anaesthetics,^{375–377} with a higher incidence in low-resource settings,³⁷⁸ in older-frailer patients, and in newborns and infants.^{379,380} Overall survival rate is > 50 %.³⁸¹ Strong predictors of intraoperative cardiac arrest are higher scores in the American Society of Anesthesiologists (ASA) physical status, sepsis, urgent or emergency case, complexity of the case, anaesthetic technique and age.³⁸¹ The leading causes are complications during cardiac surgery, major haemorrhage, bradyarrhythmias and septic shock.^{378,381}

Specifics of intraoperative cardiac arrest and its treatment

Mostly, gradual physiological deterioration leads to intraoperative cardiac arrest.³⁸² Cardiac arrest treatment in the OR follows the general ALS algorithm. However, several modifications intended to identify reversible causes are required. Key interventions include immediately calling for help, informing the surgical and anaesthesia team, and ensuring the presence of sufficiently skilled people.

High-risk surgical patients are often monitored with invasive blood pressure pre-arrest. Recent proposals suggest starting chest compression if systolic blood pressure stays below 50 mmHg despite interventions to treat the underlying cause.^{375,383,384} Sudden fall in ET_{CO₂} in such cases is a strong indicator for cardiac arrest. Adjust the position and the height of the operating table or trolley to optimise delivery of high-quality chest compressions.

Low initial end-tidal capnography values (ET_{CO₂} < 2.7 kPa or 20 mmHg) are linked to inadequate chest compression quality, indicating the need for improvement and possibly suggesting rescuer fatigue.^{385–387} For pre-arrest patients (SBP < 50 mmHg), initial incremental boluses of 50–100 µg adrenaline intravenously have been proposed, rather than the standard 1 mg bolus. In pre-arrest patients, higher adrenaline doses may induce severe hypertension or tachyarrhythmias. If a low-dose adrenaline bolus fails, the standard 1 mg of intravenous adrenaline should be given.^{375,388}

In circumstances where the probability of cardiac arrest is high, a defibrillator should be readily available in standby mode and self-adhesive defibrillation electrodes should be applied before induction of anaesthesia. Safe defibrillation should be performed immediately in case of a shockable rhythm. Ensure adequate venous access, pre-

pare resuscitation drugs and fluids, establish advanced airway management (if not already undertaken) and use a mechanical ventilator delivering 100 % oxygen as soon as possible.³⁸⁹ Current data suggest that mechanical ventilation yields a similar PaO₂ to that of manual ventilation with a self-inflating bag.^{390–393} If there is a qualified sonographer able to perform ultrasound (transthoracic/transoesophageal) with minimum interruptions to chest compressions, they should be summoned to aid diagnosis.³⁸⁹

Chest compression is optimally performed in the supine position, but in case of a cardiac arrest in prone position with an advanced airway in place, follow the ILCOR Good Practice Statement to initiate CPR in prone position.^{394,395} Consider simultaneous left lateral decubitus and head-down positioning in cases of massive air embolism^{396–398} if it does not affect chest compression quality.

Identification of reversible causes should be prioritised and treated appropriately:

- If arrest is caused by significant blood loss, chest compressions are effective only if the circulating volume is replaced simultaneously and haemorrhage control (e.g. surgery, endoscopy, endovascular techniques) is initiated immediately.³⁸⁹
- In cases of severe subdiaphragmatic exsanguination, resuscitative endovascular balloon occlusion of the aorta (REBOA) may be considered in an attempt to slow intravascular volume depletion. However, there is limited evidence on improved survival and further studies are required to elucidate the potential benefit.^{389,399,400}
- ECPR should be considered in cases where conventional CPR fails or when prolonged resuscitation is required.
- Open cardiac compressions should be performed only by trained healthcare professionals in cases of intraoperative cardiac arrest or as part of resuscitative thoracotomy for trauma patients.
- If arrest occurs during laparoscopic or robotic surgery, consider releasing any pneumoperitoneum and deflate the abdomen to enhance venous return during CPR; during thoracoscopy, stop CO₂ insufflation and rule out contralateral pneumothorax.
- For other reversible causes, consult the specific subchapters of these guidelines.

Human factors in intraoperative cardiac arrest. As in every resuscitation event, a designated team leader should direct and coordinate the resuscitation team and their assistants, focusing on high-quality chest compressions and ventilations, minimising no-flow times, addressing reversible causes simultaneously, and preventing focus on low-priority distracting tasks. Surgery must be stopped unless it addresses a reversible cause of cardiac arrest. There may be a need to cover the surgical field to enable access to the patient and to perform resuscitation tasks.

Successful management of intraoperative cardiac arrest requires not only individual technical skills and a well-organised team response, but also an institutional safety culture embedded in everyday practice through continuous education, training and multidisciplinary cooperation.^{8,401,402} Institutional protocols for responding to potential arrest situations (e.g. massive transfusion protocols) and checklists will help to optimise the response to cardiac arrest in the operating room environment.^{8,403}

There is no evidence to support the use of hypothermic temperature control after adult intraoperative cardiac arrest.⁴⁰⁴ Therefore,

post-arrest temperature control should follow local post-resuscitation protocols.

Local anaesthetic systemic toxicity

Cardiac arrest is a rare complication of local anaesthetic overdose, which is often caused by inadvertent intravascular injection. Direct action of local anaesthetics on the sodium channels of cardiac myocytes causes cardiovascular collapse, usually within 1–5 min, but onset was reported from 30 sec to 60 min.^{405,406} Early symptoms are *peri*-oral numbness, metallic taste, dizziness, tinnitus and blurred vision followed by substantial hypotension, dysrhythmias, and generalised seizures. The emergency diagnosis is often based on exclusion of other causes.^{407,408} Seizure management involves administering benzodiazepines in incremental doses (e.g. IV lorazepam 0.1 mg kg⁻¹ IV, midazolam 0.05–0.1 mg kg⁻¹ IV), followed by stepwise doses of propofol or pentothal (up to induction doses) if needed. Attention should be given to securing the airway and maintaining ventilation while preventing cardiovascular collapse during sedative drug administration (especially propofol). Intravenous lipid therapy is recommended as rescue therapy to treat cardiovascular collapse and cardiac arrest although its use is based on very low-certainty evidence,^{131,408–410} but without documented harm.⁴¹¹ A 20 % lipid emulsion (at least 4 bags of 250 mL) should be available if large doses of local anaesthetics are used (e.g. operating rooms, labour wards, emergency department).^{409,412} During cardiac arrest, after an initial 20 % lipid emulsion IV bolus (1.5 mL/kg IV over 1 min), an infusion should be started (0.25 mL kg⁻¹ min IV). If ROSC is not achieved in 5 min, the infusion rate should be doubled with two additional boluses at 5-minute intervals.^{409,412} A maximum cumula-



QR code. 4 – Cardiac arrest due to local anaesthetic toxicity management.

tive dose of 12 mL kg⁻¹ IV 20 % lipid emulsion is recommended to avoid fat overload.⁴¹¹ It is suggested that adrenaline IV be given at a lower dose (≤ 1 µg kg⁻¹) in local anaesthetic intoxication because its arrhythmogenic and acidotic effects may impede sustained ROSC – this has been shown in animal studies.^{409,413–416}

Further information on cardiac arrest due to local anaesthetic toxicity can be found at the [QR code. 4](#).

Cardiac surgery

The incidence of cardiac arrest following cardiac surgery is 2–5 %, with higher survival rates (around 50 %) that are higher than other causes.^{417–421} This is largely due to early detection of, and the high incidence of, reversible causes. Common causes of cardiac arrest in this setting include arrhythmias causing VF, accounting for up to 50 % of cases, followed by cardiac tamponade and major bleeding, which often presents as PEA.

Evidence-based recommendations for the management of cardiac arrest following cardiac surgery derive from the European Association for Cardio-Thoracic Surgery,^{422,423} the Society of Thoracic Surgeons expert consensus document for the resuscitation of patients who arrest after cardiac surgery,⁴²⁴ the British Societies

Guidelines,⁴²⁵ and the ILCOR CoSTR for the Management of Implantable Left Ventricle Assist Devices recipients.⁴²⁶

Ensure adequate initial and refresher training of staff in resuscitation technical skills, ALS and teamwork through simulated cardiac surgery scenarios, including training to perform an emergency resternotomy. Teamwork, including human factors such as situational awareness and communications skills, should be included in the training and acquired through practice in simulated scenarios. Roles should be previously allocated to staff in every setting to ensure effective coordination of resuscitation efforts.⁴²⁷

Emergency equipment should include small resternotomy sets containing only the essential elements to open the chest, be standardised, adequately marked, readily available where patients with recently performed interventions are recovering and periodically checked.^{423,424}

Safety checklists reduce complications and mortality in non-cardiac surgery and should be implemented for the management of cardiac arrest in these settings.⁴²⁸

Detect cardiac arrest and activate cardiac arrest protocol. Early signs of deterioration can be identified in the monitored post-operative patient after careful examination. Hypotension is a common sign of several post-operative complications (Table 9).^{429–431} Echocardiography should be performed in case of haemodynamic instability; consider transoesophageal echocardiography for more precise diagnosis.^{432,433} Continuous ECG monitoring enables early identification of arrhythmias; supraventricular tachycardias are the most frequent in this setting.⁴³⁴

Cardiac arrest can be detected by checking ECG rhythm, clinical examination and review of vital signs, including pressure waveforms (arterial, central venous and pulmonary artery pressures, and pulse oximetry) and ETCO₂.^{423,424}

Resuscitate and treat possible causes in cardiac surgery

Key modifications to the standard ALS algorithm include immediate correction of reversible causes and, if this is not successful, emergent resternotomy.^{423,424}

Resuscitate and treat possible causes:

- o Start resuscitation according to ALS but with modification:
- o VF/ pVT: Apply up to 3 consecutive shocks (<1 min).
- o Asystole/ extreme bradycardia: Apply early pacing – maximum output (<1 min).
- o PEA: Correct potentially reversible causes. If paced rhythm, transiently turn off pacing to exclude VF.

No ROSC:

- o Initiate chest compression and ventilation.
- o Consider POCUS/ TOE.
- o Perform early resternotomy (<5 min).
- o Consider circulatory support devices and ECPR (Fig. 17).

In patients with VF/pVT, defibrillation of up to three stacked shocks are prioritised within a maximum time of 60 s.^{435,436} If these fail, immediate resternotomy and internal defibrillation are advised.⁴³⁶ In case of asystole or extreme bradycardia, attempt epicardial pacing (DDD mode at 80 to 100 beats/min and at maximum output voltages) or transcutaneous pacing should be attempted for one minute before initiating chest compressions. PEA should trigger immediate external chest compressions, searching for reversible causes and preparing for early resternotomy. In the presence of a pulseless stimulated rhythm, pause the pacing to exclude underlying VF and, if indicated, perform defibrillation.^{423,424}

If ROSC is not achieved following defibrillation or pacing, or in case of PEA, initiate compressions and ventilations should be initiated while preparing for emergency resternotomy. Perform external compressions at 100–120 beats/min, aiming to reach a systolic blood pressure > 60 mmHg; failure to attain this value despite adequate

Table 9 – Common causes of patient deterioration after cardiac surgery and management.

• Haemorrhage • 'Medical' bleeding: post-operative coagulopathy • 'Surgical' bleeding: operative trauma	• Correct hypothermia and hypotension, avoid haemodilution • Consider blood product transfusion and use of haemostatic agents, guided by haematological tests • Check chest drains to identify active bleeding and perform echocardiography to exclude cardiac tamponade; consider early re-operation if suspected
• Low cardiac output state • Inadequate preload • Excessive afterload • Decreased ventricular contractility • Diastolic dysfunction	• Perform echocardiography to assess ventricular function • Ensure adequate ventricular filling • Correct systemic vasoconstriction • Maintain atrioventricular coordination • Correct metabolic disturbances and hypocalcaemia • Consider inotropic or mechanical circulatory support
Graft or valve failure	• Check for ECG abnormalities • Perform echocardiography • Consider percutaneous intervention or re-operation
Arrhythmias	• Correct electrolytic disturbances • Consider antiarrhythmic, electrical cardioversion or pacing
Vasodilation • Rewarming • Analgesics / sedatives • Sepsis • Anaphylaxis • Adrenal insufficiency • Vasoplegic syndrome	• Correct specific underlying causes • Consider haemodynamic-guided IV fluid therapy • Consider vasopressor support

1. Prevent & be prepared



Ensure adequate training of the staff in technical skills and ALS



Ensure availability and functioning of emergency equipment



Use safety checklists



Identify and manage deterioration in the post-operative cardiac patient

2. Detect cardiac arrest and activate cardiac arrest protocol



Confirm cardiac arrest by clinical signs and pulseless waveforms



Call for help and activate cardiac arrest protocol



- Ensure airway patency and breathing
- Deliver 100% oxygen
- Stop syringe drivers



Consider echocardiography to early identify possible reversible causes

3. Resuscitate and treat possible reversible causes

ASSESS RHYTHM




VF/pVT




Defibrillate
(apply up to 3 consecutive shocks)

Asystole / extreme bradycardia



Pace -
if wires available

PEA



Provide high-quality CPR

Start basic life support and address reversible causes

Amiodarone 300 mg IV

Consider external pacing
if wires not available

If paced, turn off pacing
to exclude underlying VF

No ROSC

- Continue advanced life support
- Prepare & perform early re sternotomy (<5 min)
- Consider circulatory support devices and ECPR

Fig. 17 – Post-cardiac surgery cardiac arrest algorithm.

ALS – advanced life support, VF – ventricular fibrillation, pVT – pulseless ventricular tachycardia, ROSC – return of spontaneous circulation, PEA – pulseless electrical activity, IV – intravenous.

performance may indicate tamponade or severe haemorrhage, requiring emergency re-sternotomy.^{423,424} Compared with external compressions, internal cardiac compression provides better coronary and systemic perfusion pressure, which may justify chest reopening.^{437,438} Airway management in this setting follows the usual indications for ALS.

In mechanically ventilated patients check the position and patency of the tracheal tube, increase inspiratory oxygen to 100 % and remove positive end-expiratory pressure. If a tension pneumothorax is suspected, emergency decompression is necessary.^{423,424}

Medications during resuscitation. As a general principle, stop all infusions other than those needed for resuscitation. Amiodarone (300 mg) or lidocaine (100 mg) may be administered intravenously after three failed shocks to treat VF/pVT.^{424,439} Injecting adrenaline (1 mg) shortly after cardiac surgery is controversial. The European

Association of Cardio-Thoracic Surgery and Society of Thoracic Surgeons discourage the routine use of adrenaline based on the concern that intense hypertension induced by adrenaline may cause bleeding or disruption of surgical anastomoses after ROSC^{423,424,439}, although lower doses (50–100 mcg boluses) may be considered in *peri-arrest* situations based on expert consensus.^{375,388,424,429,437}

POCUS and Transoesophageal echocardiography. In a *peri-arrest* patient following cardiac surgery, consider transthoracic POCUS, although it may give limited views of the posterior cardiac chambers and any surrounding posterior cardiac tamponade. Transoesophageal echocardiography (TOE) is a preferable alternative, giving clearer 360° views to include posterior cardiac structures, and enables uninterrupted chest compressions. The presence of cardiac tamponade, intrathoracic haemorrhage, pleural effusions, hypovolemia,




Fig. 18 – Left ventricular assist device ALS algorithm.

LVAD – left ventricular assist device, ALS – advanced life support, VT – ventricular tachycardia, VF – ventricular fibrillation, ECG – electrocardiography.

dynamic left ventricular outflow tract obstruction and aortic dissection can generally be identified using TOE.^{432,433,440–447} TOE can evaluate the effectiveness of chest compressions by real-time imaging of the emptying and filling of compressed cardiac chambers, potentially leading to repositioning of hands for chest compressions.^{448,449}

Early re sternotomy

Refractory cardiac arrest requires re sternotomy within 5 min, in order to perform open cardiac compressions, release any tamponade and correct underlying causes. This is a safe procedure in the ICU,⁴⁵⁰ leading to higher survival rates, especially if performed with minimal delay and in the presence of surgically repairable problem on reopening.⁴⁵¹ Re sternotomy should be attempted as part of the resuscitation protocol of cardiac patients until at least day 10 post-operatively.⁴²⁴

Circulatory support devices

Intra-aortic balloon pump. In patients supported by intra-aortic balloon pump who present in cardiac arrest, the device may contribute to improving coronary and brain perfusion if coordinated with cardiac compression (1:1 ratio, with maximal amplification).⁴⁵² The ECG trigger of the balloon is not reliable during resuscitation and should be switched to pressure trigger mode, or to internal mode at 100 beats/min if cardiac compressions is interrupted for a significant interval.

Extracorporeal CPR. ECPR may be considered if re sternotomy fails to revert cardiac arrest or as an initial approach, alternative to re sternotomy, for patients undergoing minimally invasive cardiac surgery or those who arrest > 10 days after initial sternotomy.⁴²⁴ However, there is limited data addressing this specific scenario, since most studies relate to cardiogenic shock or paediatric populations.⁴⁵³

Left ventricular assist devices. Mechanical circulatory support devices, especially left ventricular assist devices (LVAD) are being increasingly used as a bridge to heart transplantation or recovery in advanced heart failure patients. Patients with these devices often lack traditional clinical signs such as a pulse, even when haemodynamically stable and therefore assessment of the circulatory state using palpation, non-invasive blood pressure measurement and pulse oximetry may be difficult. Recognition of cardiac arrest is based on the absence of responsiveness and breathing. When performed, conventional CPR is recommended. The risk of device dislodgement during CPR appears minimal.^{425,426}

The ERC recommendations are based on the ILCOR 2025 CoSTR and the British Societies guidelines 2025^{425,426} on the management of emergencies in implantable LVAD recipients in transplant centres (Fig. 18).

- Immediate activation of specialised teams for unconscious LVAD patients.
- Start CPR while simultaneously attempting to restore the device if multiple rescuers are available. Consider delaying CPR for up to 2 min to attempt device restoration if a single rescuer is present.
- Troubleshoot device issues as a priority, following relevant protocols:
 - o Low Flow Alarms: Check for volume responsiveness with leg raise technique. If the alarm resolves and hypovolaemia is

confirmed, proceed with fluid resuscitation. Check haemoglobin values as soon as possible to evaluate the context of hypovolaemia, haemorrhage or dehydration. In cases of low flow resulting from high blood pressure, manage afterload with antihypertensives.

- o Pump Thrombosis: Administer anticoagulation or consider mechanical circulatory support like ECMO.
- o Electrical Failure: Ensure connection to a power source and integrity of powerlines. Replace the batteries or the controller.
- o Arrhythmias: Defibrillation or cardioversion should be performed if required, ensuring adequate sedation if the patient is still conscious. A functioning LVAD may provide sufficient brain perfusion even in fibrillating hearts.
- Simultaneously (if second responder available) address Airway, Breathing, Cardiac (Monitor/Defibrillator).
- Determine the adequacy of circulation. Evaluate the patient for MAP > 60 mmHg, absence of cyanosis, and presence of an audible LVAD hum. Echocardiography and invasive monitoring may be required for precise assessment. In cases of inadequate circulation, consider temporary mechanical support or corrective surgery. If the device:
 - o Is working well and patient not in circulatory failure: Perform a complete ABCDE assessment
 - o Is still not working or patient is in circulatory failure: Start CPR and
 - consider chest reopening if < 10 days from implantation
 - deliver standard ALS, addressing 4H & 4 T.

Cardiac arrest in sports

The absolute risk of experiencing of cardiac arrest during physical exercise is small.⁴⁵⁴ The incidence of cardiac death associated with sport or exercise in the general population is reported between 0.46 and 6.8 per 100,000 person-years.^{455–458} Large population-based studies of cardiac arrest in athletes from the USA and Europe indicate regional differences in the underlying causes of death.^{459,460} About a third survived; most cases occurred during non-elite competitive or recreational sports, and bystander CPR was performed in 75 %. In patients aged ≤ 35 years, premature coronary artery disease and sudden arrhythmic death syndrome predominated, followed by myocarditis. In athletes aged ≥ 35 years, coronary artery disease predominated.⁴⁶¹ The likelihood of sport-related cardiac arrest is highest for males aged between 40 and 60 and with pre-existing cardiomyopathy. In 22 % of cases no pathology was found and survival was better with early resuscitation, use of AED, and presence of professional staff.⁴⁶² Highest incidences are reported for football players during competition and for running and gym exercise during non-competition.⁴⁶³

Cardiac arrest during sport or exercise requires rapid recognition and effective treatment if the affected individual is to survive. Improved survival of cardiac arrest during sport is attributed to witnessed events, prompt resuscitation and availability of an AED.^{455,464} This evidence supports the importance of planning, adhering and implementing standard BLS in event-prone sport events.

For those athletes who have a shockable cardiac rhythm during cardiac arrest, a minimum of three shocks should be delivered on the field of play if the shockable rhythm persists.

The medical team should consider moving the athlete in cardiac arrest to a designated rendezvous point, in the safest and most effi-

cient way. If that is not possible, CPR should be continued on the field.⁴⁶⁵ These plans should be rehearsed regularly.

Unlike in professional sports, recreational athletes, supporters and fans may be less aware of the risk of cardiac arrest, highlighting the importance of awareness programs. Initiatives such as the ERC–UEFA cooperation in 2024 during the European championships demonstrate how strategic collaborations between sports and health organisations can raise awareness and improve outcomes.⁴⁶⁶ Once an athlete is successfully resuscitated, thorough cardiological assessment and follow-up are vital before considering a return to sport.⁴⁶⁷

Prevention

There is no consensus between the major organisations on preventative measures. The International Olympic Committee recommend cardiac screening for athletes. The European Society of Cardiology recommends 12-lead ECG as a screening tool for all young athletes,⁴⁶⁷ but the AHA and American College of Cardiology conclude that there is insufficient evidence to support this as a screening measure.^{468,469} Processes applied often follow economic views rather than a health care perspective. Primary prevention includes a targeted history and a physical examination, and the addition of a 12-lead ECG to at least prevent some unnecessary deaths, but this may still overlook several important conditions associated with cardiac arrest including high-risk anomalous coronary origins, aortopathies and adrenergically-mediated arrhythmias. Beyond an ECG, there are insufficient data to support additional routine testing.⁴⁷⁰

For older participants in sports and exercise, a medical evaluation of their individual risk should include the current level of physical activity, their known cardiovascular, metabolic, or renal diseases, the presence of the signs or symptoms suggestive of cardiovascular disease and the desired or anticipated exercise intensity.⁴⁷¹ In this context, screening approaches should be adapted to the specific characteristics and risk profile of the target population.^{467,470}

Comotio cordis

Comotio cordis is a rare but potentially fatal cause of cardiac arrest, triggered by a blunt, non-penetrating impact to the precordium, typically occurring within a 20-millisecond window during the rising phase of the T wave, inducing VF.^{472,473} It has historically been cited as a leading cause of cardiac arrest in young athletes, particularly in sports involving high-velocity projectiles such as baseball, lacrosse, and hockey.^{474,475} In football, comotio cordis has been reported in 9 % of cardiac arrests in players ≤ 35 years old, with 79 % of cases caused by a ball impact to the chest.⁴⁶²

Survival rates are approximately 66 % in 2008–2023.⁴⁷⁴ Witnessed CA, early recognition, immediate CPR initiation, and rapid defibrillation with an AED remain the most important factors for survival.^{474–477} Survival rates are 40 % when resuscitation occurs within 3 min, but drop to just 5 % when delayed beyond this window.^{475,476}

Protective gear, such as chest protectors in lacrosse has been introduced in attempt to prevent comotio cordis, but recent studies suggest that they do not reduce the risk of VF in critical impacts.^{478,479}

Emergency medical services (EMS) and transportation

EMS play a crucial role in the chain of survival by providing timely, high-quality care to patients experiencing life-threatening emergencies, including cardiac arrest. This chapter provides evidence-based recommendations for resuscitation practices in the context

of EMS (including specialised medical transport services) and transportation.

EMS collect valuable data that can support continuous quality improvement and provide feedback to healthcare providers involved in CPR.⁴⁸⁰ However, validity of data should be assessed before use.^{481,482} Feedback systems have a positive effect on the quality of care and medical personnel are willing to receive feedback.^{483,484} A recent systematic review showed improvements in documentation, protocol adherence, and small effects on cardiac arrest performance, clinical decision-making, ambulance times, and survival rates.⁴⁸⁵

Resuscitation during transport

CPR quality is reduced if performed during transportation, affecting correct hand position, chest compressions rate and depth, pauses, and overall CPR quality.^{201,486} Only a few indications justify ongoing resuscitation during transport (advanced interventions not available in the prehospital setting as outlined in this chapter). Whenever possible, mechanical CPR should be implemented to mitigate the risk to both patient and rescuer.

Arterial line in the prehospital setting

Invasive blood pressure measurement in OHCA patients in a HEMS situation is feasible,^{383,487–491} and can help prehospital teams to guide resuscitation and post-resuscitation care,^{383,487,489,492} specifically in haemodynamically unstable patients.^{493,494}

Resuscitation by two-member ALS crews

Teams larger than two ALS members improve CPR quality and efficiency in rhythm recognition, adrenaline administration, and intubation.⁴⁹⁵ No significant no-flow fraction differences were found between two-, three-, and four-member teams.⁴⁹⁵ Two-paramedic teams were slower and more error-prone than paramedic-emergency medical technician teams.⁴⁹⁵ The evidence is not robust enough to support formal recommendations for education, protocols, or equipment regarding ALS provided by two HCPs. However, studies suggest that pre-filled syringes and automation-assisted protocols might improve team performance.⁴⁹⁵

Inflight cardiac arrest and microgravity resuscitation

Inflight cardiac arrest

Although air travel is safe in general, physiological changes during air travel, passenger demographics, pre-existing medical conditions and the number of passengers aboard larger aircraft and long distance flights raise the probability of in-flight emergencies.^{496,497} A meta-analysis reported an incidence 0.09 cardiac arrest events per million passengers.⁴⁹⁸ Not all airlines are equipped with an AED and ALS equipment.^{499–501}

Medical professional help should be sought via in-flight announcement. The rescuer should kneel in the leg-space in front of the aisle seats to perform chest compressions if the patient cannot be transferred within a few seconds to an area with adequate floor space (e.g. galley). Overhead-CPR is a possible option in limited space environments. Airway management should be based on the equipment available and the expertise of the rescuer. If the flight plan is over open water, with high possibility of ROSC during an ongoing resuscitation, consider an early diversion. If ROSC is unlikely (pre-existing conditions, absence of an AED), the expected time for diver-

sion and landing would be too long to improve the prognosis reasonably.

Microgravity resuscitation. CPR is challenging even for trained professionals on earth, and spaceflight adds further complications due to microgravity, confined space, and limited resources. With the rise of commercial missions by private companies, clear guidelines are essential for managing medical emergencies in both long-duration and short-term space missions.

Recommendations exist from the German Society of Aerospace Medicine and the European Society of Aerospace Medicine Space Medicine Group.⁵⁰² However, a recent simulation study⁵⁰³ using underwater methods to approximate microgravity found that all commonly recommended CPR techniques⁵⁰⁴ (including the Reverse-Bear Hug, Schmitz Hinkelbein, Cologne, and Evetts-Russomano) resulted in rates below 5 % of effective chest compressions. In the absence of data from real space environment, it is not possible to endorse any particular CPR method.

The ERC key recommendations are summarised here

- Airway management, defibrillation and IV/IO access are similar to terrestrial ALS, but only once the patient is secured.
- Consider mechanical CPR
- Consult telemedicine support during cardiac arrest in low earth orbit if feasible and manpower allows.
- The crewmember with the highest medical qualification should decide on termination of resuscitation, consulting telemedicine support.

Cruise ships

There is limited evidence about treatment of OHCA on cruise ships. The ERC guidelines recommend adhering to standard BLS and ALS protocols. Outcome from cardiac arrest on cruise ships is expected to be worse, compared with the general population, because access to healthcare facilities is more complicated, on-board resources are limited, and transfers can be prolonged.⁵⁰⁵ A medical first-responder team should be available 24/7. All equipment necessary for ALS should be available onboard and readily accessible. An AED should be onboard and requested immediately. Where there are too few numbers of crew health care professionals, an onboard announcement should be made to call for further medical professional help. In most cruise ships telemedicine is available, and it should be used as early as possible.⁵⁰⁶ Qualified medical air transportation is an option to cover long distances to medical facilities.

Special patient groups

Asthma and chronic obstructive pulmonary disease

Evidence based recommendations for the management of acute life-threatening asthma are based on the British Thoracic Society, Scottish Intercollegiate Guidelines Network⁵⁰⁷ and the Global Initiative for Asthma Strategy 2021,⁵⁰⁸ while for chronic obstructive lung disease (COPD) recommendations are based on those of the Global initiative for chronic obstructive lung disease.⁵⁰⁹ A recent ILCOR Evidence Update⁸ did not reveal any new data on management of cardiac arrest in asthma patients beyond the 2021 ERC guidelines.

Patients with an exacerbation of obstructive lung disease (asthma/ COPD) are at high risk for cardiac arrest.

For COPD, the Global initiative for chronic obstructive lung disease guidelines recommend that supplemental oxygen is titrated to achieve a target saturation of 88–92 %, with frequent monitoring of blood gases to ensure adequate oxygenation without carbon dioxide retention. Pharmacological therapy comprises inhaled short-acting beta-2 agonists with or without short-acting anticholinergics (with repeated dosing as required), systemic corticosteroids and, in case a bacterial infection is suspected, antibiotics. Non-invasive ventilation is recommended in the presence of respiratory acidosis ($\text{PaCO}_2 > 6 \text{ kPa}/45 \text{ mmHg}$ and arterial $\text{pH} \leq 7.35$); severe dyspnoea with clinical signs of fatigue or increased work of breathing. Escalation to invasive ventilation may be necessary if NIV fails, if the patient cannot tolerate NIV, or there are factors such as agitation, reduced consciousness, high aspiration risk, cardiovascular instability, or life-threatening hypoxia. However, be alert to the higher risk of life-threatening hypotension after emergency intubation and mechanical ventilation in patients with raised arterial CO_2 and obstructive lung disease.^{510,511} In selected unconscious cases with severe hypoxemia, veno-venous ECMO may be an option to prevent hypoxic cardiac arrest.^{512,513}

Treatment of cardiac arrest caused by obstructive lung disease

Cardiac arrest in patients with obstructive lung disease may be caused by hypoxia, hypovolaemia, toxic agents (arrhythmias caused by stimulant drugs e.g. beta-adrenergic agonists, aminophylline), electrolyte disturbance (hypokalaemia), tension pneumothorax and/or the effects of gas trapping leading to reduced venous return and hypotension.^{514–519} Cardiac arrest in obstructive lung disease is usually associated with a non-shockable rhythm and poor survival rates.^{520,521}

Because of extremely elevated inflation pressure during severe asthma attacks there is a significant risk of gastric inflation and aspiration with simultaneous hypoventilation of the lungs when attempting to ventilate a severe asthmatic patient with bag valve mask devices. The trachea should be intubated as soon as possible during cardiac arrest caused by asthma, by someone who is trained and competent to do so.

Tension pneumothorax may develop because of high airway pressure, which, if left untreated, may cause cardiac arrest. Check for signs of tension pneumothorax and treat accordingly.^{522,523} Disconnect from positive pressure ventilation if air-trapping and hyperinflation occurs and apply pressure to manually reduce the hyperinflation. Some case reports have described ROSC in patients with air trapping when the tracheal tube was disconnected from the breathing system.^{524–530} If dynamic hyperinflation of the lungs is suspected during CPR, compression of the chest while temporarily disconnecting the tracheal tube may relieve air trapping.^{524,527,529} Although this procedure is supported by limited evidence, it is unlikely to be harmful in an otherwise desperate situation.

Ventilating the lungs with a slower respiratory rate of 8–10 breaths per minute and sufficient tidal volume to cause the chest to rise during CPR should minimise dynamic hyperinflation of the lungs (air trapping).⁵²⁸ In mechanically ventilated patients with severe asthma, increasing the expiratory time (achieved by reducing the respiratory rate and changing inspiration to expiration time ratio) provides only moderate gains in terms of reduced gas trapping when a minute volume of less than 10 L min^{-1} is used.⁵³¹

No studies evaluating the use of IV fluids for cardiac arrest caused by obstructive lung disease were identified. Consider giving intravenous fluids to patients with obstructive lung disease, particularly those with an acute exacerbation of asthma, because they be dehydrated as a result of reduced oral intake and/or increased insen-

sible losses. Give fluids cautiously to prevent potential adverse effects.⁵³²

ECPR has been used successfully in patients with life threatening asthma.^{513,533} Consistent with the ERC Guidelines 2025 ALS, ECPR may be considered if conventional therapies fail and there is immediate access to this treatment.

Cardiac arrest in haemodialysis patients

Chronic kidney disease is a global health problem affecting 1 in 10 people worldwide⁵³⁴ and the WHO projects it will become the 5th most prevalent chronic condition by 2040.⁵³⁵

Risk factors for cardiac arrest in dialysis patients are summarised in Table 10. Fluid and electrolyte disturbances are common, with the highest risk period being just before the first dialysis session of the week, i.e. on Mondays or Tuesdays.^{536–540} The risk period extends over the 12 h after start of treatment. This suggests that the non-physiological rapid removal of toxins, fluid and electrolytes, most pronounced during the first session of the week, accounts for the high risk period.^{541,542} Clinically significant arrhythmias are also more common on dialysis compared with non-dialysis days.⁵⁴³

Out-of-hospital cardiac arrest in haemodialysis patients

Cardiac arrest accounts for 47.1 % of deaths in HD patients, occurring 20 times more frequently in dialysis patients compared with the general population.^{544–546} Within a dialysis centre, the odds of VF was found to be 5-fold greater in patients during dialysis and 14-fold greater in patients who arrest after dialysis compared with events occurring before dialysis.⁵³⁷ In a cohort of dialysis patients with a wearable cardioverter defibrillator, VT/VF was the initial rhythm in 78.6 %, with most occurring during dialysis.⁵⁴⁷

Table 11 was reproduced with permission from UK Kidney Association Hyperkalaemia Guideline in adults.⁵⁰

A three-fold increase in odds to hospital discharge with favourable neurological outcome has been shown when CPR is initiated by dialysis staff rather than awaiting the arrival of emergency services.⁵⁴⁶ Following cardiac arrest, 24 % of haemodialysis patients survived to hospital discharge and 15 % still were still alive at 1 year

following cardiac arrest.⁵³⁷ The poor outcome in one report may reflect the low rate of bystander CPR and AED use.^{548,549}

In-hospital cardiac arrest in haemodialysis patients

The incidence of in-hospital cardiac arrest (IHCA) in patients receiving haemodialysis (HD) is approximately 20 times higher than in the general population (6.3 % vs. 0.3 %), with dialysis patients accounting for 17 % of all IHCA cases. The incidence of cardiac arrest specifically in dialysis clinics ranges from 3.4 to 7.8 per 100,000 dialysis sessions. Most events are witnessed, with 70–80 % occurring during treatment.^{537–539,550} Multiple studies in HD patients demonstrate ROSC in 44–72 %^{551–553} and survival to hospital discharge ranges from 22–31 %.^{551–555} Duration of CPR and older age were predictors of a poor outcome.⁵⁵⁵ The ‘Get With The Guidelines-Resuscitation’ registry shows comparable neurological outcomes in HD patients compared with non-dialysis patients (17 % vs 16 %, $p = 0.07$).⁵⁵¹ These findings suggest that outcomes for HD patients are no worse than other patients dispelling the perception of futility.⁵⁵⁶ Nevertheless, a large national registry of IHCA found that dialysis patients had lower overall resuscitation quality compared with case-matched non-dialysis patients.⁵⁵¹ Shortfalls in resuscitation practice in dialysis patients include a delay in initiation of CPR⁵⁴⁶ and failure to achieve timely first defibrillation for a shockable rhythm.^{536,546,551}

Modifications to cardiopulmonary resuscitation in haemodialysis patients

Start resuscitation by following the standard ALS algorithm. Assign a trained dialysis nurse to operate the HD machine (Fig. 10). Stop the HD machine and return blood volume to the patient with a fluid bolus. If the HD machine is not defibrillation-proof disconnect it from the patient in accordance with the International Electrotechnical Committee (IEC) standards. Keep the dialysis access open so that it can be used for drug administration. The Kidney Disease Outcomes Initiative guideline on the management of cardiovascular disease in dialysis patients recommends that all dialysis units should have on-site capability for external defibrillation.⁵⁵⁷ Most are equipped with AEDs,^{546,558} but manual defibrillators may also be used by trained staff. Dialysis clinics are predominantly nurse-led units. Staff training and confidence may influence the rate of nurse-led defibrillation. When dialysis staff apply the AED before arrival of paramedics, there is a greater proportion of shockable primary arrest rhythms.^{537,546} Given the higher chance of survival with shockable rhythm, avoid delays in defibrillation. All of the reversible causes (4 Hs and 4 Ts) apply to dialysis patients. Electrolyte imbalances and fluid shifts during dialysis are common causes of cardiac arrest. Consider hyperkalaemia if cardiac arrest occurs pre or early in the dialysis session. For management of hyperkalaemic cardiac arrest, refer to

Table 10 – Risk factors for cardiac arrest in haemodialysis patients.

1. Hyperkalaemia
2. Excessive fluid shifts during dialysis
3. The 3-day inter-dialytic interval – ‘weekend break’
 - in-centre dialysis is delivered 3 days per week (Mon / Wed / Fri or Tue / Thu / Sat)
 - highest risk period is just prior to the first session of the week (i.e. Monday or Tuesday) as K⁺ level peak and fluid accumulates
 - risk extends for 12 h after initiation of dialysis on the first session of the week after rapid fluid and electrolyte shifts from peak levels
4. Low potassium dialysate fluid (1 mmol L⁻¹)
5. History of heart disease
6. QT-prolonging medications
7. Non-compliance with diet and / or dialysis regimen

Table 11 – Cardiac arrest during dialysis in out-patient clinics.

Reproduced with permission from UK Kidney Association Hyperkalaemia Guideline in adults.

Study	N=	Before HD	During HD	After HD
Karnik 2001 ⁵³⁸	400	7 %	81 %	12 %
La France 2006 ⁵³⁹	38	8 %	78 %	14 %
Davis 2008 ⁵³⁷	110	10 %	70 %	20 %

Section *Hyperkalaemia*. Consider hypokalaemia if cardiac arrest occurs mid-late or immediately post dialysis.

Dialysis may be required in the early post-resuscitation period, guided by fluid status and serum biochemistry. Patient transfer to an intensive care unit with dialysis facilities is essential.

Prevention of cardiac arrest in dialysis patients

Avoiding hyperkalaemia and volume overload requires patient adherence and careful dialysis prescription, but may reduce the risk of cardiac arrest.^{538,559,560} The use of potassium binders on non-dialysis days to facilitate use of a higher dialysate potassium content (3 mmol) can reduce the occurrence of clinically significant arrhythmias (bradycardia, ventricular tachycardia, and/or asystole).⁵⁶¹ The higher frequency of cardiac arrest during dialysis and the equivalent survival in dialysis patients compared with non-dialysis patients may reflect a higher likelihood of a reversible cause (i.e. electrolyte or fluid disturbance).⁵⁵¹ The highest risk period is the 'weekend break.' It is conceivable that removing this inter-dialytic gap with short, frequent (4–5 per week) dialysis sessions may reduce risk of cardiac death, but this is only achievable with home haemodialysis.

Resuscitation in obese patients

Obesity is defined by WHO as a body mass index exceeding 30 kg m⁻².⁵⁶² Obesity can increase cardiovascular morbidity and mortality directly and indirectly.⁵⁶³ Given the global increase in obesity, an ILCOR scoping review assessed the evidence for cardiac arrest treatment and outcomes in obese patients.⁵⁶⁴ In adults, the effect of obesity on neurological outcome, survival to hospital discharge, longer term survival (months to years), and ROSC was variable. Few studies reported resuscitation quality indicators, and no studies reported adjustments to resuscitation techniques, or provider outcomes. Current evidence indicates that it is reasonable to use standard resuscitation protocols.

Delivery of effective CPR may be challenging in obese patients.⁵⁶⁴ Rescuer fatigue may make it advisable to change the rescuer performing chest compressions more often than every two minutes. The use of mechanical chest compression devices might be considered although body dimensions and slope of the anterior chest wall limit usability of most devices in very obese patients. Obesity is associated with difficult mask ventilation and might pose difficulties with airway management.^{565,566} Experienced providers should secure a patent airway (supraglottic airway devices or tracheal intubation) early to minimise the period of bag-mask ventilation. Although mortality seems to be higher in obese patients than normal-weighted patients receiving ECPR, it should not be withheld in obese patients.⁵⁶⁷

Resuscitation in patients with pectus excavatum

This chest wall deformation (sometimes associated with congenital heart disease) reduces available space for heart and lungs with direct pressure to these organs and increases the risk of cardiac arrest.⁵⁶⁸ The heart is often displaced leftward, which can complicate effective chest compressions. Reports suggest reduced chest compression depth (3–4 cm) to minimise the risk of organ injury in the presence of reduced anteroposterior thoracic diameter.^{569,570}

Surgically placed corrective "Nuss bars" complicate CPR, as they require significantly increased force to deliver proper chest compression depth.⁵⁷¹ Rescuer fatigue and organ damage (liver lacerations specifically) are to be considered when mechanical chest compressions

are delivered.⁵⁷² Defibrillation in the presence of a Nuss bar may divert electrical current flow through the low resistance metallic bar when defibrillation pads are placed anterolaterally.⁵⁷³

Based on very indirect data from case reports, anatomical and simulation studies reporting cardiac arrest in patients with pectus excavatum, the following deviations from the standard algorithm might be considered:

- During CPR, consider a lower chest compression depth of 3–4 cm, because the reduced anteroposterior thoracic diameter limits the effective compression depth achievable and increases the risk of direct trauma to the heart, lungs and great vessels.
- In case of a Nuss bar correction, substantially increased force is required to deliver effective chest compressions because increased chest wall rigidity, potentially accelerating rescuer fatigue.
- Use mechanical chest compressions with caution.
- If chest compressions are ineffective consider early implementation of ECPR to ensure sufficient organ perfusion.
- Use anteroposterior pad placement for defibrillation using standard energies to ensure optimal current flow through the myocardium, despite the presence of the metallic Nuss Bar.⁵⁷⁴

Cardiac arrest in pregnancy

Maternal cardiac arrest refers to cardiac arrest that occurs at any stage in pregnancy or within six weeks after birth. Maternal mortality remains high, with an estimated 287,000 cases globally in 2020 – equivalent to one death every two minutes. The majority occur in low- and middle-income countries. A significant proportion of maternal cardiac arrests occur outside maternity units. Pregnancy is not always immediately apparent, particularly in its early stages. However, considering pregnancy in reproductive-age patients is essential for identifying reversible causes such as concealed ectopic bleeding and modifiable factors such as aortocaval compression after 20 weeks' gestation.⁵⁷⁵

These Guidelines have been informed by an ILCOR scoping review³³⁴ and international guidelines.^{575–583} Most guidance is based on observational data, expert opinion and physiological principles. This update introduces a new process for management of maternal cardiac arrest along with key figures and tables to support clinical application.

US data reveal that 32 % of obstetric patients experiencing cardiac arrest have no history of a pre-existing disorder, 36.1 % had respiratory insufficiency while 33.3 % had hypotension as the most common antecedent conditions. Pulseless electrical activity is the most common rhythm.⁵⁸⁴ While there are limited data on precise causes of maternal cardiac arrest, MBRRACE-UK reports causes of maternal mortality from 2020 to 2022 to be thromboembolism (16 %), COVID-19 (14 %), cardiac disease (13 %), mental health conditions (11 %), sepsis (9 %), epilepsy and stroke (9 %), obstetric bleeding (7 %), early pregnancy disorders (5 %), cancer (3 %), and pre-eclampsia (3 %). Risk factors included Black and Asian ethnicity, social deprivation, maternal age > 35, and obesity.⁵⁷⁵ In contrast, global data from low- and middle-income countries identified haemorrhage (27.1 %), hypertensive disorders (14 %), and sepsis (10.7 %) as the leading direct causes of maternal deaths.⁵⁸⁵

The UK Obstetric Surveillance System identified 66 cases of cardiac arrest in pregnancy in a three year period with a survival rate of 58 %, poor outcomes linked to OHCA, and delays in resuscitative hysterotomy.⁵⁸⁶ The UK NAP7 survey reported an incidence of perioperative

cardiac arrest in obstetric patients of 7.9 per 100,000 anaesthetic encounters and a survival rate of 82 %, with haemorrhage, high neuraxial block, and bradyarrhythmias as the leading causes.⁵⁸⁷

Prevention of maternal cardiac arrest

Maternal cardiac arrest is usually preventable, though in certain causes, such as amniotic fluid embolism, arrest may occur before intervention.⁵⁸⁸ Approach any deteriorating pregnant or peripartum patient following the ABCDE approach. There is evidence to suggest that implementing an obstetric-specific early warning system enables early recognition of deteriorating pregnant patients; however, uptake across Europe is low.^{589–593} Early involvement of an obstetrician, along with simultaneous activation of maternal and neonatal resuscitation teams, is essential.

Aortocaval compression

Aortocaval compression (i.e., compression of the inferior vena cava and the aorta by the gravid uterus) usually manifests by the 20th week of gestation. This may significantly reduce venous return and subsequently cardiac output in the supine position.^{594,595} In healthy pregnant patients with preserved intrinsic compensatory mechanisms, the effects of aortocaval compression may be absent or minimal.⁵⁹⁶ However, in critically ill or hypotensive patients, aortocaval compression may precipitate cardiac arrest and limit the effectiveness of cardiopulmonary resuscitation.

Aortocaval compression may be relieved by manual left uterine displacement or left lateral tilt (Fig. 19), though supporting data are derived from non-cardiac arrest and simulation studies. ILCOR conducted a scoping review on this topic and found insufficient evidence to recommend either method over the other during CPR in pregnant patients.⁸ Magnetic resonance imaging suggests that a left lateral tilt of approximately 30° is needed to partially relieve inferior vena cava compression. This may be achieved by applying lateral tilt on an operating table or by placing wedges under the right hip for patients on a conventional bed.^{597–599} However, wedges are rarely available, and simulation and animal studies have questioned the effectiveness of chest compressions in the left lateral position.^{600,601}

Accordingly, the ERC suggests performing manual left uterine displacement in maternal cardiac arrest. It can be achieved by placing one or both hands below the uterus, on the patient's right side, and pushing upwards and to the left, or if standing on the patient's left, reaching across to cup the uterus from below and lifting upward and leftward (Fig. 19).^{602,603}

Definitive relief of aortocaval compression may be achieved only by resuscitative hysterotomy (sometimes known as “perimortem Caesarean section”).

Resuscitative hysterotomy

Resuscitative hysterotomy should be performed by a skilled team at the site of cardiac arrest to achieve ROSC by relieving aortocaval compression.⁶⁰⁴ Previous guidelines recommended starting the procedure at 4 min and completing uterine evacuation by 5 min.⁶⁰⁵ The maternal cardiac arrest supplement to the ALS algorithm (Table 12) shifts the focus to preparing for resuscitative hysterotomy, highlighting the time-sensitive nature of the intervention.

- The ILCOR review found insufficient evidence to support a specific time for initiating resuscitative hysterotomy.
- Time-sensitive resuscitative hysterotomy depends on clearly designated team competences, rapid system activation, and equipment readiness, all requiring training and rehearsal.^{606,607}
- If arrest results from hypotension despite optimal resuscitation, it is suggested that resuscitative hysterotomy be performed as soon as possible.
- The procedure should not be considered futile beyond 5 min, however, the benefit declines steadily the longer emptying the uterus is delayed.^{586,608}
- In exceptional cases, ECPR may be initiated immediately, with uterine evacuation postponed due to anticoagulation-related bleeding risks.^{609,610}
- In the prehospital setting, the procedure requires adequate access to the patient and a trained clinician, otherwise prioritise time-critical transport to hospital

While fixed timing for resuscitative hysterotomy is de-emphasised, early intervention remains paramount to maximise the chance of return of spontaneous circulation and reduce the risk of hypoxic brain injury.

Extracorporeal life support

A retrospective analysis of peripartum patients requiring ECPR from the International Registry of Extracorporeal Life Support Organization, identified 280 patients,⁶¹¹ with 70 % survival. Survival rates were higher when extracorporeal membranous oxygenation (ECMO) was initiated prior to cardiac arrest.

If available, early ECMO can be considered by the multidisciplinary team in pregnant patients with signs of impending circulatory collapse.




Fig. 19 – Manual left uterine displacement manoeuvre.

Table 12 – Modifications to advanced life support in obstetric patients.

ALS during pregnancy	Rationale
Detection of cardiac arrest: unconscious + abnormal breathing = suspected cardiac arrest	<i>No change</i>
Call for help – “maternal arrest team”	Consider pregnancy in any collapsed woman of childbearing age. Call maternal cardiac arrest team (including an obstetrician and neonatologist)
Manual (left) uterine displacement throughout	Relieve aortocaval compression to improve cardiac output as early as possible and maintain it throughout resuscitation. Establish manual left uterine displacement when two or more team members are available—one performing manual left uterine displacement, the other performing CPR. (Fig. 19)
Chest compression quality parameters and ventilation-compression-ratio	<i>No change</i>
Airway management	Aspiration and failed intubation risk are both increased. Use a stepwise approach (bag-mask, tracheal tube or supraglottic airway if tracheal intubation fails), according to rescuer skills. Try to achieve a ramped position. Intubation should be performed by an experienced person. Equipment considerations: - Short-handled laryngoscope for large breasts - Video laryngoscope as standard for intubation - Smaller tracheal tube with guidance
Defibrillation – Shock energy	<i>No change</i>
Pad position	No change: ensure defibrillation pads are placed beneath, not over, enlarged breast tissue
Fetal monitoring	Remove internal & external fetal monitors before defibrillation
Identify common and reversible causes	See Table 13 on 4Hs, 4Ts and 4Ps
Medications	
Vascular access early	Consider IV/IO access above the diaphragm if possible
Dosage and time of adrenaline, amiodarone and lidocaine	<i>No change</i>
Calcium chloride	10 ml IV calcium chloride 10 %: for Mg overdose, low calcium or hyperkalaemia
Magnesium	2 g IV: for polymorphic VT 4 g IV: for eclampsia
Tranexamic acid	1 g IV: for haemorrhage
Resuscitative Hysterotomy (Perimortem Caesarean section)	In patients > 20 weeks' gestation or fundus above umbilicus. Resuscitative hysterotomy is a time-sensitive intervention. Preparation to perform should be undertaken early. Resuscitative hysterotomy should be performed as soon as possible at the site of cardiac arrest by a skilled team.
Post-resuscitation care	Stabilise mother to stabilise fetus and prepare for major obstetric haemorrhage.

Relieve aortocaval compression. Aortocaval compression should be relieved as early as possible and maintained throughout resuscitation (Fig. 19).

Chest compressions. Currently, there is no strong evidence to support modifying hand position for chest compressions during pregnancy. A magnetic resonance imaging study of cardiac remodelling

in the third trimester of pregnancy showed an increase in left ventricular mass but no cephalad displacement.⁶¹² However, a subsequent transthoracic echocardiography study of left ventricular position during the third trimester demonstrated that it is located approximately 6 cm cranial to the distal tip of the xiphoid process.⁶¹³

With regard to mechanical chest compression devices, there is insufficient evidence to make a recommendation for or against their use, though in certain scenarios, practical alternatives are unavailable – for example, during extrication or transport in the prehospital setting.⁶¹⁴

Airway management. Pregnancy increases the risk of aspiration of gastric contents and difficult intubation.^{615–617} Manage the airway as directed by the ERC Guidelines 2025 ALS (bag-mask, tracheal tube or supraglottic airway if tracheal intubation fails) according to skills of the HCP. Early tracheal intubation facilitates oxygenation and ventilation, while protecting against aspiration, but it is considered an expert skill and should be performed in accordance with current obstetric-anaesthesia guidelines.⁶¹⁷

Defibrillation. When using anterolateral positioning, ensure defibrillation pads are placed beneath, not over, enlarged breast tissue. Defibrillation energy levels are the same as those for non-pregnant adults.⁶¹⁸ To avoid burns, external fetal monitoring should be removed before defibrillation.

Medications. Medications used in the resuscitation of pregnant women are the same as for the general adult population. While there is limited evidence, intravenous or intraosseous access should be established above the level of the diaphragm whenever possible, based on physiological considerations.⁶¹⁹

Reversible causes. The 4Hs and 4Ts are important considerations in all cardiac arrests, including during pregnancy, and may prompt pregnancy-specific diagnoses such as hypovolaemia from a ruptured ectopic pregnancy or pulmonary embolism, the latter being more common in pregnant and postpartum women (Table 13). However, there are also pregnancy-specific causes of cardiac arrest, which are now introduced as the 4Ps:

Pre-eclampsia and eclampsia: Follow established guidelines for management.⁶²⁰ Initiate immediate treatment for severe hypertension, defined as $\geq 160/110$ mmHg. In severe cases give 4 g IV magnesium sulphate over 5 to 15 min to treat and prevent eclampsia, followed by 1 g/h over 24 h.⁶²⁰ Consider pulmonary oedema and treat accordingly. Magnesium toxicity from treatment overdose may cause cardiac arrest (see Table 14).

Puerperal sepsis: In suspected puerperal sepsis, take blood cultures, give early broad-spectrum antibiotics, and consider obstetric sources such as chorioamnionitis, retained products, endometritis, or surgical site infection that may require intervention.

Placental and uterine complications: Examples include placental abruption, placenta praevia or accreta, uterine atony, rupture, inversion, and amniotic fluid embolism. Seek clinical signs of overt or occult haemorrhage (vaginal or intra-abdominal). Examine the abdomen for peritonism and free fluid (use ultrasound if available), assess for hypovolaemia, check haemoglobin concentration and assess clotting function. Request blood products, alert the obstetric team, and prepare for urgent surgical intervention. The management of amniotic fluid embolism is supportive: maintain oxygenation, treat haemorrhage with blood products, correct disseminated intravascu-

Table 13 – Potentially reversible causes of cardiac arrest in obstetric patients: 4Hs, 4Ts, 4Ps.

Potentially reversible causes in pregnancy	Obstetric-specific considerations
Hypoxia	- high risk of aspiration - high risk of difficult airway - reduced lung capacity - higher oxygen demand
Hypovolaemia	- aortocaval compression - haemorrhage (e.g., ectopic pregnancy and other pregnancy-related causes below, DIC) - distributive (e.g., high regional block, anaphylaxis)
Hypo-/hyperkalaemia, other electrolyte disorders	- also consider magnesium levels
Hypo-/hyperthermia	
Thrombosis (coronary and pulmonary)	- high risk for pulmonary embolism - myocardial infarction - coronary and aortic dissection - (amniotic fluid embolism)
Tamponade (cardiac)	
Tension pneumothorax	- nitrous oxide in pre-existing pneumothorax
Toxic agents (poisoning)	- local anaesthetic and other perioperative drugs
Pre-eclampsia and eclampsia	- Follow guidelines - Consider pulmonary oedema - Magnesium overdose may cause cardiac arrest
Puerperal sepsis	- take blood cultures and give early broad-spectrum antibiotics - Chorioamnionitis, retained products, endometritis, - Surgical site infection
Placental and uterine complications	- Placental abruption, placenta praevia, placenta accreta, uterine atony, uterine rupture, uterine inversion, amniotic fluid embolism)
Peripartum cardiomyopathy	- Treat as acute heart failure. - Early echocardiography - Consult cardiology experts

lar coagulation (DIC), and immediately involve obstetric, anaesthetic and critical care teams.

Peripartum cardiomyopathy: Treat as acute heart failure. Consider intravenous diuretics, vasodilators (e.g. nitrates), and inotropes if indicated. Avoid fluid overload. Seek early echocardiography and cardiology input.

Table 14 – Suggested pre-prepared equipment to perform Resuscitative Hysterotomy. We suggest as a minimum a scalpel and 3 cord clamps to be available in all resuscitation carts.

For staff	- Sterile gloves (non-latex) - Gowns - Masks
For mother	- Skin preparation solution - 3 scalpels (1: incision, 2: umbilical cord, 3: backup) - Dissection scissors - Laparotomy sponges - 4 haemostat clamps - Retractor - Gauze (ideally haemostatic)
For baby	- 3 cord clamps - Hat, 4 towels, heated incubator - Equipment to resuscitate the neonate

Cardiovascular disease

Percutaneous coronary intervention is the preferred reperfusion strategy for STEMI in pregnancy.⁵⁸² Consider thrombolysis if timely PCI is unavailable. A review of 200 cases of thrombolysis for massive pulmonary embolism in pregnancy reported maternal mortality rate of 1 % supporting its safe use in pregnancy when clinically indicated.⁶²¹ Point-of-care ultrasound is invaluable in cases where it is unclear if the cause of arrest is due to concealed bleeding or thromboembolism.⁵⁷⁵

Haemorrhage

This is a leading cause of maternal morbidity and mortality.⁶²² Life-threatening haemorrhage can occur both antenatally and postnatally. Ensure that a massive/major obstetric haemorrhage protocol is available in all units, and that it is reviewed regularly and rehearsed in collaboration with all team members including transfusion services and portering staff.⁶²³

Management includes:

- Immediate involvement of obstetric, anaesthetic, and critical care teams who should follow existing guidelines for the management of major obstetric haemorrhage.^{623,624}
- Activation of the major obstetric haemorrhage protocol, with prompt blood transfusion and correction of coagulopathy.
- Administration of tranexamic acid 1 g IV in cases of suspected bleeding.^{625–628}

Maternal considerations for post resuscitation care

Post resuscitation care in pregnancy aligns with standard adult management, with some important additional considerations. Hypothermic temperature control has been reported in isolated cases during early pregnancy, with fetal heart rate monitoring in place. There was no evidence of harm and maternal and neonatal outcomes were favourable.^{629,630} The continued focus should be to resuscitate the mother. Once the mother is stabilised, fetal resuscitation will also be optimised. This includes treating maternal pathology, optimising maternal physiology, and ensuring necessary investigations and imaging. Preparation for major obstetric haemorrhage in anticipation of ROSC is essential following resuscitative hysterotomy or throm-

bolysis, as well as for other maternal cardiac arrest causes such as amniotic fluid embolism.

Maternal critical illness requires a multidisciplinary approach making it essential to involve all relevant specialties, including obstetrics, anaesthesia, critical care and neonatology.⁶³¹

The psychological impact of adverse maternal outcomes should be acknowledged, with support offered to the patient, their family, and the staff involved in their care.

Preparation for cardiac arrest in pregnancy

Gaps in preparedness, including equipment availability and multidisciplinary coordination, remain widespread and may hinder advanced life support in pregnancy, including time-sensitive resuscitative hysterotomy and neonatal resuscitation.⁵⁹³

Pregnant and peripartum women can deteriorate to cardiac arrest anywhere and medical facilities and services must be prepared for such events. There should be^{632–638}:

- Plans and equipment in place for resuscitation of both the pregnant woman and newborn (see Table 14).
- Early involvement of obstetric, anaesthetic, critical care, and neonatal teams.
- Regular multidisciplinary training in obstetric emergencies, using simulated resuscitation scenarios.
- Review and debriefing following clinical events to support learning and team performance

Collaborator

We would like to thank Ryan Aird, cardiac arrest survivor, Member of SCAUK (Sudden Cardiac Arrest UK) for his comments and input while developing these ERC Guidelines 2025 Special Circumstances in Resuscitation.

Disclosures

Declarations of competing interests for all ERC Guidelines authors are displayed in a COI table which can be found online at <https://doi.org/10.1016/j.resuscitation.2025.110753>.

Author details

^aDepartment of Anesthesiology, University Medical Centre, Johannes-Gutenberg University, Mainz, Germany ^bCardiopulmonary Resuscitation Lab, School of Medicine, University of Crete, Heraklion, Crete, Greece ^cDepartment of Anesthesiology, Onassis Hospital, Athens, Greece ^dFaculty of Education Sciences and CLINURSID Research Group, Universidade de Santiago de Compostela, Santiago de Compostela, Spain ^eSimulation and Intensive Care Unit of Santiago (SICRUS) Research Group, Health Research Institute of Santiago, University Hospital of Santiago de Compostela-CHUS, Santiago de Compostela, Spain ^fDepartments of Renal and Internal Medicine, Victoria Hospital, Kirkcaldy, Fife, UK ^gExtreme Environments Laboratory, School of Sport, Health and Exercise Science, University of Portsmouth, UK ^hMaatschappij tot Redding van Drenkelingen, Amsterdam, the Netherlands ⁱNottingham University Hospitals NHS Trust, Nottingham, UK ^jUniv. Grenoble Alpes, Emergency Department

and Mobile Intensive Care Unit, CNRS, UMR 5525, TIMC, 38000 Grenoble, France ^kMaccabi Healthcare Services and Hebrew University, Faculty of Medicine, Israel ^lUniversity Medical Center Schleswig-Holstein, Institute for Emergency and Rescue Medicine, Kiel, Germany ^mCardiology Department, University Clinical Hospital of Santiago de Compostela, Spain ⁿInstitute of Health Research of Santiago de Compostela (IDIS), Biomedical Research Networking Centres on Cardiovascular Disease (CIBER-CV), A Coruna, Spain ^oFaculty of Medicine, University of Bern, Switzerland ^pDepartment of Anaesthesia, Intensive Care Medicine, Emergency Medicine and Pain Medicine, University Medicine Greifswald, Germany ^qDepartment of Anesthesiology and Intensive Care Medicine, University Hospital Ulm, Germany ^rInstitute of Sports and Preventive Medicine, Saarland University, Saarbrücken, Germany ^sDepartment of Anaesthesiology and Intensive Care Medicine, St. John of God Hospital, Paracelsus Medical University, Salzburg, Austria ^tDepartment of Anaesthesia and Intensive Care Medicine, Charles University, Third Faculty of Medicine and FNKV University Hospital in Prague, Czech Republic ^uDepartment of Cardiovascular Sciences, Fondazione Policlinico Universitario A. Gemelli, IRCCS, Italy ^vInstitute of Anesthesiology and Intensive Care, Catholic University of the Sacred Heart, Italy ^wDepartment of Emergency Medicine, University Hospital of Heraklion, Crete, Greece ^xBielefeld University, Medical School and University Medical Center OWL, EvKB, Department of Anesthesiology, Intensive Care, Emergency Medicine, Transfusion Medicine and Pain Therapy, Bielefeld, Germany ^yEmergency Medical Services of the Hradec Králové Region, Hradec Králové, Czech Republic ^zDepartment of Anaesthesiology and Intensive Care Medicine, Charles University in Prague, University Hospital Hradec Králové, Czech Republic ^{aa}Department of Anaesthesiology and Intensive Care Medicine, Charles University, Faculty of Medicine in Hradec Králové, Czech Republic ^{ab}Department of Anaesthetics, University Hospital Southampton, Southampton SO168AD, UK

REFERENCES

- Berg KM, Bray JE, Djarv T, et al. Executive Summary: 2025 International Liaison Committee on Resuscitation Consensus on Science With Treatment Recommendations. *Resuscitation* 2025;215 (Suppl 2):110805.
- Greif RLKG, Djarv T, Ek JE, et al. European resuscitation council guidelines 2025: executive summary. *Resuscitation* 2025;215 (Suppl 1):110770.
- Djarv TRJ, Semeraro F, Bräde L, et al. European Resuscitation Council Guidelines 2025: First Aid. *Resuscitation* 2025;215 (Suppl 1):110752.
- Panesar SS, Javad S, de Silva D, et al. The epidemiology of anaphylaxis in Europe: a systematic review. *Allergy* 2013;68 (11):1353–61. <https://doi.org/10.1111/all.12272>.
- Aurich S, Dölle-Bierke S, Francuzik W, et al. Anaphylaxis in elderly patients—data from the European Anaphylaxis Registry. *Front Immunol* 2019;10:750.
- Francuzik W, Ruëff F, Bauer A, et al. Phenotype and risk factors of venom-induced anaphylaxis: a case-control study of the European Anaphylaxis Registry. *J Allergy Clin Immunol* 2021;147(2):653–662. e9. <https://doi.org/10.1016/j.jaci.2020.06.008>.
- Turner PJ, Gowland MH, Sharma V, et al. Increase in anaphylaxis-related hospitalizations but no increase in fatalities: an analysis of United Kingdom national anaphylaxis data, 1992–2012. *J Allergy Clin Immunol* 2015;135:956–963.e1.
- Greif R, Bray JE, Djarv T, et al. 2024 international consensus on cardiopulmonary resuscitation and emergency cardiovascular care science with treatment recommendations: summary from the basic life support; advanced life support; pediatric life support; neonatal life support; education, implementation, and teams; and first aid task forces. *Resuscitation* 2024;205. <https://doi.org/10.1016/j.resuscitation.2024.110414>.
- Cardona V, Ansotegui IJ, Ebisawa M, et al. World allergy organization anaphylaxis guidance 2020. *World Allergy Organ J* 2020;13(10):100472. <https://doi.org/10.1016/j.waojou.2020.100472>.
- Muraro A, Worm M, Alviani C, et al. EAACI guidelines: anaphylaxis (2021 update). *Allergy* 2022;77(2):357–77. <https://doi.org/10.1111/all.15032>.
- Golden DBK, Wang J, Waserman S, et al. Anaphylaxis: a 2023 practice parameter update. *Ann Allergy Asthma Immunol* 2024;132(2):124–76. <https://doi.org/10.1016/j.anaai.2023.09.015>.
- (ASCI) ASoClaA. Australasian Society of Clinical Immunology and Allergy (ASCI). (https://www.allergy.org.au/images/ASCI_HP_Guidelines_Acute_Management_Anaphylaxis_2024.pdf).
- Cook TM, Kane AD, Armstrong RA, Kursumovic E, Soar J. Perioperative cardiac arrest due to suspected anaphylaxis as reported to the 7th national audit project of the royal college of anaesthetists. *Anaesthesia* 2024;79(5):498–505. <https://doi.org/10.1111/anae.16229>.
- Brown SGA. Cardiovascular aspects of anaphylaxis: implications for treatment and diagnosis. *Current Opinion in Allergy and Clinical Immunology* 2005;5(4):359–64. <https://doi.org/10.1097/01.all.0000174158.78626.35>.
- Pumphrey RS. Fatal posture in anaphylactic shock. *J Allergy Clin Immunol* 2003;112(2):451–2. <https://doi.org/10.1067/mai.2003.1614>.
- Turner PJ, Jerschow E, Umasunthar T, Lin R, Campbell DE, Boyle RJ. Fatal anaphylaxis: mortality rate and risk factors. *J Allergy Clin Immunol Pract* 2017;5(5):1169–78. <https://doi.org/10.1016/j.jaip.2017.06.031>.
- Cardona V, Ferré-Ybarz L, Guilarte M, et al. Safety of adrenaline use in anaphylaxis: a multicentre register. *Int Arch Allergy Immunol* 2017;173(3):171–7. <https://doi.org/10.1159/000477566>.
- Simons FE, Gu X, Simons KJ. Epinephrine absorption in adults: intramuscular versus subcutaneous injection. *J Allergy Clin Immunol* 2001;108(5):871–3. <https://doi.org/10.1067/mai.2001.119409>.
- Carlson JN, Cook S, Djarv T, Woodin JA, Singletary E, Zideman DA. Second dose of epinephrine for anaphylaxis in the first aid setting: a scoping review. *Cureus* 2020;12(11):e11401. <https://doi.org/10.7759/cureus.11401>.
- Campbell RL, Bellolio MF, Knutson BD, et al. Epinephrine in anaphylaxis: higher risk of cardiovascular complications and overdose after administration of intravenous bolus epinephrine compared with intramuscular epinephrine. *J Allergy Clin Immunol Pract* 2015;3:76–80.
- Ring J, Beyer K, Biedermann T, et al. Guideline (S2k) on acute therapy and management of anaphylaxis: 2021 update. *Allergo J Int* 2021;30(1):1–25. <https://doi.org/10.1007/s40629-020-00158-y>.
- Boswell B, Rudders SA, Brown JC. Emerging therapies in anaphylaxis: alternatives to intramuscular administration of epinephrine. *Curr Allergy Asthma Rep* 2021;21(3):18. <https://doi.org/10.1007/s11882-021-00994-0>.
- Dewachter P, Savic L. Perioperative anaphylaxis: pathophysiology, clinical presentation and management. *BJA Educ* 2019;19 (10):313–20. <https://doi.org/10.1016/j.bjae.2019.06.002>.
- Vincent JL, De Backer D. Circulatory shock. *N Engl J Med* 2013;369 (18):1726–34. <https://doi.org/10.1056/NEJMr1208943>.

25. O'Driscoll BR, Howard LS, Earis J, Mak V. BTS guideline for oxygen use in adults in healthcare and emergency settings. *Thorax* 2017;72 (Suppl 1):ii1–ii90. <https://doi.org/10.1136/thoraxjnl-2016-209729>.
26. Ellis BC, Brown SG. Parenteral antihistamines cause hypotension in anaphylaxis. *Emerg Med Australas* 2013;25(1):92–3. <https://doi.org/10.1111/1742-6723.12028>.
27. Alqurashi W, Ellis AK. Do corticosteroids prevent biphasic anaphylaxis?. *J Allergy Clin Immunol Pract* 2017;5(5):1194–205. <https://doi.org/10.1016/j.jaip.2017.05.022>.
28. Liyanage CK, Galappathay P, Seneviratne SL. Corticosteroids in management of anaphylaxis; a systematic review of evidence. *Eur Ann Allergy Clin Immunol* 2017;49(5):196–207. <https://doi.org/10.23822/EurAnnACI.1764-1489.15>.
29. Delli Colli L, Al Ali A, Gabrielli S, et al. Managing anaphylaxis: epinephrine, antihistamines, and corticosteroids: more than 10 years of Cross-Canada Anaphylaxis REgistry data. *Ann Allergy Asthma Immunol* 2023;131(6):752–758.e1. <https://doi.org/10.1016/j.anai.2023.08.606>.
30. Gabrielli S, Clarke A, Morris J, et al. Evaluation of prehospital management in a Canadian emergency department anaphylaxis cohort. *J Allergy Clin Immunol Pract* 2019;7(7):2232–2238.e3. <https://doi.org/10.1016/j.jaip.2019.04.018>.
31. Tafesse E, Hurst M, Sugrue D, et al. Serum potassium as a predictor of adverse clinical outcomes in patients with increasing comorbidity burden. *Eur Heart J Qual Care Clin Outcomes* 2022;8 (1):61–9. <https://doi.org/10.1093/ehjqcco/qcaa078>.
32. Jessen MK, Andersen LW, Djakow J, et al. Pharmacological interventions for the acute treatment of hyperkalaemia: a systematic review and meta-analysis. *Resuscitation* 2025;110489. <https://doi.org/10.1016/j.resuscitation.2025.110489>.
33. Conway R, Creagh D, Byrne DG, O'Riordan D, Silke B. Serum potassium levels as an outcome determinant in acute medical admissions. *Clin Med (Lond)* 2015;15(3):239–43. <https://doi.org/10.7861/clinmedicine.15-3-239>.
34. Einhorn LM, Zhan M, Hsu VD, et al. The frequency of hyperkalemia and its significance in chronic kidney disease. *Arch Intern Med* 2009;169(12):1156–62. <https://doi.org/10.1001/archinternmed.2009.132>.
35. Yu Y, Vangaveti VN, Schnetler RJ, Crowley BJ, Mallett AJ. Hyperkalaemia among hospital admissions: prevalence, risk factors, treatment and impact on length of stay. *BMC Nephrol* 2024;25(1):454. <https://doi.org/10.1186/s12882-024-03863-w>.
36. Agiro A, Mu F, Cook E, et al. Hyperkalemia and the risks of adverse cardiovascular outcomes in patients with chronic kidney disease. *J Am Heart Assoc* 2025;14(1)e035256. <https://doi.org/10.1161/JAHA.124.035256>.
37. Logan Ellis H, Al-Agil M, Kelly PA, Teo J, Sharpe C, Whyte MB. The burden of hyperkalaemia on hospital healthcare resources. *Clin Exp Med* 2024;24(1):190. <https://doi.org/10.1007/s10238-024-01452-7>.
38. Fu EL, Evans M, Clase CM, et al. Stopping renin-angiotensin system inhibitors in patients with advanced CKD and risk of adverse outcomes: a nationwide study. *J Am Soc Nephrol* 2021;32 (2):424–35. <https://doi.org/10.1681/ASN.2020050682>.
39. Svensson MK, Murohara T, Lesen E, et al. Hyperkalaemia-related reduction of RAASi treatment associates with more subsequent inpatient care. *Nephrol Dial Transplant* 2024;39(8):1258–67. <https://doi.org/10.1093/ndt/qfae016>.
40. Yang A, Shi M, Lau ESH, et al. Clinical outcomes following discontinuation of renin-angiotensin-system inhibitors in patients with type 2 diabetes and advanced chronic kidney disease: a prospective cohort study. *EClinicalMedicine* 2023;55:101751. <https://doi.org/10.1016/j.eclinm.2022.101751>.
41. Kosiborod MN, Cherney DZI, Desai AS, et al. Sodium zirconium cyclosilicate for management of hyperkalemia during spironolactone optimization in patients with heart failure. *J Am Coll Cardiol* 2024. <https://doi.org/10.1016/j.jacc.2024.11.014>.
42. Pitt B, Anker SD, Lund LH, et al. Patiromer facilitates angiotensin inhibitor and mineralocorticoid antagonist therapies in patients with heart failure and hyperkalemia. *J Am Coll Cardiol* 2024;84 (14):1295–308. <https://doi.org/10.1016/j.jacc.2024.05.079>.
43. Chothia MY, Kassum P, Zemlin A. A method comparison study of a point-of-care blood gas analyser with a laboratory auto-analyser for the determination of potassium concentrations during hyperkalaemia in patients with kidney disease. *Biochem Med (Zagreb)* 2020;30(3):030702. <https://doi.org/10.11613/BM.2020.030702>.
44. Ismail MH, Baharuddin KA, Suliman MA, Mohd Shukri MF, Che Has SN, Lo ZZ. Hyperkalemia measurement between blood gas analyser and main laboratory biochemistry analyser. *Med J Malaysia* 2021;76(2):157–63. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/33742622>.
45. Mahmoud H, Jaffar Z, Al Alawi YM, et al. Accuracy of potassium measurement using blood gas analyzer. *Cureus* 2022;14(3)e23653. <https://doi.org/10.7759/cureus.23653>.
46. Durfey N, Lehnhof B, Bergeson A, et al. Severe hyperkalemia: can the electrocardiogram risk stratify for short-term adverse events?. *West J Emerg Med* 2017;18(5):963–71. <https://doi.org/10.5811/westjem.2017.6.33033>.
47. Littmann L, Gibbs MA. Electrocardiographic manifestations of severe hyperkalemia. *J Electrocardiol* 2018;51(5):814–7. <https://doi.org/10.1016/j.jelectrocard.2018.06.018>.
48. Batterink J, Cessford TA, Taylor RAI. Pharmacological interventions for the acute management of hyperkalaemia in adults. *Cochrane Database of Systematic Reviews* 2015. <https://doi.org/10.1002/14651858.CD010344.pub2>.
49. Mahoney BA, Smith WAD, Lo DS, Tsoi K, Tonelli M, Clase CM. Emergency interventions for hyperkalaemia - art. no. CD003235. pub2. *Cochrane Database of Systematic Reviews* 2005(2) (In English). doi: ARTN CD003235. 10.1002/14651858.CD003235.pub2.
50. Alfonzo A, Harrison A, Baines R, Chu A. Treatment of acute hyperkalaemia in adults. UK Kidney Association (UKKA) Guideline 2023. Available from: <http://www.ukkidney.org>.
51. Cook ME, Tran LK, DeGrado JR, Alkazemi A, Marino KK. Evaluation of insulin dosing strategies for hyperkalemia management at an academic medical center. *Clin Ther* 2024;46 (5):382–8. <https://doi.org/10.1016/j.clinthera.2024.03.003>.
52. Finder SN, McLaughlin LB, Dillon RC. 5 versus 10 units of intravenous insulin for hyperkalemia in patients with moderate renal dysfunction. *J Emerg Med* 2022;62(3):298–305. <https://doi.org/10.1016/j.jemermed.2021.10.027>.
53. Garcia J, Pintens M, Morris A, Takamoto P, Baumgartner L, Tasaka CL. Reduced versus conventional dose insulin for hyperkalemia treatment. *J Pharm Pract* 2018897190018799220. <https://doi.org/10.1177/0897190018799220>.
54. Moussavi K, Nguyen LT, Hua H, Fitter S. Comparison of IV insulin dosing strategies for hyperkalemia in the emergency department. *Crit Care Explor* 2020;2(4)e0092. <https://doi.org/10.1097/CCE.000000000000092>.
55. Pearson SC, O'Connor K, Keller K, Hodge TJ, Nesbit R. Efficacy of standard- vs reduced-dose insulin for treatment of hyperkalemia: a quasi-experiment. *Am J Health Syst Pharm* 2022;79 (Suppl 1): S13–20. <https://doi.org/10.1093/ajhp/zxab382>.
56. Lim AKH, Crnobraja L, Metlapalli M, Govinna M, Jiang C. The effect of patient factors and cotreatments on the magnitude of potassium lowering with insulin-glucose treatment in patients with hyperkalemia. *Epidemiologia (Basel)* 2021;2(1):27–35. <https://doi.org/10.3390/epidemiologia2010003>.
57. Apel J, Reutrakul S, Baldwin D. Hypoglycemia in the treatment of hyperkalemia with insulin in patients with end-stage renal disease. *Clin Kidney J* 2014;7(3):248–50. <https://doi.org/10.1093/ckj/sfu026>.

58. Boughton CK, Dixon D, Goble E, et al. Preventing hypoglycemia following treatment of hyperkalemia in hospitalized patients. *J Hosp Med* 2019;14(5):284–7. <https://doi.org/10.12788/jhm.3145>.
59. Coca A, Valencia AL, Bustamante J, Mendiluce A, Floege J. Hypoglycemia following intravenous insulin plus glucose for hyperkalemia in patients with impaired renal function. *PLoS One* 2017;12(2):e0172961. <https://doi.org/10.1371/journal.pone.0172961>.
60. LaRue HA, Peksa GD, Shah SC. A comparison of insulin doses for the treatment of hyperkalemia in patients with renal insufficiency. *Pharmacotherapy* 2017;37(12):1516–22. <https://doi.org/10.1002/phar.2038>.
61. Pierce DA, Russell G, Pirkle Jr JL. Incidence of hypoglycemia in patients with low eGFR treated with insulin and dextrose for hyperkalemia. *Ann Pharmacother* 2015;49(12):1322–6. <https://doi.org/10.1177/1060028015607559>.
62. Scott NL, Klein LR, Cales E, Driver BE. Hypoglycemia as a complication of intravenous insulin to treat hyperkalemia in the emergency department. *Am J Emerg Med* 2019;37(2):209–13. <https://doi.org/10.1016/j.ajem.2018.05.016>.
63. Chothia MY, Humphrey T, Schoonees A, Chikte UME, Davids MR. Hypoglycaemia due to insulin therapy for the management of hyperkalaemia in hospitalised adults: a scoping review. *PLoS One* 2022;17(5):e0268395. <https://doi.org/10.1371/journal.pone.0268395>.
64. Crnobrnja L, Metlapalli M, Jiang C, Govinna M, Lim AKH. The association of insulin-dextrose treatment with hypoglycemia in patients with hyperkalemia. *Sci Rep* 2020;10(1):22044. <https://doi.org/10.1038/s41598-020-79180-7>.
65. Humphrey TJL, James G, Wilkinson IB, Hiemstra TF. Clinical outcomes associated with the emergency treatment of hyperkalaemia with intravenous insulin-dextrose. *Eur J Intern Med* 2022;95:87–92. <https://doi.org/10.1016/j.ejim.2021.09.018>.
66. Kijprasert W, Tarudeeyathaworn N, Loketkrawee C, Pimpaporn T, Pattarasettaseranee P, Tangsuwanaruk T. Predicting hypoglycemia after treatment of hyperkalemia with insulin and glucose (Glu-K60 score). *BMC Emerg Med* 2022;22(1):179. <https://doi.org/10.1186/s12873-022-00748-9>.
67. Tee SA, Devine K, Potts A, et al. Iatrogenic hypoglycaemia following glucose-insulin infusions for the treatment of hyperkalaemia. *Clin Endocrinol (Oxf)* 2021;94(2):176–82. <https://doi.org/10.1111/cen.14343>.
68. Ibarra Jr F, Fountain C, Fallert T. Current hyperkalemia interventions co-administered with a dextrose 10% solution significantly lower hypoglycemic rates (CHICA-D10). *Am J Emerg Med* 2024;84:120–3. <https://doi.org/10.1016/j.ajem.2024.07.061>.
69. Allon M, Copkney C. Albuterol and insulin for treatment of hyperkalemia in hemodialysis patients. *Kidney International* 1990;38(5):869–72. <https://doi.org/10.1038/ki.1990.284>.
70. Ahmed J, Weisberg LS. Hyperkalemia in dialysis patients. *Semin Dialysis* 2001;14(5):348–56. <https://doi.org/10.1046/j.1525-139X.2001.00087.x> [In English].
71. Vallentin MF, Granfeldt A, Meilandt C, et al. Effect of intravenous or intraosseous calcium vs saline on return of spontaneous circulation in adults with out-of-hospital cardiac arrest: a randomized clinical trial. *JAMA* 2021;326(22):2268–76. <https://doi.org/10.1001/jama.2021.20929>.
72. Piktet JS, Wan X, Kouk S, Laurita KR, Wilson LD. Beneficial effect of calcium treatment for hyperkalemia is not due to “membrane stabilization”. *Crit Care Med* 2024;52(10):1499–508. <https://doi.org/10.1097/CCM.0000000000006376>.
73. Fishel Bartal M, Sibai BM. Eclampsia in the 21st century. *Am J Obstet Gynecol* 2022;226(2S):S1237–53. <https://doi.org/10.1016/j.ajog.2020.09.037>.
74. Kosiborod M, Rasmussen HS, Lavin P, et al. Effect of sodium zirconium cyclosilicate on potassium lowering for 28 days among outpatients with hyperkalemia: the HARMONIZE randomized clinical trial. *JAMA* 2014;312(21):2223–33. <https://doi.org/10.1001/jama.2014.15688>.
75. Wallmuller C, Meron G, Kurkciyan I, Schober A, Stratil P, Sterz F. Causes of in-hospital cardiac arrest and influence on outcome. *Resuscitation* 2012;83(10):1206–11. <https://doi.org/10.1016/j.resuscitation.2012.05.001>.
76. Wang CH, Huang CH, Chang WT, et al. The effects of calcium and sodium bicarbonate on severe hyperkalaemia during cardiopulmonary resuscitation: a retrospective cohort study of adult in-hospital cardiac arrest. *Resuscitation* 2016;98:105–11. <https://doi.org/10.1016/j.resuscitation.2015.09.384>.
77. Saarinen S, Nurmi J, Toivio T, Fredman D, Virkkunen I, Castrén M. Does appropriate treatment of the primary underlying cause of PEA during resuscitation improve patients' survival? *Resuscitation* 2012;83(7):819–22. <https://doi.org/10.1016/j.resuscitation.2011.12.018>.
78. Chiu CC, Yen HH, Chen YL, Siao FY. Severe hyperkalemia with refractory ventricular fibrillation: successful resuscitation using extracorporeal membrane oxygenation. *Am J Emerg Med* 2014;32(8):943.e5–6. <https://doi.org/10.1016/j.ajem.2014.01.016>.
79. Kao KC, Huang CC, Tsai YH, Lin MC, Tsao TC. Hyperkalemic cardiac arrest successfully reversed by hemodialysis during cardiopulmonary resuscitation: case report. *Chang Gung Med J* 2000;23(9):555–9. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/11092145>.
80. Kose N, Bilgin F. Successful treatment of a patient with cardiac arrest due to hyperkalemia by prolonged cardiopulmonary resuscitation along with hemodialysis: a case report and review of the literature. *Medicina (Kaunas)* 2021;57(8). <https://doi.org/10.3390/medicina57080810>.
81. Tijssen JA, Filler G. When CRRT on ECMO is not enough for potassium clearance: a case report. *Can J Kidney Health Dis* 2017;4:2054358117722559. <https://doi.org/10.1177/2054358117722559>.
82. Lee GT, Jeong D, Park JE, et al. Efficacy of anti-hyperkalemic agents during cardiopulmonary resuscitation in out-of-hospital cardiac arrest. *Heliyon* 2024;10(16):e36345. <https://doi.org/10.1016/j.heliyon.2024.e36345>.
83. Kardalas E, Paschou SA, Anagnostis P, Muscogiuri G, Siasos G, Vryonidou A. Hypokalemia: a clinical update. *Endocr Connect* 2018;7(4):R135–46. <https://doi.org/10.1530/EC-18-0109>.
84. Tse G, Li KHC, Cheung CKY, et al. Arrhythmogenic mechanisms in hypokalaemia: insights from pre-clinical models. *Front Cardiovasc Med* 2021;8:620539. <https://doi.org/10.3389/fcvm.2021.620539>.
85. Fan Y, Wu M, Li X, et al. Potassium levels and the risk of all-cause and cardiovascular mortality among patients with cardiovascular diseases: a meta-analysis of cohort studies. *Nutr J* 2024;23(1):8. <https://doi.org/10.1186/s12937-023-00888-z>.
86. Lima I, Nunes JT, de Oliveira IH, et al. Association of potassium disorders with the mode of death and etiology in patients with chronic heart failure: the INCOR-HF study. *Sci Rep* 2024;14(1):30167. <https://doi.org/10.1038/s41598-024-74928-x>.
87. Hagengaard L, Sogaard P, Espersen M, et al. Association between serum potassium levels and short-term mortality in patients with atrial fibrillation or flutter co-treated with diuretics and rate- or rhythm-controlling drugs. *Eur Heart J Cardiovasc Pharmacother* 2020;6(3):137–44. <https://doi.org/10.1093/ehjcvp/pvz024>.
88. Ravn Jacobsen M, Jabbari R, Glinge C, et al. Potassium disturbances and risk of ventricular fibrillation among patients with st-segment-elevation myocardial infarction. *J Am Heart Assoc* 2020;9(4):e014160. <https://doi.org/10.1161/JAHA.119.014160>.
89. Huang BP, Zhao L, Zhao XM, et al. Serum potassium levels and mortality in hospitalized heart failure patients. *Rev Cardiovasc Med* 2023;24(8):228. <https://doi.org/10.31083/j.rcm.2408228>.
90. Cohn JN, Kowey PR, Whelton PK, Prisant LM. New guidelines for potassium replacement in clinical practice: a contemporary review by the national council on potassium in clinical practice. *Arch Intern*

- Med 2000;160(16):2429–36. <https://doi.org/10.1001/archinte.160.16.2429>.
91. Huang CL, Kuo E. Mechanism of hypokalemia in magnesium deficiency. *J Am Soc Nephrol* 2007;18(10):2649–52. <https://doi.org/10.1681/ASN.2007070792>.
 92. Epstein Y, Yanovich R. Heatstroke. *N Engl J Med* 2019;380(25):2449–59. <https://doi.org/10.1056/NEJMra1810762>.
 93. Hayashida K, Shimizu K, Yokota H. Severe heatwave in Japan. *Acute Med Surg* 2019;6(2):206–7. <https://doi.org/10.1002/ams2.387>.
 94. Hopkins PM, Girard T, Dalay S, et al. Malignant hyperthermia 2020: guideline from the association of anaesthetists. *Anaesthesia* 2021;76(5):655–64. <https://doi.org/10.1111/anae.15317>.
 95. Rüffert H, Bastian B, Bendixen D, et al. Consensus guidelines on perioperative management of malignant hyperthermia suspected or susceptible patients from the European malignant hyperthermia group. *Br J Anaesth* 2021;126(1):120–30. <https://doi.org/10.1016/j.bja.2020.09.029>.
 96. Gong X. Malignant hyperthermia when dantrolene is not readily available. *BMC Anesthesiol* 2021;21(1):119. <https://doi.org/10.1186/s12871-021-01328-3>.
 97. Cong Z, Wan T, Wang J, et al. Epidemiological and clinical features of malignant hyperthermia: a scoping review. *Clin Genet* 2024;105(3):233–42. <https://doi.org/10.1111/cge.14475>.
 98. Eifling KP, Gaudio FG, Dumke C, et al. Wilderness medical society clinical practice guidelines for the prevention and treatment of heat illness: 2024 update. *Wilderness Environ Med* 2024;35(1_suppl):112s–27s. <https://doi.org/10.1177/10806032241227924>.
 99. Lott C, Truhlář A, Alfonzo A, et al. European resuscitation council guidelines 2021: cardiac arrest in special circumstances. *Resuscitation* 2021;161:152–219. <https://doi.org/10.1016/j.resuscitation.2021.02.011>.
 100. Singletary EM, Zideman DA, Bendall JC, et al. 2020 international consensus on first aid science with treatment recommendations. *Resuscitation* 2020;156:A240–82. <https://doi.org/10.1016/j.resuscitation.2020.09.016>.
 101. Zideman DA, Singletary EM, Borra V, et al. European resuscitation council guidelines 2021: first aid. *Resuscitation* 2021;161:270–90. <https://doi.org/10.1016/j.resuscitation.2021.02.013>.
 102. Roberts WO, Armstrong LE, Sawka MN, Yeargin SW, Heled Y, O'Connor FG. ACSM expert consensus statement on exertional heat illness: recognition, management, and return to activity. *Curr Sports Med Rep* 2023;22(4):134–49. <https://doi.org/10.1249/jsr.0000000000001058>.
 103. Strapazzon G, Procter E, Paal P, Brugger H. Pre-hospital core temperature measurement in accidental and therapeutic hypothermia. *High Alt Med Biol* 2014;15(2):104–11. <https://doi.org/10.1089/ham.2014.1008>.
 104. Barletta JF, Palmieri TL, Toomey SA, et al. Society of critical care medicine guidelines for the treatment of heat stroke. *Crit Care Med* 2025;53(2):e490–500. <https://doi.org/10.1097/ccm.0000000000006551>.
 105. Douma MJ, Aves T, Allan KS, et al. First aid cooling techniques for heat stroke and exertional hyperthermia: a systematic review and meta-analysis. *Resuscitation* 2020;148:173–90. <https://doi.org/10.1016/j.resuscitation.2020.01.007>.
 106. Hew-Butler T, Rosner MH, Fowkes-Godek S, et al. Statement of the 3rd international exercise-associated hyponatremia consensus development conference, Carlsbad, California, 2015. *Br J Sports Med* 2015;49(22):1432–46. <https://doi.org/10.1136/bisports-2015-095004>.
 107. Bouchama A, Dehbi M, Chaves-Carballo E. Cooling and hemodynamic management in heatstroke: practical recommendations. *Crit Care* 2007;11(3):R54. <https://doi.org/10.1186/cc5910>.
 108. Litman RS, Griggs SM, Dowling JJ, Riazi S. Malignant hyperthermia susceptibility and related diseases. *Anesthesiology* 2018;128(1):159–67. <https://doi.org/10.1097/aln.0000000000001877>.
 109. Riazi S, Kraeva N, Hopkins PM. Malignant hyperthermia in the post-genomics era: new perspectives on an old concept. *Anesthesiology* 2018;128(1):168–80. <https://doi.org/10.1097/aln.0000000000001878>.
 110. Hall AP, Henry JA. Acute toxic effects of 'Ecstasy' (MDMA) and related compounds: overview of pathophysiology and clinical management. *Br J Anaesth* 2006;96(6):678–85. <https://doi.org/10.1093/bja/ael078>.
 111. Kollmann-Camaïora A, Alsina E, Domínguez A, et al. Clinical protocol for the management of malignant hyperthermia. *Rev Esp Anestesiol Reanim* 2017;64(1):32–40. <https://doi.org/10.1016/j.redar.2016.06.004> [In eng spa].
 112. Centres accredited by the European Malignant Hyperthermia Group.
 113. Kyotani Y, Zhao J, Nakahira K, Yoshizumi M. The role of antipsychotics and other drugs on the development and progression of neuroleptic malignant syndrome. *Sci Rep* 2023;13(1):18459. <https://doi.org/10.1038/s41598-023-45783-z>.
 114. Ngo V, Guerrero A, Lanum D, et al. Emergent treatment of neuroleptic malignant syndrome induced by antipsychotic monotherapy using dantrolene. *Clin Pract Cases Emerg Med* 2019;3(1):16–23. <https://doi.org/10.5811/cpcem.2018.11.39667>.
 115. Rosenberg MR, Green M. Neuroleptic malignant syndrome: review of response to therapy. *Arch Int Med* 1989;149(9):1927–31. <https://doi.org/10.1001/archinte.1989.00390090009002>.
 116. Wijdicks EFM, Ropper AH. Neuroleptic malignant syndrome. *New England Journal of Medicine* 2024;391(12):1130–8. <https://doi.org/10.1056/NEJMra2404606>.
 117. Zhang Y, Deng W, Wang M, Luo S, Li S. A real-world pharmacovigilance study of neuroleptic malignant syndrome based on FDA adverse event reporting system. *Front Pharmacol* 2024;15. <https://doi.org/10.3389/fphar.2024.1438661> (Original Research) [In English].
 118. Chiew AL, Isbister GK. Management of serotonin syndrome (toxicity). *British Journal of Clinical Pharmacology*;n/a(n/a). <https://doi.org/10.1111/bcp.16152>.
 119. Gillman PK. Serotonin syndrome treated with chlorpromazine. *J Clin Psychopharmacol* 1997;17(2):128–9. <https://doi.org/10.1097/00004714-199704000-00017>.
 120. Mason PJ, Morris VA, Balcezak TJ. Serotonin syndrome. Presentation of 2 cases and review of the literature. *Medicine (Baltimore)* 2000;79(4):201–9. <https://doi.org/10.1097/00005792-200007000-00001>.
 121. Morarasu BC, Coman AE, Bologa C, et al. Recognition and management of serotonin toxidrome in the emergency department-case based review. *J Pers Med* 2022;12(12). <https://doi.org/10.3390/jpm12122069>.
 122. Nguyen H, Pan A, Smollin C, Cantrell LF, Kearney T. An 11-year retrospective review of cyproheptadine use in serotonin syndrome cases reported to the California Poison Control System. *J Clin Pharm Ther* 2019;44(2):327–34. <https://doi.org/10.1111/jcpt.12796>.
 123. Boehrer JD, Moliterno DJ, Willard JE, Hillis LD, Lange RA. Influence of labetalol on cocaine-induced coronary vasoconstriction in humans. *Am J Med* 1993;94(6):608–10. [https://doi.org/10.1016/0002-9343\(93\)90212-8](https://doi.org/10.1016/0002-9343(93)90212-8).
 124. Callaway CW, Clark RF. Hyperthermia in psychostimulant overdose. *Ann Emerg Med* 1994;24(1):68–76. [https://doi.org/10.1016/s0196-0644\(94\)70165-2](https://doi.org/10.1016/s0196-0644(94)70165-2).
 125. Castro AL, Tarelho S, Almeida D, Sousa L, Franco JM, Teixeira HM. MDMA intoxication in a potential organ donor with cardiac arrest. *J Anal Toxicol* 2020;44(8):923–6. <https://doi.org/10.1093/jat/bkaa042>.
 126. Chen JP. Methamphetamine-associated acute myocardial infarction and cardiogenic shock with normal coronary arteries: refractory global coronary microvascular spasm. *J Invasive Cardiol* 2007;19(4):E89–92.

127. Grunau BE, Wiens MO, Brubacher JR. Dantrolene in the treatment of MDMA-related hyperpyrexia: a systematic review. *Cjem* 2010;12 (5):435–42. <https://doi.org/10.1017/s1481803500012598>.
128. Hollander JE, Carter WA, Hoffman RS. Use of phentolamine for cocaine-induced myocardial ischemia. *N Engl J Med* 1992;327 (5):361. <https://doi.org/10.1056/nejm199207303270517>.
129. Hysek C, Schmid Y, Rickli A, et al. Carvedilol inhibits the cardiostimulant and thermogenic effects of MDMA in humans. *Br J Pharmacol* 2012;166(8):2277–88. <https://doi.org/10.1111/j.1476-5381.2012.01936.x>.
130. Kiyatkin EA, Ren S. Clubbing with ecstasy. *Temperature (Austin)* 2014;1(3):160–1. <https://doi.org/10.4161/23328940.2014.980137>.
131. Lavonas EJ, Akpunonu PD, Arens AM, et al. 2023 American heart association focused update on the management of patients with cardiac arrest or life-threatening toxicity due to poisoning: an update to the american heart association guidelines for cardiopulmonary resuscitation and emergency cardiovascular care. *Circulation* 2023;148(16):e149–84. <https://doi.org/10.1161/CIR.0000000000001161>.
132. Liechti ME, Kunz I, Kupferschmidt H. Acute medical problems due to Ecstasy use. Case-series of emergency department visits. *Swiss Med Wkly* 2005;135(43–44):652–7. <https://doi.org/10.4414/smw.2005.11231>.
133. Lucyk SN. Acute cardiovascular toxicity of cocaine. *Canadian J Cardiol* 2022;38(9):1384–94. <https://doi.org/10.1016/j.cjca.2022.05.003>.
134. Matsumoto RR, Seminerio MJ, Turner RC, et al. Methamphetamine-induced toxicity: an updated review on issues related to hyperthermia. *Pharmacol Ther* 2014;144(1):28–40. <https://doi.org/10.1016/j.pharmthera.2014.05.001>.
135. Mende L, Böhm R, Regenthal R, Klein N, Grond S, Radke J. Cardiac arrest caused by an ecstasy intoxication. *Anesthesiol Intensivmed Notfallmed Schmerzther* 2005;40(12):762–5. <https://doi.org/10.1055/s-2005-870500> [In ger].
136. Mueller PD, Korey WS. Death by “ecstasy”: the serotonin syndrome?. *Ann Emerg Med* 1998;32(3 Pt 1):377–80. [https://doi.org/10.1016/s0196-0644\(98\)70018-6](https://doi.org/10.1016/s0196-0644(98)70018-6).
137. Yap CYL, Taylor DM, Knott JC, et al. Intravenous midazolam-droperidol combination, droperidol or olanzapine monotherapy for methamphetamine-related acute agitation: subgroup analysis of a randomized controlled trial. *Addiction* 2017;112(7):1262–9. <https://doi.org/10.1111/add.13780>.
138. Burns MJ, Linden CH, Graudins A, Brown RM, Fletcher KE. A comparison of physostigmine and benzodiazepines for the treatment of anticholinergic poisoning. *Annals of Emergency Medicine* 2000;35(4):374–81. [https://doi.org/10.1016/S0196-0644\(00\)70057-6](https://doi.org/10.1016/S0196-0644(00)70057-6).
139. Dawson AH, Buckley NA. Pharmacological management of anticholinergic delirium - theory, evidence and practice. *Br J Clin Pharmacol* 2016;81(3):516–24. <https://doi.org/10.1111/bcp.12839>.
140. Green R, Sitar DS, Tenenbein M. Effect of anticholinergic drugs on the efficacy of activated charcoal. *J Toxicol Clin Toxicol* 2004;42 (3):267–72. <https://doi.org/10.1081/clt-120037426>.
141. Jamshidi N, Dawson A. The hot patient: acute drug-induced hyperthermia. *Aust Prescr* 2019;42(1):24–8. <https://doi.org/10.18773/austprescr.2019.006>.
142. Rasimas JJ, Sachdeva KK, Donovan JW. Revival of an antidote: bedside experience with physostigmine. *Toxicology Communications* 2018;2(1):85–101. <https://doi.org/10.1080/24734306.2018.1535538>.
143. Haller CA, Benowitz NL. Adverse cardiovascular and central nervous system events associated with dietary supplements containing ephedra alkaloids. *New England Journal of Medicine* 2000;343(25):1833–8. <https://doi.org/10.1056/NEJM200012213432502>.
144. Pentel P. Toxicity of over-the-counter stimulants. *JAMA* 1984;252 (14):1898–903. <https://doi.org/10.1001/jama.1984.03350140044023>.
145. Rhidian R. Running a risk? Sport supplement toxicity with ephedrine in an amateur marathon runner, with subsequent rhabdomyolysis. *BMJ Case Rep* 2011;2011. <https://doi.org/10.1136/bcr.11.2011.5093>.
146. Chyka PA, Erdman AR, Christianson G, et al. Salicylate poisoning: An evidence-based consensus guideline for out-of-hospital management. *Clin Toxicol* 2007;45(2):95–131. <https://doi.org/10.1080/15563650600907140>.
147. Juurlink DN, Gosselin S, Kielstein JT, et al. Extracorporeal treatment for salicylate poisoning: systematic review and recommendations from the EXTRIP workgroup. *Annals of Emergency Medicine* 2015;66(2):165–81. <https://doi.org/10.1016/j.annemergmed.2015.03.031>.
148. Grundlingh J, Dargan PI, El-Zanfaly M, Wood DM. 2,4-dinitrophenol (DNP): a weight loss agent with significant acute toxicity and risk of death. *J Med Toxicol* 2011;7(3):205–12. <https://doi.org/10.1007/s13181-011-0162-6>.
149. Kopec KT, Kim T, Mowry J, Aks S, Kao L. Role of dantrolene in dinitrophenol (DNP) overdose: a continuing question?. *Am J Emerg Med* 2019;37(6):1216.e1–2. <https://doi.org/10.1016/j.ajem.2019.03.035>.
150. Pasquier M, Cools E, Zafren K, Carron PN, Frochoux V, Rousson V. Vital signs in accidental hypothermia. *High Alt Med Biol* 2021;22 (2):142–7. <https://doi.org/10.1089/ham.2020.0179>.
151. Frei C, Darocha T, Debaty G, et al. Clinical characteristics and outcomes of witnessed hypothermic cardiac arrest: a systematic review on rescue collapse. *Resuscitation* 2019;137:41–8. <https://doi.org/10.1016/j.resuscitation.2019.02.001>.
152. Podsiadlo P, Darocha T, Svendsen OS, et al. Outcomes of patients suffering unwitnessed hypothermic cardiac arrest rewarmed with extracorporeal life support: a systematic review. *Artif Organs* 2021;45(3):222–9. <https://doi.org/10.1111/aor.13818>.
153. Mydske S, Thomassen O. Is prehospital use of active external warming dangerous for patients with accidental hypothermia: a systematic review. *Scand J Trauma Resusc Emerg Med* 2020;28 (1):77. <https://doi.org/10.1186/s13049-020-00773-2>.
154. Fujimoto Y, Matsuyama T, Morita S, et al. Indoor versus outdoor occurrence in mortality of accidental hypothermia in japan: the j-point registry. *Ther Hypothermia Temp Manag* 2020;10(3):159–64. <https://doi.org/10.1089/ther.2019.0017>.
155. Mroczek T, Gladki M, Skalski J. Successful resuscitation from accidental hypothermia of 11.8 degrees C: where is the lower bound for human beings?. *Eur J Cardiothorac Surg* 2020;58 (5):1091–2. <https://doi.org/10.1093/ejcts/ezaa159>.
156. Musi ME, Sheets A, Zafren K, et al. Clinical staging of accidental hypothermia: the revised swiss system: recommendation of the international commission for mountain emergency medicine (ICAR MedCom). *Resuscitation* 2021;162:182–7. <https://doi.org/10.1016/j.resuscitation.2021.02.038>.
157. Paal P, Pasquier M, Darocha T, et al. Accidental Hypothermia: 2021 Update. *Int J Environ Res Public Health* 2022;19(1). <https://doi.org/10.3390/ijerph19010501>.
158. Pasquier M, Hugli O, Paal P, et al. Hypothermia outcome prediction after extracorporeal life support for hypothermic cardiac arrest patients: the HOPE score. *Resuscitation* 2018;126:58–64. <https://doi.org/10.1016/j.resuscitation.2018.02.026>.
159. Pasquier M, Paal P, Kosinski S, Brown D, Podsiadlo P, Darocha T. Esophageal temperature measurement. *N Engl J Med* 2020;383 (16):e93. <https://doi.org/10.1056/NEJMvcm1900481>.
160. Pasquier M, Rousson V, Darocha T, et al. Hypothermia outcome prediction after extracorporeal life support for hypothermic cardiac arrest patients: an external validation of the HOPE score. *Resuscitation* 2019;139:321–8. <https://doi.org/10.1016/j.resuscitation.2019.03.017>.
161. Podsiadlo P, Brozek T, Balik M, et al. Predictors of cardiac arrest in severe accidental hypothermia. *Am J Emerg Med* 2024;78:145–50. <https://doi.org/10.1016/j.ajem.2024.01.031>.

162. Woodhall B, Sealy WC, Hall KD, Floyd WL. Craniotomy under conditions of quinidine-protected cardioplegia and profound hypothermia. *Ann Surg* 1960;152(1):37–44. <https://doi.org/10.1097/0000658-196007000-00006>.
163. Pasquier M, Zurrón N, Weith B, et al. Deep accidental hypothermia with core temperature below 24 degrees c presenting with vital signs. *High Alt Med Biol* 2014;15(1):58–63. <https://doi.org/10.1089/ham.2013.1085>.
164. Soar J, Bottiger BW, Carli P, et al. European Resuscitation Council Guidelines 2025: Adult Advanced Life Support. 2025;215 (Suppl 1):110769.
165. Pasquier M ea. Estimation of the survival probability using HOPE. (<https://www.hypothermiascore.org>).
166. Gordon L, Paal P. Normothermic and hypothermic cardiac arrest-Beware of Jekyll and Hyde. *Resuscitation* 2018;129:e10–1. <https://doi.org/10.1016/j.resuscitation.2018.04.020>.
167. Wood SC. Interactions between hypoxia and hypothermia. *Annu Rev Physiol* 1991;53:71–85. <https://doi.org/10.1146/annurev.ph.53.030191.000443>.
168. Mair P, Gasteiger L, Mair B, Stroehle M, Walpoth B. Successful defibrillation of four hypothermic patients with witnessed cardiac arrest. *High Alt Med Biol* 2019;20(1):71–7. <https://doi.org/10.1089/ham.2018.0084>.
169. Stoner J, Martin G, O'Mara K, Ehlers J, Tomlanovich M. Amiodarone and bretylium in the treatment of hypothermic ventricular fibrillation in a canine model. *Acad Emerg Med* 2003;10(3):187–91. <https://doi.org/10.1111/j.1553-2712.2003.tb01988.x>.
170. Krismer AC, Lindner KH, Kornberger R, et al. Cardiopulmonary resuscitation during severe hypothermia in pigs: does epinephrine or vasopressin increase coronary perfusion pressure?. *Anesth Analg* 2000;90(1):69–73. <https://doi.org/10.1097/00000539-200001000-00017>.
171. Kornberger E, Lindner KH, Mayr VD, et al. Effects of epinephrine in a pig model of hypothermic cardiac arrest and closed-chest cardiopulmonary resuscitation combined with active rewarming. *Resuscitation* 2001;50(3):301–8. [https://doi.org/10.1016/s0300-9572\(01\)00353-7](https://doi.org/10.1016/s0300-9572(01)00353-7).
172. Cools E, Meyer M, Courvoisier D, Walpoth B. Successful pre-rewarming resuscitation after cardiac arrest in severe hypothermia: a retrospective cohort study from the international hypothermia registry. *Int J Environ Res Public Health* 2022;19(7). <https://doi.org/10.3390/ijerph19074059>.
173. Gordon L, Paal P, Ellertson JA, Brugger H, Peek GJ, Zafren K. Delayed and intermittent CPR for severe accidental hypothermia. *Resuscitation* 2015;90:46–9. <https://doi.org/10.1016/j.resuscitation.2015.02.017>.
174. Takauji S, Hayakawa M, Yamada D, et al. Outcome of extracorporeal membrane oxygenation use in severe accidental hypothermia with cardiac arrest and circulatory instability: a multicentre, prospective, observational study in Japan (ICE-CRASH study). *Resuscitation* 2023;182:109663. <https://doi.org/10.1016/j.resuscitation.2022.12.001>.
175. Ruttman E, Weissenbacher A, Ulmer H, et al. Prolonged extracorporeal membrane oxygenation-assisted support provides improved survival in hypothermic patients with cardiocirculatory arrest. *J Thorac Cardiovasc Surg* 2007;134(3):594–600. <https://doi.org/10.1016/j.jtcvs.2007.03.049>.
176. Mendrala K, Darocha T, Pluta M, et al. Outcomes of extracorporeal life support in hypothermic cardiac arrest: revisiting ELSO guidelines. *Resuscitation* 2024;205:110424. <https://doi.org/10.1016/j.resuscitation.2024.110424>.
177. Gruber E, Beikircher W, Pizzini R, et al. Non-extracorporeal rewarming at a rate of 6.8 degrees C per hour in a deeply hypothermic arrested patient. *Resuscitation* 2014;85(8):e119–20. <https://doi.org/10.1016/j.resuscitation.2014.05.011>.
178. Kuhnke M, Albrecht R, Scheffold JC, Paal P. Successful resuscitation from prolonged hypothermic cardiac arrest without extracorporeal life support: a case report. *J Med Case Rep* 2019;13(1):354. <https://doi.org/10.1186/s13256-019-2282-6>.
179. Okada Y, Matsuyama T, Morita S, et al. The development and validation of a “5A” severity scale for predicting in-hospital mortality after accidental hypothermia from J-point registry data. *J Intensive Care* 2019;7:27. <https://doi.org/10.1186/s40560-019-0384-2>.
180. Okada Y, Matsuyama T, Hayashida K, Takauji S, Kanda J, Yokobori S. External validation of 5A score model for predicting in-hospital mortality among the accidental hypothermia patients: JAAM-hypothermia study 2018-2019 secondary analysis. *J Intensive Care* 2022;10(1):24. <https://doi.org/10.1186/s40560-022-00616-5>.
181. Swol J, Darocha T, Paal P, et al. Extracorporeal life support in accidental hypothermia with cardiac arrest-a narrative review. *ASAIO J* 2022;68(2):153–62. <https://doi.org/10.1097/MAT.0000000000001518>.
182. Rauch S, Brugger H, Falk M, et al. Avalanche survival rates in Switzerland, 1981-2020. *JAMA Netw Open* 2024;7(9):e2435253. <https://doi.org/10.1001/jamanetworkopen.2024.35253>.
183. Boue Y, Payen JF, Brun J, et al. Survival after avalanche-induced cardiac arrest. *Resuscitation* 2014;85(9):1192–6. <https://doi.org/10.1016/j.resuscitation.2014.06.015>.
184. Moroder L, Mair B, Brugger H, Voelckel W, Mair P. Outcome of avalanche victims with out-of-hospital cardiac arrest. *Resuscitation* 2015;89:114–8. <https://doi.org/10.1016/j.resuscitation.2015.01.019>.
185. Metrailler-Mermoud J, Hugli O, Carron PN, et al. Avalanche victims in cardiac arrest are unlikely to survive despite adherence to medical guidelines. *Resuscitation* 2019;141:35–43. <https://doi.org/10.1016/j.resuscitation.2019.05.037>.
186. Procter E, Strapazzon G, Dal Cappello T, et al. Burial duration, depth and air pocket explain avalanche survival patterns in Austria and Switzerland. *Resuscitation* 2016;105:173–6. <https://doi.org/10.1016/j.resuscitation.2016.06.001>.
187. Rauch S, Koppenberg J, Josi D, et al. Avalanche survival depends on the time of day of the accident: a retrospective observational study. *Resuscitation* 2022;174:47–52. <https://doi.org/10.1016/j.resuscitation.2022.03.023>.
188. Pasquier M, Strapazzon G, Kottmann A, et al. On-site treatment of avalanche victims: scoping review and 2023 recommendations of the international commission for mountain emergency medicine (ICAR MedCom). *Resuscitation* 2023;184:109708. <https://doi.org/10.1016/j.resuscitation.2023.109708>.
189. Genswein M, Macias D, McIntosh S, Reiweger I, Hetland A, Paal P. AvaLife-A new multi-disciplinary approach supported by accident and field test data to optimize survival chances in rescue and first aid of avalanche patients. *Int J Environ Res Public Health* 2022;19(9). <https://doi.org/10.3390/ijerph19095257>.
190. Egger A, Niederer M, Tscherny K, et al. Influence of physical strain at high altitude on the quality of cardiopulmonary resuscitation. *Scand J Trauma Resusc Emerg Med* 2020;28(1):19. <https://doi.org/10.1186/s13049-020-0717-0>.
191. Putzer G, Braun P, Zimmermann A, et al. LUCAS compared to manual cardiopulmonary resuscitation is more effective during helicopter rescue-a prospective, randomized, cross-over manikin study. *Am J Emerg Med* 2013;31(2):384–9. <https://doi.org/10.1016/j.ajem.2012.07.018>.
192. Gassler H, Kurka L, Rauch S, Seewald S, Kulla M, Fischer M. Mechanical chest compression devices under special circumstances. *Resuscitation* 2022;179:183–8. <https://doi.org/10.1016/j.resuscitation.2022.06.014>.
193. Konstantinides SV, Meyer G, Becattini C, et al. 2019 ESC Guidelines for the diagnosis and management of acute pulmonary embolism developed in collaboration with the European Respiratory Society (ERS). *Eur Heart J* 2020;41(4):543–603. <https://doi.org/10.1093/eurheartj/ehz405>.
194. Javaudin F, Lascarrou JB, Le Bastard Q, et al. Thrombolysis during resuscitation for out-of-hospital cardiac arrest caused by pulmonary embolism increases 30-day survival: findings from the French

- national cardiac arrest registry. *Chest* 2019;156(6):1167–75. <https://doi.org/10.1016/j.chest.2019.07.015>.
195. Bottiger BW, Arntz HR, Chamberlain DA, et al. Thrombolysis during resuscitation for out-of-hospital cardiac arrest. *N Engl J Med* 2008;359(25):2651–62. <https://doi.org/10.1056/NEJMoa070570>.
 196. Bergum D, Nordseth T, Mjølstad OC, Skogvoll E, Haugen BO. Causes of in-hospital cardiac arrest - incidences and rate of recognition. *Resuscitation* 2015;87:63–8. <https://doi.org/10.1016/j.resuscitation.2014.11.007>.
 197. Kurciciyan I, Meron G, Sterz F, et al. Pulmonary embolism as a cause of cardiac arrest: presentation and outcome. *Arch Intern Med* 2000;160(10):1529–35. <https://doi.org/10.1001/archinte.160.10.1529>.
 198. Pudil J, Rob D, Smalcova J, et al. Pulmonary embolism-related refractory out-of-hospital cardiac arrest and extracorporeal cardiopulmonary resuscitation: prague OHCA study post hoc analysis. *Eur Heart J Acute Cardiovasc Care* 2023;12(8):507–12. <https://doi.org/10.1093/ehjacc/zuad052>.
 199. Soar J, Berg KM, Andersen LW, et al. Adult advanced life support: 2020 international consensus on cardiopulmonary resuscitation and emergency cardiovascular care science with treatment recommendations. *Resuscitation* 2020;156:A80–A119. <https://doi.org/10.1016/j.resuscitation.2020.09.012>.
 200. Soar J, Maconochie I, Wyckoff MH, et al. 2019 international consensus on cardiopulmonary resuscitation and emergency cardiovascular care science with treatment recommendations. *Resuscitation* 2019;145:95–150. <https://doi.org/10.1016/j.resuscitation.2019.10.016>.
 201. Wyckoff MH, Greif R, Morley PT, et al. 2022 international consensus on cardiopulmonary resuscitation and emergency cardiovascular care science with treatment recommendations: summary from the basic life support; advanced life support; pediatric life support; neonatal life support; education, implementation, and teams; and first aid task forces. *Resuscitation* 2022;181:208–88. <https://doi.org/10.1016/j.resuscitation.2022.10.005>.
 202. Heradstveit BE, Sunde K, Sunde GA, Wentzel-Larsen T, Heltne JK. Factors complicating interpretation of capnography during advanced life support in cardiac arrest—a clinical retrospective study in 575 patients. *Resuscitation* 2012;83(7):813–8. <https://doi.org/10.1016/j.resuscitation.2012.02.021>.
 203. Aagaard R, Lofgren B, Caap P, Mygind-Klausen T, Botker MT, Granfeldt A. A low end-tidal CO(2)/arterial CO(2) ratio during cardiopulmonary resuscitation suggests pulmonary embolism. *Resuscitation* 2018;133:137–40. <https://doi.org/10.1016/j.resuscitation.2018.10.008>.
 204. Bova C, Greco F, Misuraca G, et al. Diagnostic utility of echocardiography in patients with suspected pulmonary embolism. *Am J Emerg Med* 2003;21(3):180–3. [https://doi.org/10.1016/s0735-6757\(02\)42257-7](https://doi.org/10.1016/s0735-6757(02)42257-7).
 205. Geibel A, Zehender M, Kasper W, Olschewski M, Klima C, Konstantinides SV. Prognostic value of the ECG on admission in patients with acute major pulmonary embolism. *Eur Respir J* 2005;25(5):843–8. <https://doi.org/10.1183/09031936.05.00119704>.
 206. Wan S, Quinlan DJ, Agnelli G, Eikelboom JW. Thrombolysis compared with heparin for the initial treatment of pulmonary embolism: a meta-analysis of the randomized controlled trials. *Circulation* 2004;110(6):744–9. <https://doi.org/10.1161/01.CIR.0000137826.09715.9C>.
 207. Ruiz-Bailen M, Aguayo-de-Hoyos E, Serrano-Corcoles MC, et al. Thrombolysis with recombinant tissue plasminogen activator during cardiopulmonary resuscitation in fulminant pulmonary embolism. A case series. *Resuscitation* 2001;51(1):97–101. [https://doi.org/10.1016/s0300-9572\(01\)00384-7](https://doi.org/10.1016/s0300-9572(01)00384-7).
 208. Sharifi M, Berger J, Beeston P, et al. Pulseless electrical activity in pulmonary embolism treated with thrombolysis (from the “PEAPETT” study). *Am J Emerg Med* 2016;34(10):1963–7. <https://doi.org/10.1016/j.ajem.2016.06.094>.
 209. Wu JP, Gu DY, Wang S, Zhang ZJ, Zhou JC, Zhang RF. Good neurological recovery after rescue thrombolysis of presumed pulmonary embolism despite prior 100 minutes CPR. *J Thorac Dis* 2014;6(12):E289–93. <https://doi.org/10.3978/j.issn.2072-1439.2014.12.23>.
 210. Summers K, Schultheis J, Raiff D, Dahhan T. Evaluation of rescue thrombolysis in cardiac arrest secondary to suspected or confirmed pulmonary embolism. *Ann Pharmacother* 2019;53(7):711–5. <https://doi.org/10.1177/1060028019828423>.
 211. Fava M, Loyola S, Bertoni H, Dougnac A. Massive pulmonary embolism: percutaneous mechanical thrombectomy during cardiopulmonary resuscitation. *J Vasc Interv Radiol* 2005;16(1):119–23. <https://doi.org/10.1097/01.RVI.0000146173.85401.BA>.
 212. Pasirja C, Kronfli A, Rouse M, et al. Outcomes after surgical pulmonary embolectomy for acute submassive and massive pulmonary embolism: a single-center experience. *J Thorac Cardiovasc Surg* 2018;155(3):1095–1106 e2. <https://doi.org/10.1016/j.jtcvs.2017.10.139>.
 213. Konstantinov IE, Saxena P, Koniuszko MD, Alvarez J, Newman MA. Acute massive pulmonary embolism with cardiopulmonary resuscitation: management and results. *Tex Heart Inst J* 2007;34(1):41–5. discussion 45–6 Available from: <https://www.ncbi.nlm.nih.gov/pubmed/17420792>.
 214. Goldhaber SZ, Haire WD, Feldstein ML, et al. Alteplase versus heparin in acute pulmonary embolism: randomised trial assessing right-ventricular function and pulmonary perfusion. *Lancet* 1993;341(8844):507–11. [https://doi.org/10.1016/0140-6736\(93\)90274-k](https://doi.org/10.1016/0140-6736(93)90274-k).
 215. Maj G, Melisurgo G, De Bonis M, Pappalardo F. ECLS management in pulmonary embolism with cardiac arrest: which strategy is better?. *Resuscitation* 2014;85(10):e175–6. <https://doi.org/10.1016/j.resuscitation.2014.03.309>.
 216. Debaty G, Babaz V, Durand M, et al. Prognostic factors for extracorporeal cardiopulmonary resuscitation recipients following out-of-hospital refractory cardiac arrest: A systematic review and meta-analysis. *Resuscitation* 2017;112:1–10. <https://doi.org/10.1016/j.resuscitation.2016.12.011>.
 217. Richardson ASC, Tonna JE, Nanjaya V, et al. Extracorporeal cardiopulmonary resuscitation in adults. Interim guideline consensus statement from the extracorporeal life support organization. *ASAIO J* 2021;67(3):221–8. <https://doi.org/10.1097/MAT.0000000000001344>.
 218. Sakuraya M, Hifumi T, Inoue A, Sakamoto T, Kuroda Y Group S-JS. Neurological outcomes and reperfusion strategies in out-of-hospital cardiac arrest patients due to pulmonary embolism who underwent venoarterial extracorporeal membrane oxygenation: a post-hoc analysis of a multicenter retrospective cohort study. *Resuscitation* 2023;191:109926. <https://doi.org/10.1016/j.resuscitation.2023.109926>.
 219. Hayashi M, Shimizu W, Albert CM. The spectrum of epidemiology underlying sudden cardiac death. *Circ Res* 2015;116(12):1887–906. <https://doi.org/10.1161/CIRCRESAHA.116.304521>.
 220. Geri G, Passouant O, Dumas F, et al. Etiological diagnoses of out-of-hospital cardiac arrest survivors admitted to the intensive care unit: Insights from a French registry. *Resuscitation* 2017;117:66–72. <https://doi.org/10.1016/j.resuscitation.2017.06.006>.
 221. Vahatalo J, Holmstrom L, Pakanen L, et al. Coronary artery disease as the cause of sudden cardiac death among victims < 50 years of age. *Am J Cardiol* 2021;147:33–8. <https://doi.org/10.1016/j.amicard.2021.02.012>.
 222. Zachariassardottir S, Risgaard B, Agesen FN, et al. Sudden cardiac death and coronary disease in the young: a nationwide cohort study in Denmark. *Int J Cardiol* 2017;236:16–22. <https://doi.org/10.1016/j.ijcard.2017.01.118>.
 223. Scquizzato T, Sofia R, Gazzato A, et al. Coronary angiography findings in resuscitated and refractory out-of-hospital cardiac arrest: a systematic review and meta-analysis. *Resuscitation*

- 2023;189:109869. <https://doi.org/10.1016/j.resuscitation.2023.109869>.
224. Visseren FLJ, Mach F, Smulders YM, et al. 2021 ESC Guidelines on cardiovascular disease prevention in clinical practice. *Eur Heart J* 2021;42(34):3227–337. <https://doi.org/10.1093/eurheartj/ehab484>.
 225. Byrne RA, Rossello X, Coughlan JJ, et al. 2023 ESC Guidelines for the management of acute coronary syndromes. *Eur Heart J* 2023;44(38):3720–826. <https://doi.org/10.1093/eurheartj/ehad191>.
 226. Farquharson B, Abhyankar P, Smith K, et al. Reducing delay in patients with acute coronary syndrome and other time-critical conditions: a systematic review to identify the behaviour change techniques associated with effective interventions. *Open Heart* 2019;6(1):e000975. <https://doi.org/10.1136/openhrt-2018-000975>.
 227. Cartanya-Bonvehi J, Pericas-Vila A, Subirana I, Garcia-Garcia C, Tizon-Marcos H, Elosua R. Effectiveness of STEMI networks with out-of-hospital triage: a systematic review and meta-analysis. *Rev Esp Cardiol (Engl Ed)* 2024. <https://doi.org/10.1016/j.rec.2024.07.008>.
 228. Baldi E, Schnaubelt S, Caputo ML, et al. Association of timing of electrocardiogram acquisition after return of spontaneous circulation with coronary angiography findings in patients with out-of-hospital cardiac arrest. *JAMA Netw Open* 2021;4(1):e2032875. <https://doi.org/10.1001/jamanetworkopen.2020.32875>.
 229. McFadden P, Reynolds JC, Maddler RD, Brown M. Diagnostic test accuracy of the initial electrocardiogram after resuscitation from cardiac arrest to indicate invasive coronary angiographic findings and attempted revascularization: a systematic review and meta-analysis. *Resuscitation* 2021;160:20–36. <https://doi.org/10.1016/j.resuscitation.2020.11.039>.
 230. Kim YJ, Min SY, Lee DH, et al. The role of post-resuscitation electrocardiogram in patients with ST-segment changes in the immediate post-cardiac arrest period. *JACC Cardiovasc Interv* 2017;10(5):451–9. <https://doi.org/10.1016/j.jcin.2016.11.046>.
 231. Garcia-Tejada J, Jurado-Roman A, Rodriguez J, et al. Post-resuscitation electrocardiograms, acute coronary findings and in-hospital prognosis of survivors of out-of-hospital cardiac arrest. *Resuscitation* 2014;85(9):1245–50. <https://doi.org/10.1016/j.resuscitation.2014.06.001>.
 232. Drennan IR, Nikolaou N, Netherton S, et al. Early Coronary Angiography Post-ROSC: Consensus on Science with Treatment Recommendations. [Internet] Brussels, Belgium: International Liaison Committee on Resuscitation (ILCOR) Advanced Life Support Task Force. <http://ilcor.org>.
 233. Elfwen L, Lagedal R, James S, et al. Coronary angiography in out-of-hospital cardiac arrest without ST elevation on ECG-Short- and long-term survival. *Am Heart J* 2018;200:90–5. <https://doi.org/10.1016/j.ahj.2018.03.009>.
 234. Patel N, Patel NJ, Macon CJ, et al. Trends and outcomes of coronary angiography and percutaneous coronary intervention after out-of-hospital cardiac arrest associated with ventricular fibrillation or pulseless ventricular tachycardia. *JAMA Cardiol* 2016;1(8):890–9. <https://doi.org/10.1001/jamacardio.2016.2860>.
 235. Nikolaou NI, Netherton S, Welsford M, et al. A systematic review and meta-analysis of the effect of routine early angiography in patients with return of spontaneous circulation after out-of-hospital cardiac arrest. *Resuscitation* 2021;163:28–48. <https://doi.org/10.1016/j.resuscitation.2021.03.019>.
 236. Elfwen L, Lagedal R, Nordberg P, et al. Direct or subacute coronary angiography in out-of-hospital cardiac arrest (DISCO)-an initial pilot-study of a randomized clinical trial. *Resuscitation* 2019;139:253–61. <https://doi.org/10.1016/j.resuscitation.2019.04.027>.
 237. Lemkes JS, Janssens GN, van der Hoeven NW, et al. Coronary angiography after cardiac arrest without ST-segment elevation. *N Engl J Med* 2019;380(15):1397–407. <https://doi.org/10.1056/NEJMoa1816897>.
 238. Kern KB, Radsel P, Jentzer JC, et al. Randomized pilot clinical trial of early coronary angiography versus no early coronary angiography after cardiac arrest without ST-segment elevation: the PEARL study. *Circulation* 2020;142(21):2002–12. <https://doi.org/10.1161/CIRCULATIONAHA.120.049569>.
 239. Desch S, Freund A, Akin I, et al. Angiography after out-of-hospital cardiac arrest without ST-segment elevation. *N Engl J Med* 2021;385(27):2544–53. <https://doi.org/10.1056/NEJMoa2101909>.
 240. Hauw-Berlemont C, Lamhaut L, Diehl JL, et al. Emergency vs delayed coronary angiogram in survivors of out-of-hospital cardiac arrest: results of the randomized, multicentric EMERGE trial. *JAMA Cardiol* 2022;7(7):700–7. <https://doi.org/10.1001/jamacardio.2022.1416>.
 241. Viana-Tejedor A, Andrea-Riba R, Scardino C, et al. Coronary angiography in patients without ST-segment elevation following out-of-hospital cardiac arrest. COUPE clinical trial. *Rev Esp Cardiol (Engl Ed)* 2023;76(2):94–102. <https://doi.org/10.1016/j.rec.2022.05.013>.
 242. Pocock H, Nicholson T, Szarpak L, Soar J, Berg KM, Force. obotILCoRALST. Mechanical CPR Devices: Consensus on Science with Treatment Recommendations. [Internet] Brussels, Belgium: International Liaison Committee on Resuscitation (ILCOR) Advanced Life Support Task Force. <http://ilcor.org>.
 243. Holmberg MJ, Granfeldt A, Guerguerian AM, et al. Extracorporeal cardiopulmonary resuscitation for cardiac arrest: an updated systematic review. *Resuscitation* 2023;182:109665. <https://doi.org/10.1016/j.resuscitation.2022.12.003>.
 244. Yannopoulos D, Bartos J, Raveendran G, et al. Advanced reperfusion strategies for patients with out-of-hospital cardiac arrest and refractory ventricular fibrillation (ARREST): a phase 2, single centre, open-label, randomised controlled trial. *Lancet* 2020;396(10265):1807–16. [https://doi.org/10.1016/S0140-6736\(20\)32338-2](https://doi.org/10.1016/S0140-6736(20)32338-2).
 245. Belohlavek J, Smalcova J, Rob D, et al. Effect of intra-arrest transport, extracorporeal cardiopulmonary resuscitation, and immediate invasive assessment and treatment on functional neurologic outcome in refractory out-of-hospital cardiac arrest: a randomized clinical trial. *JAMA* 2022;327(8):737–47. <https://doi.org/10.1001/jama.2022.1025>.
 246. Suverein MM, Delnoij TSR, Lorusso R, et al. Early extracorporeal CPR for refractory out-of-hospital cardiac arrest. *N Engl J Med* 2023;388(4):299–309. <https://doi.org/10.1056/NEJMoa2204511>.
 247. Grunau B, O'Neil BJ, Giustini D, Drennan IR, Lavonas EJ. Opioid-associated cardiac arrest: a systematic review of intra-arrest naloxone and other opioid-specific advanced life-support therapies. *Resusc Plus* 2025;100906.
 248. Beaulieu J, St-Onge M. Antidote use for cardiac arrest or hemodynamic instability due to cardiac glycoside poisoning: a narrative review. *Resusc Plus* 2024;19:100690.
 249. Maier S, Rösner L, Saemann L. Extracorporeal membrane oxygenation in intoxication and overdoses: a systematic review. *Thorac Cardiovasc Surg* 2024;72(04):288–95.
 250. Lalande E, Burwash-Brennan T, Burns K, et al. Is point-of-care ultrasound a reliable predictor of outcome during traumatic cardiac arrest? A systematic review and meta-analysis from the SHoC investigators. *Resuscitation* 2021;167:128–36. <https://doi.org/10.1016/j.resuscitation.2021.08.027>.
 251. Liu A, Nguyen J, Ehrlich H, et al. Emergency resuscitative thoracotomy for civilian thoracic trauma in the field and emergency department settings: a systematic review and meta-analysis. *J Surg Res* 2022;273:44–55. <https://doi.org/10.1016/j.jss.2021.11.012>.
 252. Narvestad JK, Meskinfarnard M, Søreide K. Emergency resuscitative thoracotomy performed in European civilian trauma patients with blunt or penetrating injuries: a systematic review. *Eur J Trauma Emerg Surg* 2016;42(6):677–85. <https://doi.org/10.1007/s00068-015-0559-z>.
 253. Shi D, McLaren C, Evans C. Neurological outcomes after traumatic cardiopulmonary arrest: a systematic review. *Trauma Surg Acute*

- Care Open 2021;6(1)e000817. <https://doi.org/10.1136/tsaco-2021-000817>.
254. Tran A, Fernando SM, Rochweg B, et al. Pre-arrest and intra-arrest prognostic factors associated with survival following traumatic out-of-hospital cardiac arrest – A systematic review and meta-analysis. *Resuscitation* 2020;153:119–35. <https://doi.org/10.1016/j.resuscitation.2020.05.052>.
 255. Vianen NJ, Van Lieshout EMM, Maissan IM, et al. Prehospital traumatic cardiac arrest: a systematic review and meta-analysis. *Eur J Trauma Emerg Surg* 2022;48(4):3357–72. <https://doi.org/10.1007/s00068-022-01941-y>.
 256. Wongtanasarasin W, Thepchinda T, Kasirawat C, Saetiao S, Leungvorawat J, Kittivorakanchai N. Treatment outcomes of epinephrine for traumatic out-of-hospital cardiac arrest: a systematic review and meta-analysis. *J Emerg Trauma Shock* 2021;14(4):195–200. https://doi.org/10.4103/jets.Jets_35_21.
 257. Yang Z, Song Z, Hou M. Target temperature management versus normal temperature management for cardiac arrest after traumatic brain injury patient: a meta-analysis and systemic review. *Ther Hypothermia Temp Manag* 2022;12(3):139–45. <https://doi.org/10.1089/ther.2022.0007>.
 258. Morgan KM, Abou Khalil E, Feeney EV, et al. The efficacy of low-titer group O whole blood compared with component therapy in civilian trauma patients: a meta-analysis. *Crit Care Med* 2024;52(7):e390–404. <https://doi.org/10.1097/CCM.0000000000006244>.
 259. Grasner JT, Wnent J, Herlitz J, et al. Survival after out-of-hospital cardiac arrest in Europe - Results of the EuReCa TWO study. *Resuscitation* 2020;148:218–26. <https://doi.org/10.1016/j.resuscitation.2019.12.042>.
 260. Talmy T, Greenstein I, Gendler S, et al. Survival following prehospital traumatic cardiac arrest resuscitation in the israel defense forces: a retrospective study. *Prehosp Emerg Care* 2024;28(3):438–47. <https://doi.org/10.1080/10903127.2023.2241542>.
 261. Zwingmann J, Lefering R, Feucht M, Sudkamp NP, Stroh PC, Hammer T. Outcome and predictors for successful resuscitation in the emergency room of adult patients in traumatic cardiorespiratory arrest. *Crit Care* 2016;20(1):282. <https://doi.org/10.1186/s13054-016-1463-6>.
 262. Alqudah Z, Nehme Z, Alrawashdeh A, Williams B, Oteir A, Smith K. Paediatric traumatic out-of-hospital cardiac arrest: a systematic review and meta-analysis. *Resuscitation* 2020;149:65–73. <https://doi.org/10.1016/j.resuscitation.2020.01.037>.
 263. Cera SM, Mostafa G, Sing RF, Sarafin JL, Matthews BD, Heniford BT. Physiologic predictors of survival in post-traumatic arrest. *Am Surg* 2003;69(2):140–4. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/12641355>.
 264. Chen YC, Wu KH, Hsiao KY, et al. Factors associated with outcomes in traumatic cardiac arrest patients without prehospital return of spontaneous circulation. *Injury* 2019;50(1):4–9. <https://doi.org/10.1016/j.injury.2018.07.010>.
 265. Djarv T, Axelsson C, Herlitz J, Stromsoe A, Israelsson J, Claesson A. Traumatic cardiac arrest in Sweden 1990–2016 – A population-based national cohort study. *Scand J Trauma Resusc Emerg Med* 2018;26(1):30. <https://doi.org/10.1186/s13049-018-0500-7>.
 266. Israr S, Cook AD, Chapple KM, et al. Pulseless electrical activity following traumatic cardiac arrest: sign of life or death?. *Injury* 2019;50(9):1507–10. <https://doi.org/10.1016/j.injury.2019.05.025>.
 267. Obara T, Yumoto T, Bunya N, et al. Association between signs of life and survival in traumatic cardiac arrest patients: a nationwide, retrospective cohort study. *Resusc Plus* 2024;19:100701. <https://doi.org/10.1016/j.resplu.2024.100701>.
 268. Seamon MJ, Haut ER, Van Arendonk K, et al. An evidence-based approach to patient selection for emergency department thoracotomy: a practice management guideline from the Eastern Association for the Surgery of Trauma. *J Trauma Acute Care Surg* 2015;79(1):159–73. <https://doi.org/10.1097/TA.0000000000000648>.
 269. Seewald S, Wnent J, Grasner JT, et al. Survival after traumatic cardiac arrest is possible—a comparison of German patient-registries. *BMC Emerg Med* 2022;22(1):158. <https://doi.org/10.1186/s12873-022-00714-5>.
 270. Stratton SJ, Brickett K, Crammer T. Prehospital pulseless, unconscious penetrating trauma victims: field assessments associated with survival. *J Trauma* 1998;45(1):96–100. <https://doi.org/10.1097/00005373-199807000-00021>.
 271. Millin MG, Galvagno SM, Khandker SR, et al. Withholding and termination of resuscitation of adult cardiopulmonary arrest secondary to trauma: resource document to the joint NAEMSP-ACSCOT position statements. *J Trauma Acute Care Surg* 2013;75(3):459–67. <https://doi.org/10.1097/TA.0b013e31829cfaea>.
 272. Leis CC, Hernandez CC, Blanco MJ, Paterna PC, Hernandez Rde E, Torres EC. Traumatic cardiac arrest: should advanced life support be initiated?. *J Trauma Acute Care Surg* 2013;74(2):634–8. <https://doi.org/10.1097/TA.0b013e31827d5d3c>.
 273. Rossaint R, Afshari A, Bouillon B, et al. The European guideline on management of major bleeding and coagulopathy following trauma: sixth edition. *Crit Care* 2023;27(1):80. <https://doi.org/10.1186/s13054-023-04327-7>.
 274. Kleber C, Giesecke MT, Tsokos M, Haas NP, Buschmann CT. Trauma-related preventable deaths in Berlin 2010: need to change prehospital management strategies and trauma management education. *World J Surg* 2013;37(5):1154–61. <https://doi.org/10.1007/s00268-013-1964-2>.
 275. Davis JS, Satahoo SS, Butler FK, et al. An analysis of prehospital deaths: who can we save?. *J Trauma Acute Care Surg* 2014;77(2):213–8. <https://doi.org/10.1097/TA.0000000000000292>.
 276. Beck B, Smith K, Mercier E, et al. Potentially preventable trauma deaths: a retrospective review. *Injury* 2019;50(5):1009–16. <https://doi.org/10.1016/j.injury.2019.03.003>.
 277. Lott C, Araujo R, Cassar MR, et al. The European Trauma Course (ETC) and the team approach: past, present and future. *Resuscitation* 2009;80(10):1192–6. <https://doi.org/10.1016/j.resuscitation.2009.06.023>.
 278. Kleber C, Giesecke MT, Lindner T, Haas NP, Buschmann CT. Requirement for a structured algorithm in cardiac arrest following major trauma: epidemiology, management errors, and preventability of traumatic deaths in Berlin. *Resuscitation* 2014;85(3):405–10. <https://doi.org/10.1016/j.resuscitation.2013.11.009>.
 279. Luna GK, Pavlin EG, Kirkman T, Copass MK, Rice CL. Hemodynamic effects of external cardiac massage in trauma shock. *J Trauma* 1989;29(10):1430–3. <https://doi.org/10.1097/00005373-198910000-00022>.
 280. Jeffcoach DR, Gallegos JJ, Jesty SA, et al. Use of CPR in hemorrhagic shock, a dog model. *J Trauma Acute Care Surg* 2016;81(1):27–33. <https://doi.org/10.1097/TA.0000000000001001>.
 281. Watts S, Smith JE, Gwyther R, Kirkman E. Closed chest compressions reduce survival in an animal model of haemorrhage-induced traumatic cardiac arrest. *Resuscitation* 2019;140:37–42. <https://doi.org/10.1016/j.resuscitation.2019.04.048>.
 282. Endo A, Kojima M, Hong ZJ, Otomo Y, Coimbra R. Open-chest versus closed-chest cardiopulmonary resuscitation in trauma patients with signs of life upon hospital arrival: a retrospective multicenter study. *Crit Care* 2020;24(1):541. <https://doi.org/10.1186/s13054-020-03259-w>.
 283. Ji C, Pocock H, Deakin CD, et al. Adrenaline for traumatic cardiac arrest: a post hoc analysis of the PARAMEDIC2 trial. *Resuscitation Plus* 2025;22:100890. <https://doi.org/10.1016/j.resplu.2025.100890>.
 284. Witt CE, Shatz DV, Robinson BRH, et al. Epinephrine in prehospital traumatic cardiac arrest—life saving or false hope?. *Prehosp Emerg Care* 2025;1–9. <https://doi.org/10.1080/10903127.2025.2461283>.

285. Weegeenaar C, Perkins Z, Lockey D. Pre-hospital management of traumatic cardiac arrest 2024 position statement: faculty of prehospital care, royal college of surgeons of Edinburgh. *Scand J Trauma Resusc Emerg Med* 2024;32(1):139. <https://doi.org/10.1186/s13049-024-01304-z>.
286. Ferrada P, Garcia A, Duchesne J, et al. Comparing outcomes in patients with exsanguinating injuries: an Eastern Association for the Surgery of Trauma (EAST), multicenter, international trial evaluating prioritization of circulation over intubation (CAB over ABC). *World J Emerg Surg* 2024;19(1):15. <https://doi.org/10.1186/s13017-024-00545-8>.
287. Jamali B, Nouri S, Amidi S. Local and systemic hemostatic agents: a comprehensive review. *Cureus* 2024;16(10):e72312. <https://doi.org/10.7759/cureus.72312>.
288. Collaborators. C-, Roberts I, Shakur H, et al. The importance of early treatment with tranexamic acid in bleeding trauma patients: an exploratory analysis of the CRASH-2 randomised controlled trial. *Lancet* 2011;377(9771):1096–101, 1101 e1–2. [https://doi.org/10.1016/S0140-6736\(11\)60278-X](https://doi.org/10.1016/S0140-6736(11)60278-X).
289. Bulger EM, Perina DG, Qasim Z, et al. Clinical use of resuscitative endovascular balloon occlusion of the aorta (REBOA) in civilian trauma systems in the USA, 2019: a joint statement from the American College of Surgeons Committee on Trauma, the American College of Emergency Physicians, the National Association of Emergency Medical Services Physicians and the National Association of Emergency Medical Technicians. *Trauma Surg Acute Care Open* 2019;4(1):e000376. <https://doi.org/10.1136/tsaco-2019-000376>.
290. Jansen JO, Hudson J, Cochran C, et al. Emergency department resuscitative endovascular balloon occlusion of the aorta in trauma patients with exsanguinating hemorrhage: the UK-REBOA randomized clinical trial. *JAMA* 2023;330(19):1862–71. <https://doi.org/10.1001/jama.2023.20850>.
291. Sperry JL, Guyette FX, Brown JB, et al. Prehospital plasma during air medical transport in trauma patients at risk for hemorrhagic shock. *N Engl J Med* 2018;379(4):315–26. <https://doi.org/10.1056/NEJMoa1802345>.
292. Guyette FX, Sperry JL, Peitzman AB, et al. Prehospital blood product and crystalloid resuscitation in the severely injured patient: a secondary analysis of the prehospital air medical plasma trial. *Ann Surg* 2021;273(2):358–64. <https://doi.org/10.1097/SLA.0000000000003324>.
293. Braverman MA, Schauer SG, Ciaraglia A, et al. The impact of prehospital whole blood on hemorrhaging trauma patients: a multi-center retrospective study. *J Trauma Acute Care Surg* 2023;95(2):191–6. <https://doi.org/10.1097/TA.0000000000003908>.
294. Atkinson JL. The neglected prehospital phase of head injury: apnea and catecholamine surge. *Mayo Clin Proc* 2000;75(1):37–47. <https://doi.org/10.4065/75.1.37>.
295. Wilson MH, Hinds J, Grier G, Burns B, Carley S, Davies G. Impact brain apnoea – A forgotten cause of cardiovascular collapse in trauma. *Resuscitation* 2016;105:52–8. <https://doi.org/10.1016/j.resuscitation.2016.05.007>.
296. Barnard E, Yates D, Edwards A, Fragoso-Iniguez M, Jenks T, Smith JE. Epidemiology and aetiology of traumatic cardiac arrestland and Wales – A retrospective database analysis. *Resuscitation* 2017;110:90–4. <https://doi.org/10.1016/j.resuscitation.2016.11.001>.
297. Houwen T, Popal Z, de Bruijn MAN, et al. Outcomes after prehospital traumatic cardiac arrest in the Netherlands: a retrospective cohort study. *Injury* 2021;52(5):1117–22. <https://doi.org/10.1016/j.injury.2021.02.088>.
298. Hudson AJ, Strandenes G, Bjerkvig CK, Svanevik M, Glassberg E. Airway and ventilation management strategies for hemorrhagic shock. To tube, or not to tube, that is the question!. *J Trauma Acute Care Surg* 2018;84(6S Suppl 1):S77–82. <https://doi.org/10.1097/TA.0000000000001822>.
299. El-Sayed AA, Arafa SK, El-Demerdash AM. Pressure-controlled ventilation could decrease intraoperative blood loss and improve airway pressure measures during lumbar discectomy in the prone position: a comparison with volume-controlled ventilation mode. *J Anaesthesiol Clin Pharmacol* 2019;35(4):468–74. <https://doi.org/10.4103/joacp.JOACP.288.18>.
300. Pepe PE, Roppolo LP, Fowler RL. The detrimental effects of ventilation during low-blood-flow states. *Curr Opin Crit Care* 2005;11(3):212–8. <https://doi.org/10.1097/01.ccx.0000163650.80601.24>.
301. Davis JA, Manoach S, Heerdt P, Berlin DA. Management of respiratory failure in hemorrhagic shock. *Ann Am Thorac Soc* 2024;21(7):993–7. <https://doi.org/10.1513/AnnalsATS.202310-905CME>.
302. Roberts DJ, Leigh-Smith S, Faris PD, et al. Clinical presentation of patients with tension pneumothorax: a systematic review. *Ann Surg* 2015;261(6):1068–78. <https://doi.org/10.1097/SLA.0000000000001073>.
303. Deakin CD, Davies G, Wilson A. Simple thoracostomy avoids chest drain insertion in prehospital trauma. *J Trauma* 1995;39(2):373–4. <https://doi.org/10.1097/00005373-199508000-00031>.
304. Escott ME, Gleisberg GR, Kimmel K, Karrer A, Cosper J, Monroe BJ. Simple thoracostomy. Moving beyond needle decompression in traumatic cardiac arrest. *JEMS* 2014;39(4):26–32. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/24779096>.
305. Clemency BM, Tanski CT, Rosenberg M, May PR, Consiglio JD, Lindstrom HA. Sufficient catheter length for pneumothorax needle decompression: a meta-analysis. *Prehosp Disaster Med* 2015;30(3):249–53. <https://doi.org/10.1017/S1049023X15004653>.
306. Hecker M, Hegenscheid K, Volzke H, et al. Needle decompression of tension pneumothorax: population-based epidemiologic approach to adequate needle length in healthy volunteers in Northeast Germany. *J Trauma Acute Care Surg* 2016;80(1):119–24. <https://doi.org/10.1097/TA.0000000000000878>.
307. Wise D, Davies G, Coats T, Lockey D, Hyde J, Good A. Emergency thoracotomy: “how to do it”. *Emerg Med J* 2005;22(1):22–4. <https://doi.org/10.1136/emj.2003.012963>.
308. Flaris AN, Simms ER, Prat N, Reynard F, Caillot JL, Voiglio EJ. Clamshell incision versus left anterolateral thoracotomy. Which one is faster when performing a resuscitative thoracotomy? The tortoise and the hare revisited. *World J Surg* 2015;39(5):1306–11. <https://doi.org/10.1007/s00268-014-2924-1>.
309. Lee TH, Ouellet JF, Cook M, Schreiber MA, Kortbeek JB. Pericardiocentesis in trauma: a systematic review. *J Trauma Acute Care Surg* 2013;75(4):543–9. <https://doi.org/10.1097/TA.0b013e3182a1fea2>.
310. Paulich S, Lockey D. Resuscitative thoracotomy. *BJA Educ* 2020;20(7):242–8. <https://doi.org/10.1016/j.bjae.2020.03.005>.
311. Perkins ZB, Greenhalgh R, Ter Avest E, et al. Prehospital resuscitative thoracotomy for traumatic cardiac arrest. *JAMA Surg* 2025. <https://doi.org/10.1001/jamasurg.2024.7245>.
312. Cannon JW, Khan MA, Raja AS, et al. Damage control resuscitation in patients with severe traumatic hemorrhage: a practice management guideline from the Eastern Association for the Surgery of Trauma. *J Trauma Acute Care Surg* 2017;82(3):605–17. <https://doi.org/10.1097/TA.0000000000001333>.
313. Holcomb JB, Tilley BC, Baraniuk S, et al. Transfusion of plasma, platelets, and red blood cells in a 1:1:1 vs a 1:1:2 ratio and mortality in patients with severe trauma: the PROPPR randomized clinical trial. *JAMA* 2015;313(5):471–82. <https://doi.org/10.1001/jama.2015.12>.
314. Leibner E, Andreae M, Galvagno SM, Scalea T. Damage control resuscitation. *Clin Exp Emerg Med* 2020;7(1):5–13. <https://doi.org/10.15441/ceem.19.089>.
315. Nolan JP, Ohshima S, Nabecker S, et al. Cardiac arrest in the Cardiac Intervention Laboratory a Scoping Review: Consensus on Science with Treatment Recommendations. [Internet] Brussels, Belgium: International Liaison Committee on Resuscitation (ILCOR) Advanced Life Support Task Force. <http://ilcor.org>.

316. Sharma R, Bews H, Mahal H, et al. In-hospital cardiac arrest in the cardiac catheterization laboratory: effective transition from an ICU to CCU-Led resuscitation team. *J Interv Cardiol* 2019;2019:1686350. <https://doi.org/10.1155/2019/1686350>.
317. Sprung J, Ritter MJ, Rihal CS, et al. Outcomes of cardiopulmonary resuscitation and predictors of survival in patients undergoing coronary angiography including percutaneous coronary interventions. *Anesth Analg* 2006;102(1):217–24. <https://doi.org/10.1213/01.ane.0000189082.54614.26>.
318. Addala S, Kahn JK, Moccia TF, et al. Outcome of ventricular fibrillation developing during percutaneous coronary interventions in 19,497 patients without cardiogenic shock. *Am J Cardiol* 2005;96(6):764–5. <https://doi.org/10.1016/j.amjcard.2005.04.057>.
319. Webb JG, Solankhi NK, Chugh SK, et al. Incidence, correlates, and outcome of cardiac arrest associated with percutaneous coronary intervention. *Am J Cardiol* 2002;90(11):1252–4. [https://doi.org/10.1016/s0002-9149\(02\)02846-1](https://doi.org/10.1016/s0002-9149(02)02846-1).
320. Huang JL, Ting CT, Chen YT, Chen SA. Mechanisms of ventricular fibrillation during coronary angioplasty: increased incidence for the small orifice caliber of the right coronary artery. *Int J Cardiol* 2002;82(3):221–8. [https://doi.org/10.1016/s0167-5273\(01\)00596-4](https://doi.org/10.1016/s0167-5273(01)00596-4).
321. Mehta RH, Harjai KJ, Grines L, et al. Sustained ventricular tachycardia or fibrillation in the cardiac catheterization laboratory among patients receiving primary percutaneous coronary intervention: incidence, predictors, and outcomes. *J Am Coll Cardiol* 2004;43(10):1765–72. <https://doi.org/10.1016/j.jacc.2003.09.072>.
322. Mehta RH, Starr AZ, Lopes RD, et al. Incidence of and outcomes associated with ventricular tachycardia or fibrillation in patients undergoing primary percutaneous coronary intervention. *JAMA* 2009;301(17):1779–89. <https://doi.org/10.1001/jama.2009.600>.
323. Henriques JP, Gheeraert PJ, Ottervanger JP, et al. Ventricular fibrillation in acute myocardial infarction before and during primary PCI. *Int J Cardiol* 2005;105(3):262–6. <https://doi.org/10.1016/j.ijcard.2004.12.044>.
324. Giglioli C, Margheri M, Valente S, et al. Timing, setting and incidence of cardiovascular complications in patients with acute myocardial infarction submitted to primary percutaneous coronary intervention. *Can J Cardiol* 2006;22(12):1047–52. [https://doi.org/10.1016/s0828-282x\(06\)70320-8](https://doi.org/10.1016/s0828-282x(06)70320-8).
325. Demidova MM, Carlson J, Erlinge D, Platonov PG. Predictors of ventricular fibrillation at reperfusion in patients with acute ST-elevation myocardial infarction treated by primary percutaneous coronary intervention. *Am J Cardiol* 2015;115(4):417–22. <https://doi.org/10.1016/j.amjcard.2014.11.025>.
326. Elkaryoni A, Tran AT, Saad M, et al. Patient characteristics and survival outcomes of cardiac arrest in the cardiac catheterization laboratory: insights from get with the Guidelines(R)-Resuscitation registry. *Resuscitation* 2022;180:121–7. <https://doi.org/10.1016/j.resuscitation.2022.08.002>.
327. Al-Mukhtar O, Bilgrami I, Noaman S, et al. Cardiac arrest in the cardiac catheterization laboratory: initial experience with the role of simulation setup and training. *Am J Med Qual* 2021;36(4):238–46. <https://doi.org/10.1177/1062860620950805>.
328. Doorey AJ, Turi ZG, Lazzara EH, et al. Safety gaps in medical team communication: closing the loop on quality improvement efforts in the cardiac catheterization lab. *Catheter Cardiovasc Interv* 2022;99(7):1953–62. <https://doi.org/10.1002/ccd.30189>.
329. Lindsay AC, Bishop J, Harron K, Davies S, Haxby E. Use of a safe procedure checklist in the cardiac catheterisation laboratory. *BMJ Open Qual* 2018;7(3):e000074. <https://doi.org/10.1136/bmjog-2017-000074>.
330. Agricola E, Meucci F, Ancona F, Pardo Sanz A, Zamorano JL. Echocardiographic guidance in transcatheter structural cardiac interventions. *EuroIntervention* 2022;17(15):1205–26. <https://doi.org/10.4244/EIJ-D-21-00582>.
331. Dunning J, Archbold A, de Bono JP, et al. Joint British Societies' guideline on management of cardiac arrest in the cardiac catheter laboratory. *Heart* 2022;108(12):e3.
332. Reynolds JC, Nicholson TC, O'Neil BJ, et al. Diagnostic test accuracy with point-of-care ultrasound during cardiopulmonary resuscitation to indicate the etiology of cardiac arrest: consensus on science with treatment recommendations. [Internet] Brussels, Belgium: International Liaison Committee on Resuscitation (ILCOR) Advance Life Support Task Force. <http://ilcor.org>.
333. Mazzeffi M, Zaaqoq A, Curley J, et al. Survival after extracorporeal cardiopulmonary resuscitation based on in-hospital cardiac arrest and cannulation location: an analysis of the extracorporeal life support organization registry. *Crit Care Med* 2024;52(12):1906–17. <https://doi.org/10.1097/CCM.0000000000006439>.
334. ILCOR CoSTR. <https://costr.ilcor.org/>.
335. WHO. Global status report on drowning prevention 2024. World Health Organization; 2024.
336. Peden AE, Daw S, Lawes JC. Preliminary evaluation of the impact of mandatory life jacket laws at declared high-risk rock platforms on unintentional rock fishing drowning deaths. *Inj Prev* 2022;28(6):560–3. <https://doi.org/10.1136/ip-2022-044724>.
337. De Buck E, Vanhove ACOD, Veys K, Lang E, Vandekerckhove P. Day care as a strategy for drowning prevention in children under 6 years of age in low- and middle-income countries. *Cochrane Database Syst Rev* 2021;4(4):CD014955. <https://doi.org/10.1002/14651858.CD014955>.
338. Bierens J, Hoogenboezem J. Fatal drowning statistics from the Netherlands - an example of an aggregated demographic profile. *BMC Public Health* 2022;22(1):339. <https://doi.org/10.1186/s12889-022-12620-3>.
339. Queiroga AC, Seabra R, Franklin RC, Peden AE. Trends in drowning mortality in Portugal from 1992 to 2019: comparing Global Burden of Disease and national data. *Inj Prev* 2022;28(4):318–24. <https://doi.org/10.1136/injuryprev-2021-044415>.
340. Peden AE, Passmore J, Queiroga AC, Sweeney R, Jagnoor J. Closing the gap for drowning prevention across Europe. *Lancet Public Health* 2022;7(9):e728–9. [https://doi.org/10.1016/S2468-2667\(22\)00193-1](https://doi.org/10.1016/S2468-2667(22)00193-1).
341. Koon W, Clemens T, Bierens J, Quan L. Studying outcome predictors of drowning at the scene: why do we have so few answers?. *Am J Emerg Med* 2021;46:361–6. <https://doi.org/10.1016/j.ajem.2020.10.011>.
342. Bierens J, Abelairas-Gomez C, Barcala Furelos R, et al. Resuscitation and emergency care in drowning: a scoping review. *Resuscitation* 2021;162:205–17. <https://doi.org/10.1016/j.resuscitation.2021.01.033>.
343. Wyckoff MH, Singletary EM, Soar J, et al. 2021 international consensus on cardiopulmonary resuscitation and emergency cardiovascular care science with treatment recommendations: summary from the basic life support; Advanced life support; Neonatal life support; Education, implementation, and teams; First aid task forces; and the COVID-19 working group. *Resuscitation* 2022;169:229–311. <https://doi.org/10.1016/j.resuscitation.2021.10.040>.
344. International Life Saving Federation. <https://www.ilsf.org/>.
345. van Beeck EF, Branche CM, Szpilman D, Modell JH, Bierens JJ. A new definition of drowning: towards documentation and prevention of a global public health problem. *Bull World Health Organ* 2005;83(11):853–6. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/16302042>.
346. Abelairas-Gomez C, Tipton MJ, Gonzalez-Salvado V, Bierens J. Drowning: epidemiology, prevention, pathophysiology, resuscitation, and hospital treatment. *Emergencias* 2019;31(4):270–80. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/31347808>.

347. Bierens JJ, Lunetta P, Tipton M, Warner DS. Physiology of drowning: a review. *Physiology* (Bethesda) 2016;31(2):147–66. <https://doi.org/10.1152/physiol.00002.2015>.
348. Quan L, Bierens JJ, Lis R, Rowhani-Rahbar A, Morley P, Perkins GD. Predicting outcome of drowning at the scene: a systematic review and meta-analyses. *Resuscitation* 2016;104:63–75. <https://doi.org/10.1016/j.resuscitation.2016.04.006>.
349. Olasveengen TM, Mancini ME, Perkins GD, et al. Adult basic life support: 2020 international consensus on cardiopulmonary resuscitation and emergency cardiovascular care science with treatment recommendations. *Resuscitation* 2020;156:A35–79. <https://doi.org/10.1016/j.resuscitation.2020.09.010>.
350. Berger S, Siekmeyer M, Petzold-Quinque S, Kiess W, Merckenschlager A. Drowning and nonfatal drowning in children and adolescents: a subsequent retrospective data analysis. *Children* (Basel) 2024;11(4). <https://doi.org/10.3390/children11040439>.
351. Reizine F, Michelet P, Delbove A, et al. Development and validation of a clinico-biological score to predict outcomes in patients with drowning-associated cardiac arrest. *Am J Emerg Med* 2024;81:69–74. <https://doi.org/10.1016/j.ajem.2024.04.032>.
352. Shattock MJ, Tipton MJ. 'Autonomic conflict': a different way to die during cold water immersion?. *J Physiol* 2012;590(14):3219–30. <https://doi.org/10.1113/jphysiol.2012.229864>.
353. Tipton MJ, Collier N, Massey H, Corbett J, Harper M. Cold water immersion: kill or cure?. *Exp Physiol* 2017;102(11):1335–55. <https://doi.org/10.1113/EP086283>.
354. Tipton MJ, Golden FS. A proposed decision-making guide for the search, rescue and resuscitation of submersion (head under) victims based on expert opinion. *Resuscitation* 2011;82(7):819–24. <https://doi.org/10.1016/j.resuscitation.2011.02.021>.
355. Romlin BS, Winberg H, Janson M, et al. Excellent outcome with extracorporeal membrane oxygenation after accidental profound hypothermia (13.8 degrees C) and drowning. *Crit Care Med* 2015;43(11):e521–5. <https://doi.org/10.1097/CCM.0000000000001283>.
356. Barcala-Furelos R, Graham D, Abelairas-Gomez C, Rodriguez-Nunez A. Lay-rescuers in drowning incidents: a scoping review. *Am J Emerg Med* 2021;44:38–44. <https://doi.org/10.1016/j.ajem.2021.01.069>.
357. ILS. Medical position statement - MPS 20. Ability to perform CPR post rescue. International Life Saving Federation. <https://medical.ilsf.org/ability-to-perform-cpr-post-rescue/>.
358. Claesson A, Karlsson T, Thoren AB, Herlitz J. Delay and performance of cardiopulmonary resuscitation in surf lifeguards after simulated cardiac arrest due to drowning. *Am J Emerg Med* 2011;29(9):1044–50. <https://doi.org/10.1016/j.ajem.2010.06.026>.
359. Abelairas-Gomez C, Romo-Perez V, Barcala-Furelos R, Palacios-Aguilar J. Effect of lifeguard fatigue on the first 4 minutes of cardiopulmonary resuscitation after water rescue. *Emergencias* 2013;25(3):184–90 [In Spanish] (<Go to ISI>://WOS:000320143900005).
360. Thom O, Roberts K, Leggat PA, Devine S, Peden AE, Franklin R. Cervical spine immobilisation is only required in drowning patients at high risk of axial loading of the spine. *Emerg Med Australas* 2023;35(1):18–24. <https://doi.org/10.1111/1742-6723.14036>.
361. Breindahl N, Bierens JLM, Wiberg S, Barcala-Furelos R, Maschmann C. Prehospital guidelines on in-water traumatic spinal injuries for lifeguards and prehospital emergency medical services: an international Delphi consensus study. *Scand J Trauma Resusc Emerg Med* 2024;32(1):76. <https://doi.org/10.1186/s13049-024-01249-3>.
362. Bierens J, Bray J, Abelairas-Gomez C, et al. A systematic review of interventions for resuscitation following drowning. *Resusc Plus* 2023;14:100406. <https://doi.org/10.1016/j.resplu.2023.100406>.
363. Kaneto Y, Owada H, Kamikura T, Nakashima K, Ushimoto T, Inaba H. Advantages of bystander-performed conventional cardiopulmonary resuscitation in out-of-hospital cardiac arrest presumably caused by drowning in Japan: a propensity score-matching analysis using an extended nationwide database. *BMJ Open* 2024;14(5):e080579. <https://doi.org/10.1136/bmjopen-2023-080579>.
364. Tobin JM, Ramos WD, Greenshields J, et al. Outcome of conventional bystander cardiopulmonary resuscitation in cardiac arrest following drowning. *Prehosp Disaster Med* 2020;35(2):141–7. <https://doi.org/10.1017/S1049023X20000060>.
365. Berg KM, Bray JE, Ng KC, et al. 2023 international consensus on cardiopulmonary resuscitation and emergency cardiovascular care science with treatment recommendations: summary from the basic life support; Advanced life support; Pediatric life support; Neonatal life support; Education, implementation, and teams; and first aid task forces. *Resuscitation* 2024;195:109992. <https://doi.org/10.1016/j.resuscitation.2023.109992>.
366. Messelink DM, van der Ploeg GJ, van der Linden T, Flameling RD, Bierens J. Medical emergencies at sea: an analysis of ambulance-supported and autonomously performed operations by lifeboat crews. *BMC Emerg Med* 2023;23(1):108. <https://doi.org/10.1186/s12873-023-00879-7>.
367. Seesink J, Nieuwenburg SAV, van der Linden T, Bierens J. Circumstances, outcome and quality of cardiopulmonary resuscitation by lifeboat crews: why not always use an AED?. *Resuscitation* 2022;176:53–4. <https://doi.org/10.1016/j.resuscitation.2022.05.011>.
368. Dezfulian C, McCallin TE, Bierens J, et al. 2024 American heart association and american academy of pediatrics focused update on special circumstances: resuscitation following drowning: an update to the american heart association guidelines for cardiopulmonary resuscitation and emergency cardiovascular care. *Circulation* 2024;150(23):e501–16. <https://doi.org/10.1161/CIR.0000000000001274>.
369. Yao Y, DiNenna MA, Chen L, Jin S, He S, He J. Hypothesized mechanisms of death in swimming: a systematic review. *BMC Sports Sci Med Rehabil* 2024;16(1):6. <https://doi.org/10.1186/s13102-023-00799-w>.
370. Thom O, Roberts K, Devine S, Leggat PA, Franklin RC. Impact of lifeguard oxygen therapy on the resuscitation of drowning victims: results from an utstein style for drowning study. *Emerg Med Australas* 2024;36(6):841–8. <https://doi.org/10.1111/1742-6723.14454>.
371. Yoshimura S, Kiguchi T, Nishioka N, et al. Association of pre-hospital tracheal intubation with outcomes after out-of-hospital cardiac arrest by drowning comparing to supraglottic airway device: a nationwide propensity score-matched cohort study. *Resuscitation* 2024;197:110129. <https://doi.org/10.1016/j.resuscitation.2024.110129>.
372. Topjian AA, Berg RA, Bierens JJ, et al. Brain resuscitation in the drowning victim. *Neurocrit Care* 2012;17(3):441–67. <https://doi.org/10.1007/s12028-012-9747-4>.
373. Bitzer K, Breindahl N, Kelly B, et al. The role of accidental hypothermia in drowning patients with out-of-hospital cardiac arrest: a nationwide registry-based cohort study. *Resuscitation* 2025;207:110486. <https://doi.org/10.1016/j.resuscitation.2024.110486>.
374. Tipton M. Cold water immersion. In: Tipton M, Wooler A, editors. *The Science of Beach Lifeguarding*. Boca Raton, FL: Taylor & Francis Group; 2016. p. 87–98.
375. Nolan JP, Armstrong RA, Kane AD, et al. Advanced life support interventions during intra-operative cardiac arrest among adults as reported to the 7th national audit project of the royal college of anaesthetists. *Anaesthesia* 2024;79(9):914–23. <https://doi.org/10.1111/anae.16310>.
376. Penketh J, Nolan JP. Intra-operative cardiac arrest – We need to do better. *Anaesthesia* 2025;80(1):3–8. <https://doi.org/10.1111/anae.16487>.

377. Sunborger M, Jakobsson JG. Peri-operative cardiac arrests in Sweden 2013-2022: data analysis of incidence and trends. *Anaesthesia* 2024;79(10):1128–9. <https://doi.org/10.1111/anae.16396>.
378. Morais AC, Braz JRC, Soares JVA, et al. Eighteen-year trends in the rates of intra-operative cardiac arrest and associated mortality at a public university hospital in Brazil. *Anaesthesia* 2025;80(1):18–28. <https://doi.org/10.1111/anae.16450>.
379. Moppett IK, Kane AD, Armstrong RA, Kursumovic E, Soar J, Cook TM. Peri-operative cardiac arrest in the older frail patient as reported to the 7th national audit project of the royal college of anaesthetists. *Anaesthesia* 2024;79(8):810–20. <https://doi.org/10.1111/anae.16267>.
380. Tiradentes TAA, Einav S, Braz JRC, et al. Global anaesthesia-related cardiac arrest rates in children: a systematic review and meta-analysis. *Br J Anaesth* 2023;131(5):901–13. <https://doi.org/10.1016/j.bja.2023.08.023>.
381. Armstrong RA, Soar J, Kane AD, et al. Peri-operative cardiac arrest: epidemiology and clinical features of patients analysed in the 7th national audit project of the royal college of anaesthetists. *Anaesthesia* 2024;79(1):18–30. <https://doi.org/10.1111/anae.16156>.
382. Moitra VK, Einav S, Thies KC, et al. Cardiac arrest in the operating room: resuscitation and management for the anesthesiologist: part 1. *Anesth Analg* 2018;126(3):876–88. <https://doi.org/10.1213/ane.0000000000002596>.
383. Aziz S, Barratt J, Starr Z, et al. The association between intra-arrest arterial blood pressure and return of spontaneous circulation in out-of-hospital cardiac arrest. *Resuscitation* 2024;205:110426. <https://doi.org/10.1016/j.resuscitation.2024.110426>.
384. Harper NJN, Nolan JP, Soar J, Cook TM. Why chest compressions should start when systolic arterial blood pressure is below 50 mm Hg in the anaesthetised patient. *Br J Anaesth* 2020;124(3):234–8. <https://doi.org/10.1016/j.bja.2019.11.005>.
385. Morgan RW, Reeder RW, Bender D, et al. Associations between end-tidal carbon dioxide during pediatric cardiopulmonary resuscitation, cardiopulmonary resuscitation quality, and survival. *Circulation* 2024;149(5):367–78. <https://doi.org/10.1161/circulationaha.123.066659>.
386. Sheak KR, Wiebe DJ, Leary M, et al. Quantitative relationship between end-tidal carbon dioxide and CPR quality during both in-hospital and out-of-hospital cardiac arrest. *Resuscitation* 2015;89:149–54. <https://doi.org/10.1016/j.resuscitation.2015.01.026>.
387. Skulec R, Vojtisek P, Cerny V. Correlation between end-tidal carbon dioxide and the degree of compression of heart cavities measured by transthoracic echocardiography during cardiopulmonary resuscitation for out-of-hospital cardiac arrest. *Crit Care* 2019;23(1):334. <https://doi.org/10.1186/s13054-019-2607-2>.
388. Soar J. Epinephrine for cardiac arrest: knowns, unknowns and controversies. *Curr Opin Crit Care* 2020;26(6):590–5. <https://doi.org/10.1097/mcc.0000000000000763>.
389. Hinkelbein J, Andres J, Böttiger BW, et al. Cardiac arrest in the perioperative period: a consensus guideline for identification, treatment, and prevention from the European society of anaesthesiology and intensive care and the european society for trauma and emergency surgery. *Eur J Anaesthesiol* 2023;40(10):724–36. <https://doi.org/10.1097/eja.0000000000001813>.
390. Hernández-Tejedor A, González Puebla V, Corral Torres E, Benito Sánchez A, Pinilla López R, Galán Calategui MD. Ventilatory improvement with mechanical ventilator versus bag in non-traumatic out-of-hospital cardiac arrest: SYMEVECA study, phase 1. *Resuscitation* 2023;192:109965. <https://doi.org/10.1016/j.resuscitation.2023.109965>.
391. Malinverni S, Wilmin S, de Longueville D, et al. A retrospective comparison of mechanical cardio-pulmonary ventilation and manual bag valve ventilation in non-traumatic out-of-hospital cardiac arrests: a study from the Belgian cardiac arrest registry. *Resuscitation* 2024;199:110203. <https://doi.org/10.1016/j.resuscitation.2024.110203>.
392. Orso D, Vetrugno L, Federici N, et al. Mechanical ventilation management during mechanical chest compressions. *Respir Care* 2021;66(2):334–46. <https://doi.org/10.4187/respcare.07775>.
393. Tangpaisarn T, Tosibphanom J, Sata R, Kotruchin P, Drumheller B, Phungoen P. The effects of mechanical versus bag-valve ventilation on gas exchange during cardiopulmonary resuscitation in emergency department patients: a randomized controlled trial (CPR-VENT). *Resuscitation* 2023;193:109966. <https://doi.org/10.1016/j.resuscitation.2023.109966>.
394. Anez C, Becerra-Bolaños Á, Vives-Lopez A, Rodríguez-Pérez A. Cardiopulmonary resuscitation in the prone position in the operating room or in the intensive care unit: a systematic review. *Anesth Analg* 2021;132(2):285–92. <https://doi.org/10.1213/ane.0000000000005289>.
395. Berg K, Hsu CH, Considine J, et al. Cardiopulmonary Resuscitation and Defibrillation for Cardiac Arrest when Patients are in the Prone Position Consensus on Science with Treatment Recommendations [Internet]. Brussels, Belgium: International Liaison Committee on Resuscitation (ILCOR) Advanced Life Support and Basic Life Task Forces. <https://costr.ilcor.org/document/prone-cpr-als-systematic-review>.
396. Malik N, Claus PL, Illman JE, et al. Air embolism: diagnosis and management. *Future Cardiol* 2017;13(4):365–78. <https://doi.org/10.2217/fca-2017-0015>.
397. Marsh PL, Moore EE, Moore HB, et al. Iatrogenic air embolism: pathoanatomy, thromboinflammation, endotheliopathy, and therapies. *Front Immunol* 2023;14:1230049. <https://doi.org/10.3389/fimmu.2023.1230049>.
398. Rahman ZU, Murtaza G, Pourmorteza M, et al. Cardiac arrest as a consequence of air embolism: a case report and literature review. *Case Rep Med* 2016;2016:8236845. <https://doi.org/10.1155/2016/8236845>.
399. Castellini G, Gianola S, Biffi A, et al. Resuscitative endovascular balloon occlusion of the aorta (REBOA) in patients with major trauma and uncontrolled haemorrhagic shock: a systematic review with meta-analysis. *World J Emerg Surg* 2021;16(1):41. <https://doi.org/10.1186/s13017-021-00386-9>.
400. Ye Y, Li J, Liu S, et al. Efficacy of resuscitative endovascular balloon occlusion of the aorta for hemorrhage control in patients with abnormally invasive placenta: a historical cohort study. *BMC Pregnancy Childbirth* 2023;23(1):333. <https://doi.org/10.1186/s12884-023-05649-8>.
401. Chan PS, Greif R, Anderson T, et al. Ten steps toward improving in-hospital cardiac arrest quality of care and outcomes. *Resuscitation* 2023;193:109996. <https://doi.org/10.1016/j.resuscitation.2023.109996>.
402. Farquharson B, Cortegiani A, Lauridsen KG, Yeung J, Greif R, Nabecker S. Teaching team competencies within resuscitation training: a systematic review. *Resusc Plus* 2024;19:100687. <https://doi.org/10.1016/j.resplu.2024.100687>.
403. Nabecker S, Nation K, Gilfoyle E, et al. Cognitive aids used in simulated resuscitation: a systematic review. *Resusc Plus* 2024;19:100675. <https://doi.org/10.1016/j.resplu.2024.100675>.
404. Nolan JP, Sandroni C, Cariou A, et al. European Resuscitation Council and European Society of Intensive Care Medicine Guidelines 2025: Post-resuscitation Care. *Resuscitation* 2025;215 (Suppl 1):110809.
405. Waldinger R, Weinberg G, Gitman M. Local anesthetic toxicity in the geriatric population. *Drugs Aging* 2020;37(1):1–9. <https://doi.org/10.1007/s40266-019-00718-0>.
406. Wolfe JW, Butterworth JF. Local anesthetic systemic toxicity: update on mechanisms and treatment. *Curr Opin Anaesthesiol* 2011;24(5):561–6. <https://doi.org/10.1097/ACO.0b013e32834a9394>.
407. Cao D, Heard K, Foran M, Koyfman A. Intravenous lipid emulsion in the emergency department: a systematic review of recent literature.

- J Emerg Med 2015;48(3):387–97. <https://doi.org/10.1016/j.jemermed.2014.10.009>.
408. Macfarlane AJR, Gitman M, Bornstein KJ, El-Boghdady K, Weinberg G. Updates in our understanding of local anaesthetic systemic toxicity: a narrative review. *Anaesthesia* 2021;76 (Suppl 1):27–39. <https://doi.org/10.1111/anae.15282>.
 409. Neal JM, Neal EJ, Weinberg GL. American society of regional anesthesia and pain medicine local anesthetic systemic toxicity checklist: 2020 version. *Reg Anesth Pain Med* 2021;46(1):81–2. <https://doi.org/10.1136/rapm-2020-101986>.
 410. Ozcan MS, Weinberg G. Intravenous lipid emulsion for the treatment of drug toxicity. *J Intensive Care Med* 2014;29(2):59–70. <https://doi.org/10.1177/0885066612445978>.
 411. Hayes BD, Gosselin S, Calello DP, et al. Systematic review of clinical adverse events reported after acute intravenous lipid emulsion administration. *Clin Toxicol (Phila)* 2016;54(5):365–404. <https://doi.org/10.3109/15563650.2016.1151528>.
 412. Neal JM, Woodward CM, Harrison TK. The American society of regional anesthesia and pain medicine checklist for managing local anesthetic systemic toxicity: 2017 version. *Reg Anesth Pain Med* 2018;43(2):150–3. <https://doi.org/10.1097/AAP.0000000000000726>.
 413. de Queiroz SM, Chassard D, Musard H, et al. Resuscitation with lipid, epinephrine, or both in levobupivacaine-induced cardiac toxicity in newborn piglets. *Br J Anaesth* 2014;112(4):729–34. <https://doi.org/10.1093/bjae/aet327>.
 414. Harvey M, Cave G. Bupivacaine-induced cardiac arrest: fat is good—is epinephrine really bad?. *Anesthesiology* 2009;111 (3):467–9. <https://doi.org/10.1097/ALN.0b013e3181afe083>.
 415. Harvey M, Cave G, Prince G, Lahner D. Epinephrine injection in lipid-based resuscitation from bupivacaine-induced cardiac arrest: transient circulatory return in rabbits. *Anesth Analg* 2010;111 (3):791–6. <https://doi.org/10.1213/ANE.0b013e3181e66050>.
 416. Hiller David B, Gregorio Guido D, Ripper R, et al. Epinephrine impairs lipid resuscitation from bupivacaine overdose: a threshold effect. *Anesthesiology* 2009;111(3):498–505. <https://doi.org/10.1097/ALN.0b013e3181afde0a>.
 417. Adam Z, Adam S, Everngam RL, et al. Resuscitation after cardiac surgery: results of an international survey. *Eur J Cardiothorac Surg* 2009;36(1):29–34. <https://doi.org/10.1016/j.ejcts.2009.02.050>.
 418. Gupta P, Rettiganti M, Jeffries HE, et al. Risk factors and outcomes of in-hospital cardiac arrest following pediatric heart operations of varying complexity. *Resuscitation* 2016;105:1–7. <https://doi.org/10.1016/j.resuscitation.2016.04.022>.
 419. Kim KM, Arghami A, Habib R, et al. The society of thoracic surgeons adult cardiac surgery database: 2022 update on outcomes and research. *Ann Thorac Surg* 2023;115(3):566–74. <https://doi.org/10.1016/j.athoracsur.2022.12.033>.
 420. LaPar DJ, Ghanta RK, Kern JA, et al. Hospital variation in mortality from cardiac arrest after cardiac surgery: an opportunity for improvement?. *Ann Thorac Surg* 2014;98(2):534–9. <https://doi.org/10.1016/j.athoracsur.2014.03.030>. discussion 539–40 [In eng].
 421. Vakil K, Kealhofer JV, Alraies MC, et al. Long-term outcomes of patients who had cardiac arrest after cardiac operations. *Ann Thorac Surg* 2016;102(2):512–7. <https://doi.org/10.1016/j.athoracsur.2016.01.092>.
 422. Clark SC, Dunning J, Alfieri OR, et al. EACTS guidelines for the use of patient safety checklists. *Eur J Cardiothorac Surg* 2012;41 (5):993–1004. <https://doi.org/10.1093/ejcts/ezs009>.
 423. Dunning J, Fabbri A, Kolh PH, et al. Guideline for resuscitation in cardiac arrest after cardiac surgery. *Eur J Cardiothorac Surg* 2009;36(1):3–28. <https://doi.org/10.1016/j.ejcts.2009.01.033>.
 424. Surgery SoTSTFoRAC. The Society of Thoracic Surgeons Expert Consensus for the Resuscitation of Patients Who Arrest After Cardiac Surgery. *Ann Thorac Surg* 2017;103(3):1005–1020. (In eng). <https://doi.org/10.1016/j.athoracsur.2016.10.033>.
 425. Akhtar W, Baston VR, Berman M, et al. British societies guideline on the management of emergencies in implantable left ventricular assist device recipients in transplant centres. *Intensive Care Med* 2024;50(4):493–501. <https://doi.org/10.1007/s00134-024-07382-y>.
 426. Moskowitz A, Pocock H, Lagina A, Ng KC, Scholefield BR, Zelop CM, Bray J, Rossano J, Johnson NJ, Dunning J, Olasveengen T, Raymond T, Morales DLS, Carlese A, Elias M, Berg KM, Drennan I; ILCOR Advanced Life Support, Basical Life Support, and Pediatric Life Support Task Forces. Resuscitation of patients with durable mechanical circulatory support with acutely altered perfusion or cardiac arrest: A scoping review. *Resuscitation*. 2024 Oct;203:110389. doi: 10.1016/j.resuscitation.2024.110389. Epub 2024 Sep 7. PMID: 39245405; PMCID: PMC11466699.
 427. Dunning J, Nandi J, Ariffin S, Jerstice J, Danitsch D, Levine A. The cardiac surgery advanced life support course (CALS): delivering significant improvements in emergency cardiothoracic care. *Ann Thorac Surg* 2006;81(5):1767–72. <https://doi.org/10.1016/j.athoracsur.2005.12.012>.
 428. Haynes AB, Weiser TG, Berry WR, et al. A surgical safety checklist to reduce morbidity and mortality in a global population. *N Engl J Med* 2009;360(5):491–9. <https://doi.org/10.1056/NEJMsa0810119>.
 429. Brand J, McDonald A, Dunning J. Management of cardiac arrest following cardiac surgery. *BJA Educ* 2018;18(1):16–22. <https://doi.org/10.1016/j.bjae.2017.11.002>.
 430. Lomivorotov VV, Efremov SM, Kirov MY, Fominskiy EV, Karaskov AM. Low-cardiac-output syndrome after cardiac surgery. *J Cardiothorac Vasc Anesth* 2017;31(1):291–308. <https://doi.org/10.1053/j.jvca.2016.05.029>.
 431. Pagano D, Milojevic M, Meesters MI, et al. 2017 EACTS/EACTA guidelines on patient blood management for adult cardiac surgery. *Eur J Cardiothorac Surg* 2018;53(1):79–111. <https://doi.org/10.1093/ejcts/ezx325>.
 432. Flachs Kampf FA, Wouters PF, Edvardsen T, et al. Recommendations for transoesophageal echocardiography: EACVI update 2014. *Eur Heart J Cardiovasc. Imaging* 2014;15(4):353–65. <https://doi.org/10.1093/ehjci/ieu015>.
 433. Teran F, Prats MI, Nelson BP, et al. Focused transesophageal echocardiography during cardiac arrest resuscitation: JACC review topic of the week. *J Am Coll Cardiol* 2020;76(6):745–54. <https://doi.org/10.1016/j.jacc.2020.05.074>.
 434. Peretto G, Durante A, Limite LR, Cianflone D. Postoperative arrhythmias after cardiac surgery: incidence, risk factors, and therapeutic management. *Cardiol Res Pract* 2014;2014:615987. <https://doi.org/10.1155/2014/615987>.
 435. Lockowandt U, Levine A, Strang T, Dunning J. If a patient arrests after cardiac surgery is it acceptable to delay cardiopulmonary resuscitation until you have attempted either defibrillation or pacing?. *Interact Cardiovasc Thorac Surg* 2008;7(5):878–85. <https://doi.org/10.1510/icvts.2008.182980>.
 436. Richardson L, Dissanayake A, Dunning J. What cardioversion protocol for ventricular fibrillation should be followed for patients who arrest shortly post-cardiac surgery?. *Interact Cardiovasc Thorac Surg* 2007;6(6):799–805. <https://doi.org/10.1510/icvts.2007.163899>.
 437. International Liaison Committee on Resuscitation. 2005 International Consensus on Cardiopulmonary Resuscitation and Emergency Cardiovascular Care Science with Treatment Recommendations. Part 4: Advanced life support. *Resuscitation* 2005;67(2-3):213–47. (In eng). <https://doi.org/10.1016/j.resuscitation.2005.09.018>.
 438. Twomey D, Das M, Subramanian H, Dunning J. Is internal massage superior to external massage for patients suffering a cardiac arrest after cardiac surgery?. *Interact Cardiovasc Thorac Surg* 2008;7 (1):151–6. <https://doi.org/10.1510/icvts.2007.170399>.
 439. Soar J, Donnino MW, Maconochie I, et al. 2018 international consensus on cardiopulmonary resuscitation and emergency cardiovascular care science with treatment recommendations summary. *Resuscitation* 2018;133:194–206. <https://doi.org/10.1016/j.resuscitation.2018.10.017>.

440. Anderson KL, Castaneda MG, Boudreau SM, Sharon DJ, Bebartá VS. Left ventricular compressions improve hemodynamics in a swine model of out-of-hospital cardiac arrest. *Prehosp Emerg Care* 2017;21(2):272–80. <https://doi.org/10.1080/10903127.2016.1241328>.
441. Fair 3rd J, Mallin MP, Adler A, et al. Transesophageal echocardiography during cardiopulmonary resuscitation is associated with shorter compression pauses compared with transthoracic echocardiography. *Ann Emerg Med* 2019;73(6):610–6. <https://doi.org/10.1016/j.annemergmed.2019.01.018>.
442. Gottlieb M, Alerhand S. Managing cardiac arrest using ultrasound. *Annals of Emergency Medicine* 2023;81(5):532–42. <https://doi.org/10.1016/j.annemergmed.2022.09.016>.
443. Hussein L, Rehman MA, Jelic T, et al. Transoesophageal echocardiography in cardiac arrest: a systematic review. *Resuscitation* 2021;168:167–75. <https://doi.org/10.1016/j.resuscitation.2021.08.001>.
444. Hwang SO, Jung WJ, Roh YI, Cha KC. Intra-arrest transesophageal echocardiography during cardiopulmonary resuscitation. *Clin Exp Emerg Med* 2022;9(4):271–80. <https://doi.org/10.15441/ceem.22.399>.
445. Long B, Alerhand S, Maliel K, Koyfman A. Echocardiography in cardiac arrest: an emergency medicine review. *Am J Emerg Med* 2018;36(3):488–93. <https://doi.org/10.1016/j.ajem.2017.12.031>.
446. Riendeau Beaulac G, Teran F, Lecluyse V, et al. Transesophageal echocardiography in patients in cardiac arrest: the heart and beyond. *Can J Cardiol* 2023;39(4):458–73. <https://doi.org/10.1016/j.cjca.2022.12.027>.
447. Teran F, Dean AJ, Centeno C, et al. Evaluation of out-of-hospital cardiac arrest using transesophageal echocardiography in the emergency department. *Resuscitation* 2019;137:140–7. <https://doi.org/10.1016/j.resuscitation.2019.02.013>.
448. Hwang SO, Zhao PG, Choi HJ, et al. Compression of the left ventricular outflow tract during cardiopulmonary resuscitation. *Acad Emerg Med* 2009;16(10):928–33. <https://doi.org/10.1111/j.1553-2712.2009.00497.x>.
449. Prager R, Bowdridge J, Pratte M, Cheng J, McInnes MD, Indications AR. Clinical impact, and complications of critical care transesophageal echocardiography: a scoping review. *J Intensive Care Med* 2023;38(3):245–72. <https://doi.org/10.1177/08850666221115348>.
450. Charalambous CP, Zipitis CS, Keenan DJ. Chest reexploration in the intensive care unit after cardiac surgery: a safe alternative to returning to the operating theater. *Ann Thorac Surg* 2006;81(1):191–4. <https://doi.org/10.1016/j.athoracsurg.2005.06.024>.
451. Mackay JH, Powell SJ, Osgathorp J, Rozario CJ. Six-year prospective audit of chest reopening after cardiac arrest. *Eur J Cardiothorac Surg* 2002;22(3):421–5. [https://doi.org/10.1016/s1010-7940\(02\)00294-4](https://doi.org/10.1016/s1010-7940(02)00294-4).
452. Dogan EM, Dogan EA, Nilsson KF, Edström M. Intra-aortic balloon pump synchronized with chest compressions improves outcome during cardiopulmonary resuscitation in experimental cardiac arrest. *Resuscitation* 2024;205:110433. <https://doi.org/10.1016/j.resuscitation.2024.110433>.
453. Zhao Y, Xing J, Du Z, Liu F, Jia M, Hou X. Extracorporeal cardiopulmonary resuscitation for adult patients who underwent post-cardiac surgery. *Eur J Med Res* 2015;20:83. <https://doi.org/10.1186/s40001-015-0179-4>.
454. Franklin BA, Thompson PD, Al-Zaiti SS, et al. Exercise-related acute cardiovascular events and potential deleterious adaptations following long-term exercise training: placing the risks into perspective—an update: a scientific statement from the american heart association. *Circulation* 2020;141(13):e705–36. <https://doi.org/10.1161/cir.0000000000000749>.
455. Marjion E, Bouguin W, Karam N, et al. Survival from sports-related sudden cardiac arrest: in sports facilities versus outside of sports facilities. *Am Heart J* 2015;170(2):339–345.e1. <https://doi.org/10.1016/j.ahj.2015.03.022>.
456. Ackerman M, Atkins DL, Triedman JK. Sudden cardiac death in the young. *Circulation* 2016;133(10):1006–26. <https://doi.org/10.1161/circulationaha.115.020254>.
457. Finocchiaro G, Papadakis M, Robertus JL, et al. Etiology of sudden death in sports: insights from a United Kingdom regional registry. *J Am Coll Cardiol* 2016;67(18):2108–15. <https://doi.org/10.1016/j.jacc.2016.02.062>.
458. Malhotra A, Dhutia H, Finocchiaro G, et al. Outcomes of cardiac screening in adolescent soccer players. *N Engl J Med* 2018;379(6):524–34. <https://doi.org/10.1056/NEJMoa1714719>.
459. Maron BJ, Haas TS, Doerer JJ, Thompson PD, Hodges JS. Comparison of U.S. and Italian experiences with sudden cardiac deaths in young competitive athletes and implications for preparticipation screening strategies. *Am J Cardiol* 2009;104(2):276–80. <https://doi.org/10.1016/j.amjcard.2009.03.037>.
460. Bohm P, Scharhag J, Meyer T. Data from a nationwide registry on sports-related sudden cardiac deaths in Germany. *Eur. J Prev Cardiol* 2016;23(6):649–56. <https://doi.org/10.1177/2047487315594087>.
461. Bohm P, Scharhag J, Egger F, et al. Sports-Related Sudden Cardiac Arrest in Germany. *Can J Cardiol* 2021;37(1):105–12. <https://doi.org/10.1177/2047487315594087>. doi: 10.1016/j.cjca.2020.03.021 [In eng].
462. Egger F, Scharhag J, Kästner A, Dvořák J, Bohm P, Meyer T. FIFA Sudden Death Registry (FIFA-SDR): a prospective, observational study of sudden death in worldwide football from 2014 to 2018. *Br J Sports Med* 2022;56(2):80–7. <https://doi.org/10.1136/bisports-2020-102368>.
463. Landry CH, Allan KS, Connelly KA, Cunningham K, Morrison LJ, Dorian P. Sudden cardiac arrest during participation in competitive sports. *N Engl J Med* 2017;377(20):1943–53. <https://doi.org/10.1056/NEJMoa1615710>.
464. Drezner JA, Toresdahl BG, Rao AL, Huszti E, Harmon KG. Outcomes from sudden cardiac arrest in US high schools: a 2-year prospective study from the national registry for AED use in sports. *Br J Sports Med* 2013;47(18):1179–83. <https://doi.org/10.1136/bisports-2013-092786>.
465. Smith CM, Moore F, Drezner JA, et al. Resuscitation on the field of play: a best-practice guideline from resuscitation council UK. *Br J Sports Med* 2024;58(19):1098–106. <https://doi.org/10.1136/bisports-2024-108440>.
466. Lott C, Bahtijarević Z, Klomp P, et al. Increasing CPR awareness in Europe through EURO 2024: lessons from “Get trained save lives”. Outcomes of cardiac screening in adolescent soccer players. *Resuscitation* 2025;208(6):110532. <https://doi.org/10.1016/j.resuscitation.2025.110532>.
467. Pelliccia A, Sharma S, Gati S, et al. 2020 ESC Guidelines on sports cardiology and exercise in patients with cardiovascular disease. *Eur Heart J* 2021;42(1):17–96. <https://doi.org/10.1093/eurheartj/ehaa605>.
468. Maron BJ, Zipes DP, Kovacs RJ. Eligibility and disqualification recommendations for competitive athletes with cardiovascular abnormalities: preamble, principles, and general considerations: a scientific statement from the american heart association and american college of cardiology. *J Am Coll Cardiol* 2015;66(21):2343–9. <https://doi.org/10.1016/j.jacc.2015.09.032>.
469. Mont L, Pelliccia A, Sharma S, et al. Pre-participation cardiovascular evaluation for athletic participants to prevent sudden death: Position paper from the EHRA and the EACPR, branches of the ESC. Endorsed by APhRS, HRS, and SOLAECE. *Eur. J Prev Cardiol* 2017;24(1):41–69. <https://doi.org/10.1177/2047487316676042>.
470. Kim JH, Martinez MW, Guseh JS, et al. A contemporary review of sudden cardiac arrest and death in competitive and recreational

- athletes. *Lancet* 2024;404(10468):2209–22. [https://doi.org/10.1016/s0140-6736\(24\)02086-5](https://doi.org/10.1016/s0140-6736(24)02086-5).
471. Armstrong M, Paternostro-Bayles M, Conroy MB, et al. Preparticipation screening prior to physical activity in community lifestyle interventions. Data from a nationwide registry on sports-related sudden cardiac deaths in Germany. *Transl J Am Coll Sports Med* 2018;3(22):176–80. <https://doi.org/10.1249/tjx.0000000000000073>. DOI: 10.1177/2047487315594087.
 472. Link MS, Wang PJ, Pandian NG, et al. An experimental model of sudden death due to low-energy chest-wall impact (commotio cordis). *N Engl J Med* 1998;338(25):1805–11. <https://doi.org/10.1056/nejm199806183382504>.
 473. Maron BJ, Gohman TE, Kyle SB, Estes 3rd NA, Link MS. Clinical profile and spectrum of commotio cordis. *Jama* 2002;287(9):1142–6. <https://doi.org/10.1001/jama.287.9.1142>.
 474. Shore E, Moseley GA, DeLong R, et al. Incidents and patterns of commotio cordis among athletes in the USA from 1982 to 2023. Resuscitation on the field of play: a best-practice guideline from resuscitation council UK. *Inj Prev* 2024;58(19):1098–106. <https://doi.org/10.1136/ip-2024-045374>. doi: 10.1136/bjsports-2024-108440 [In eng].
 475. Peng T, Derry LT, Yogeswaran V, Goldschlager NF. Commotio Cordis in 2023. *Sports Med* 2023;53(8):1527–36. <https://doi.org/10.1007/s40279-023-01873-6>.
 476. Maron BJ, Haas TS, Ahluwalia A, Garberich RF, Estes NA. 3rd, Link MS. Increasing survival rate from commotio cordis. *Heart Rhythm* 2013;10(2):219–23. <https://doi.org/10.1016/j.hrthm.2012.10.034>.
 477. Okorare O, Alugba G, Olusiji S, et al. Sudden cardiac death: an update on commotio cordis. *Cureus* 2023;15(4):e38087. <https://doi.org/10.7759/cureus.38087>.
 478. Dickey GJ, Bian K, Khan HR, Mao H. Developing commotio cordis injury metrics for baseball safety: unravelling the connection between chest force and rib deformation to left ventricle strain and pressure. *Comput Methods Biomech Biomed Engin* 2022;25(3):247–56. <https://doi.org/10.1080/10255842.2021.1948022>.
 479. Dickey GJ, Bian K, Islam SU, Khan HR, Rohr S, Mao H. Advancing commotio cordis safety standards using the total human models for safety (THUMS). *Ann Biomed Eng* 2023;51(9):2070–85. <https://doi.org/10.1007/s10439-023-03235-9>.
 480. Shah A. Using data for improvement. *BMJ* 2019;364:l189. <https://doi.org/10.1136/bmj.l189>.
 481. Mashoufi M, Ayatollahi H, Khorasani-Zavareh D. A review of data quality assessment in emergency medical services. *Open Med Inform J* 2018;12:19–32. <https://doi.org/10.2174/1874431101812010019>.
 482. Mashoufi M, Ayatollahi H, Khorasani-Zavareh D, Talebi Azad Boni T. Data quality assessment in emergency medical services: an objective approach. *BMC Emerg Med* 2023;23(1):10. <https://doi.org/10.1186/s12873-023-00781-2>.
 483. Riyapan S, Sanyanuban P, Chantanakomes J, et al. Enhancing survival outcomes in developing emergency medical service system: continuous quality improvement for out-of-hospital cardiac arrest. *Resusc Plus* 2024;19:100683.
 484. Wilson C, Howell AM, Janes G, Benn J. The role of feedback in emergency ambulance services: a qualitative interview study. *BMC Health Serv Res* 2022;22(1):296. <https://doi.org/10.1186/s12913-022-07676-1>.
 485. Wilson C, Janes G, Lawton R, Benn J. Types and effects of feedback for emergency ambulance staff: a systematic mixed studies review and meta-analysis. *BMJ Qual Saf* 2023;32(10):573–88.
 486. Ho AFW, Ong MEH. Transportation during and after cardiac arrest: who, when, how and where?. *Curr Opin Crit Care* 2021;27(3):223–31.
 487. Butterfield ED, Price J, Bonsano M, et al. Prehospital invasive arterial blood pressure monitoring in critically ill patients attended by a UK helicopter emergency medical service- a retrospective observational review of practice. *Scand J Trauma Resusc Emerg Med* 2024;32(1):20.
 488. Ringen E, Haugland H, Brede JR. Characteristics of the most severely ill and injured patients in a Norwegian helicopter emergency medical service: a retrospective cohort study. *BMC Emerg Med* 2024;24(1):35.
 489. Aziz S, Lachowycz K, Major R, Rees P, Barratt J. Implementation of advanced vascular access, physiological monitoring and goal-directed resuscitation during OHCA in a helicopter emergency medical service. *J Vasc Access* 2024;11297298241242157.
 490. Yin RT, Berve PO, Skaalhegg T, et al. Recovery of arterial blood pressure after chest compression pauses in patients with out-of-hospital cardiac arrest. *Resuscitation* 2024;201:110311.
 491. Morton S, Avery P, Payne J, Omeara M. Arterial blood gases and arterial lines in the prehospital setting. *Air Med J* 2022;41(2):201–8.
 492. Standiford C, Wassermann M, Lauria MJ. Initiation of invasive arterial pressure monitoring by critical care transport crews. *Air Med J* 2022;41(2):248–51.
 493. Perera Y, Raitt J, Poole K, Metcalfe D, Lewinsohn A. Non-invasive versus arterial pressure monitoring in the pre-hospital critical care environment: a paired comparison. *Scand J Trauma Resusc Emerg Med* 2024;32(1):77.
 494. Kamboj N, Chang K, Metcalfe K, Chu CH, Conway A. Accuracy and precision of continuous non-invasive arterial pressure monitoring in critical care: a systematic review. *Intensive Crit Care Nurs* 2021;67:103091.
 495. Keselica M, Peřan D, Renza M, et al. Efficiency of two-member crews in delivering prehospital advanced life support cardiopulmonary resuscitation: a scoping review. *Resusc Plus* 2024;18:100661.
 496. Graf J, Stüben U, Pump S. In-flight medical emergencies. *Dtsch Arztebl Int* 2012;109(37):591–601.
 497. Katoch T, Pinnamaneni S, Medatwal R, et al. Hearts in the sky: understanding the cardiovascular implications of air travel. *Future Cardiol* 2024;20(11–12):651–60.
 498. Borges do Nascimento IJ, Jerončić A, Arantes AJR, et al. The global incidence of in-flight medical emergencies: a systematic review and meta-analysis of approximately 1.5 billion airline passengers. *Am J Emerg Med* 2021;48:156–64.
 499. Battineni G, Arcese A, Chintalapudi N, Di Canio M, Sibilio F, Amenta F. Approaches to medical emergencies on commercial flights. *Medicina (Kaunas)* 2024;60(5).
 500. Hinkelbein J, Schmitz J, Kerkhoff S, et al. On-board emergency medical equipment of European airlines. *Travel Med Infect Dis* 2021;40:101982.
 501. Oliveira ATB. Worldwide regulation of the medical emergency kit and first aid kit. *Aerosp Med Hum Perform* 2024;95(6):321–6.
 502. Hinkelbein J, Kerkhoff S, Adler C, et al. Cardiopulmonary resuscitation (CPR) during spaceflight – A guideline for CPR in microgravity. *Scand J Trauma Resusc Emerg Med* 2020;28(1):108.
 503. Vanni V, Hu JL, Currao A, Savastano S. Chest compressions techniques in microgravity: light years from a good quality resuscitation. *Resuscitation Plus* 2025;100996. <https://doi.org/10.1016/j.resplu.2025.100996>.
 504. Overbeek R, Liebold F, Johnson Kolaparambil Varghese L, et al. Alternative techniques for cardiopulmonary resuscitation in extreme environments – A scoping review. *Resusc Plus* 2024;20:100762.
 505. Dillard JS, Maynard W, Kashyap R. The epidemiology of maritime patients requiring medical evacuation: a literature review. *Cureus* 2023;15(11):e49606.
 506. Mohammadzadeh N, Gholamzadeh M. Requirements, challenges, and key components to improve onboard medical care using maritime telemedicine: narrative review. *Int J Telemed Appl* 2023;2023:9389286.
 507. British Thoracic Society SIGN. <https://www.sign.ac.uk/media/2269/sign-158-2024-update-final.pdf>.
 508. Reddel HK, Bacharier LB, Bateman ED, et al. Global initiative for asthma strategy 2021: executive summary and rationale for key

- changes. *European Respiratory Journal* 2022;59(1):2102730. <https://doi.org/10.1183/13993003.02730-2021>.
509. (GOLD) Gifcold.
 510. Franklin C, Samuel J, Hu TC. Life-threatening hypotension associated with emergency intubation and the initiation of mechanical ventilation. *Am J Emerg Med* 1994;12(4):425–8. [https://doi.org/10.1016/0735-6757\(94\)90053-1](https://doi.org/10.1016/0735-6757(94)90053-1).
 511. Laher AE, Buchanan SK. Mechanically ventilating the severe asthmatic. *J Intensive Care Med* 2018;33(9):491–501. <https://doi.org/10.1177/0885066617740079>.
 512. Tonna JE, Abrams D, Brodie D, et al. Management of adult patients supported with venovenous extracorporeal membrane oxygenation (VV ECMO): guideline from the extracorporeal life support organization (ELSO). *Asaio J* 2021;67(6):601–10. <https://doi.org/10.1097/mat.0000000000001432>.
 513. Ekechukwu N, Batra S, Orsi D, Rahmanian M, Bangar M, Mohamed A. Outcomes of extracorporeal life support (ECLS) in acute severe asthma: a narrative review. *Lung* 2024;202(2):91–6. <https://doi.org/10.1007/s00408-023-00667-x>.
 514. Boucher A, Payen C, Garayt C, et al. Salbutamol misuse or abuse with fatal outcome: a case-report. *Hum Exp Toxicol* 2011;30(11):1869–71. <https://doi.org/10.1177/09603271110388957>.
 515. Lemaître RN, Siscovick DS, Psaty BM, et al. Inhaled beta-2 adrenergic receptor agonists and primary cardiac arrest. *Am J Med* 2002;113(9):711–6. [https://doi.org/10.1016/s0002-9343\(02\)01384-0](https://doi.org/10.1016/s0002-9343(02)01384-0).
 516. Rosero SZ, Zareba W, Moss AJ, et al. Asthma and the risk of cardiac events in the Long QT syndrome. Long QT Syndrome Investigative Group. *Am J Cardiol* 1999;84(12):1406–11. [https://doi.org/10.1016/s0002-9149\(99\)00586-x](https://doi.org/10.1016/s0002-9149(99)00586-x).
 517. Salpeter SR, Ormiston TM, Salpeter EE. Cardiovascular effects of beta-agonists in patients with asthma and COPD: a meta-analysis. *Chest* 2004;125(6):2309–21. <https://doi.org/10.1378/chest.125.6.2309>.
 518. van den Berg ME, Stricker BH, Brusselle GG, Lahousse L. Chronic obstructive pulmonary disease and sudden cardiac death: A systematic review. *Trends Cardiovasc Med* 2016;26(7):606–13. <https://doi.org/10.1016/j.tcm.2016.04.001>.
 519. Charlton NP BD, Djäv T, Carlson J on behalf of the International Liaison Committee on Resuscitation First Aid Task Force. Potential Harm from Bronchodilators for Respiratory Concerns First Aid Task Force Synthesis of a Scoping Review. 24 November ed2022.
 520. Granfeldt A, Wissenberg M, Hansen SM, et al. Severity of chronic obstructive pulmonary disease and presenting rhythm in patients with out-of-hospital cardiac arrest. *Resuscitation* 2018;126:111–7. <https://doi.org/10.1016/j.resuscitation.2018.03.006>.
 521. Herlitz J, Rosenfelt M, Bång A, et al. Prognosis among patients with out-of-hospital cardiac arrest judged as being caused by deterioration of obstructive pulmonary disease. *Resuscitation* 1996;32(3):177–84. [https://doi.org/10.1016/0300-9572\(96\)00970-Z](https://doi.org/10.1016/0300-9572(96)00970-Z).
 522. Hostetler MA, Davis CO. Bilateral localized tension pneumothoraces refractory to needle decompression. *Pediatr Emerg Care* 1999;15(5):322–4. <https://doi.org/10.1097/00006565-199910000-00005>.
 523. Williams-Johnson J, Williams EW, Hart N, Maycock C, Bullock K, Ramphal P. Simultaneous spontaneous bilateral pneumothoraces in an asthmatic. *West Indian Med J* 2008;57(5):508–10.
 524. Eason J, Tayler D, Cottam S, et al. Manual chest compression for total bronchospasm. *Lancet* 1991;337(8737):366. [https://doi.org/10.1016/0140-6736\(91\)90998-5](https://doi.org/10.1016/0140-6736(91)90998-5).
 525. Fisher MM, Bowey CJ, Ladd-Hudson K. External chest compression in acute asthma: a preliminary study. *Crit Care Med* 1989;17(7):686–7. <https://doi.org/10.1097/00003246-198907000-00018>.
 526. Fisher MM, Whaley AP, Pye RR. External chest compression in the management of acute severe asthma—a technique in search of evidence. *Prehosp Disaster Med* 2001;16(3):124–7. <https://doi.org/10.1017/s1049023x00025863>.
 527. Harrison R. Chest compression first aid for respiratory arrest due to acute asphyxic asthma. *Emerg Med J* 2010;27(1):59–61. <https://doi.org/10.1136/emj.2007.056119>.
 528. Myles PS, Weeks AM. Cardiopulmonary resuscitation in obstructive airways disease. *Lancet* 1993;341(8854):1217. [https://doi.org/10.1016/0140-6736\(93\)91048-g](https://doi.org/10.1016/0140-6736(93)91048-g).
 529. Rezoagli E, Bastia L, Brochard L, Bellani G. Physical manoeuvres in patients with ARDS and low compliance: bedside approaches to detect lung hyperinflation and optimise mechanical ventilation. *European Respiratory Journal* 2023;61(5):2202169. <https://doi.org/10.1183/13993003.02169-2022>.
 530. Rosengarten PL, Tuxen DV, Dziukas L, Scheinkestel C, Merrett K, Bowes G. Circulatory arrest induced by intermittent positive pressure ventilation in a patient with severe asthma. *Anaesth Intensive Care* 1991;19(1):118–21. <https://doi.org/10.1177/0310057x9101900126>.
 531. Leatherman JW, McArthur C, Shapiro RS. Effect of prolongation of expiratory time on dynamic hyperinflation in mechanically ventilated patients with severe asthma. *Crit Care Med* 2004;32(7):1542–5. <https://doi.org/10.1097/01.ccm.0000130993.43076.20>.
 532. Orea-Tejeda A, Gómez-Martínez M, González-Islas D, et al. The impact of hydration status and fluid distribution on pulmonary function in COPD patients. *Sci Rep* 2022;12(1):1216. <https://doi.org/10.1038/s41598-022-05192-0>.
 533. Yeo HJ, Kim D, Jeon D, Kim YS, Rycus P, Cho WH. Extracorporeal membrane oxygenation for life-threatening asthma refractory to mechanical ventilation: analysis of the extracorporeal life support organization registry. *Crit Care* 2017;21(1):297. <https://doi.org/10.1186/s13054-017-1886-8>.
 534. Kovesdy CP. Epidemiology of chronic kidney disease: an update 2022. *Kidney Int Suppl* (2011) 2022;12(1):7–11. <https://doi.org/10.1016/j.kisu.2021.11.003>.
 535. Borg R, Carlson N, Sondergaard J, Persson F. The growing challenge of chronic kidney disease: an overview of current knowledge. *Int J Nephrol* 2023;2023:9609266. <https://doi.org/10.1155/2023/9609266>.
 536. Obremaska M, Madziarska K, Zysko D, et al. Out-of-hospital cardiac arrest in dialysis patients. *Int Urol Nephrol* 2021;53(3):563–9. <https://doi.org/10.1007/s12555-020-02694-6>.
 537. Davis TR, Young BA, Eisenberg MS, Rea TD, Copass MK, Cobb LA. Outcome of cardiac arrests attended by emergency medical services staff at community outpatient dialysis centers. *Kidney Int* 2008;73(8):933–9. <https://doi.org/10.1038/sj.ki.5002749>.
 538. Karnik JA, Young BS, Lew NL, et al. Cardiac arrest and sudden death in dialysis units. *Kidney Int* 2001;60(1):350–7. <https://doi.org/10.1046/j.1523-1755.2001.00806.x>.
 539. Lafrance JP, Nolin L, Senecal L, Leblanc M. Predictors and outcome of cardiopulmonary resuscitation (CPR) calls in a large haemodialysis unit over a seven-year period. *Nephrol Dial Transplant* 2006;21(4):1006–12. <https://doi.org/10.1093/ndt/gfk007>.
 540. Foley RN, Gilbertson DT, Murray T, Collins AJ. Long interdialytic interval and mortality among patients receiving hemodialysis. *N Engl J Med* 2011;365(12):1099–107. <https://doi.org/10.1056/NEJMoa1103313>.
 541. Pun PH. Listening to the rhythm of arrhythmias among patients maintained on hemodialysis. *Kidney Med* 2024;6(4):100803. <https://doi.org/10.1016/j.xkme.2024.100803>.
 542. Genovesi S, Boriani G, Covic A, et al. Sudden cardiac death in dialysis patients: different causes and management strategies. *Nephrol Dial Transplant* 2021;36(3):396–405. <https://doi.org/10.1093/ndt/gfz182>.

543. Soomro QH, Koplan BA, Costea AI, et al. Arrhythmia and time of day in maintenance hemodialysis: secondary analysis of the monitoring in dialysis study. *Kidney Med* 2024;6(4):100799. <https://doi.org/10.1016/j.xkme.2024.100799>.
544. United States Renal Data System (USRDS) Annual Data Report 2024: Epidemiology of kidney disease in the United States. National Institutes of Health, National Institute of Diabetes and Digestive and Kidney Diseases, Bethesda, MD.
545. Makar MS, Pun PH. Sudden cardiac death among hemodialysis patients. *Am J Kidney Dis* 2017;69(5):684–95. <https://doi.org/10.1053/j.ajkd.2016.12.006>.
546. Pun PH, Dupre ME, Starks MA, et al. Outcomes for hemodialysis patients given cardiopulmonary resuscitation for cardiac arrest at outpatient dialysis clinics. *J Am Soc Nephrol* 2019. <https://doi.org/10.1681/ASN.2018090911>.
547. Wan C, Herzog CA, Zareba W, Szymkiewicz SJ. Sudden cardiac arrest in hemodialysis patients with wearable cardioverter defibrillator. *Ann Noninvasive Electrocardiol* 2014;19(3):247–57. <https://doi.org/10.1111/anec.12119>.
548. Yang W, Kim JG, Kang GH, et al. Prognostic effect of underlying chronic kidney disease and renal replacement therapy on the outcome of patients after out-of-hospital cardiac arrest: a nationwide observational study. *Medicina (Kaunas)* 2022;58(3). <https://doi.org/10.3390/medicina58030444>.
549. Hsieh MS, Chattopadhyay A, Lu TP, et al. Effect of end-stage kidney disease on the return of spontaneous circulation in Taiwanese adults with out-of-hospital cardiac arrest. *Sci Rep* 2023;13(1):7905. <https://doi.org/10.1038/s41598-023-35024-8>.
550. Moss AH, Holley JL, Upton MB. Outcomes of cardiopulmonary resuscitation in dialysis patients. *J Am Soc Nephrol* 1992;3(6):1238–43.
551. Starks MA, Wu J, Peterson ED, et al. In-hospital cardiac arrest resuscitation practices and outcomes in maintenance dialysis patients. *Clin J Am Soc Nephrol* 2020. <https://doi.org/10.2215/CJN.05070419>.
552. Saeed F, Murad HF, Wing RE, Li J, Schold JD, Fiscella KA. Outcomes following in-hospital cardiopulmonary resuscitation in people receiving maintenance dialysis. *Kidney Med* 2022;4(1):100380. <https://doi.org/10.1016/j.xkme.2021.08.014>.
553. Sharma S, Raman P, Sinha M, Deo AS. Factors affecting outcomes of cardiopulmonary resuscitation in a nephro-urology unit: a retrospective analysis. *Indian J Crit Care Med* 2022;26(3):322–6. <https://doi.org/10.5005/ip-journals-10071-24146>.
554. Wong SP, Kreuter W, Curtis JR, Hall YN, O'Hare AM. Trends in in-hospital cardiopulmonary resuscitation and survival in adults receiving maintenance dialysis. *JAMA Intern Med* 2015;175(6):1028–35. <https://doi.org/10.1001/jamainternmed.2015.0406>.
555. Saeed F, Adil MM, Malik AA, Schold JD, Holley JL. Outcomes of in-hospital cardiopulmonary resuscitation in maintenance dialysis patients. *J Am Soc Nephrol* 2015;26(12):3093–101. <https://doi.org/10.1681/ASN.2014080766>.
556. Pun PH. In-hospital cardiopulmonary resuscitation in patients receiving maintenance dialysis: glass half full or half empty?. *Kidney Med* 2022;4(1):100399. <https://doi.org/10.1016/j.xkme.2021.10.012>.
557. Workgroup KD. K/DOQI clinical practice guidelines for cardiovascular disease in dialysis patients. *Am J Kidney Dis* 2005;45(4 Suppl 3):S1–153. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/15806502>.
558. Leirich RW, Pun PH, Tanenbaum ND, Smith SR, Middleton JP. Automated external defibrillators and survival from cardiac arrest in the outpatient hemodialysis clinic. *J Am Soc Nephrol* 2007;18(1):312–20. <https://doi.org/10.1681/ASN.2006040392>.
559. Pun PH, Leirich RW, Honeycutt EF, Herzog CA, Middleton JP. Modifiable risk factors associated with sudden cardiac arrest within hemodialysis clinics. *Kidney Int* 2011;79(2):218–27. <https://doi.org/10.1038/ki.2010.315>.
560. Jadoul M, Thumma J, Fuller DS, et al. Modifiable practices associated with sudden death among hemodialysis patients in the dialysis outcomes and practice patterns study. *Clin J Am Soc Nephrol* 2012;7(5):765–74. <https://doi.org/10.2215/CJN.08850811>.
561. Charytan DM, Winkelmayer WC, Granger CB, et al. Effects of dialysate potassium concentration of 3.0 mmol/l with sodium zirconium cyclosilicate on dialysis-free days versus dialysate potassium concentration of 2.0 mmol/l alone on rates of cardiac arrhythmias in hemodialysis patients with hyperkalemia. *Kidney Int* 2025;107(1):169–79. <https://doi.org/10.1016/j.kint.2024.10.010>.
562. Kluge HHP. WHO European regional obesity report 2022. Copenhagen: World Health Organization; 2022.
563. Welsh A, Hammad M, Piña IL, Kulinski J. Obesity and cardiovascular health. *European Journal of Preventive Cardiology* 2024;31(8):1026–35.
564. Considine J, Couper K, Greif R, et al. Cardiopulmonary resuscitation in obese patients: a scoping review. *Resuscitation Plus* 2024;20:100820.
565. Detsky ME, Jivraj N, Adhikari NK, et al. Will this patient be difficult to intubate? The rational clinical examination systematic review. *JAMA* 2019;321(5):493–503.
566. Hung KC, Chuang MH, Kang FC, et al. Prevalence and risk factors of difficult mask ventilation: a systematic review and meta-analysis. *Journal of Clinical Anesthesia* 2023;90:111197.
567. Kosmopoulos M, Kalra R, Alexy T, et al. The impact of BMI on arrest characteristics and survival of patients with out-of-hospital cardiac arrest treated with extracorporeal cardiopulmonary resuscitation. *Resuscitation* 2023;188:109842.
568. Kelly Jr RE. Pectus excavatum: historical background, clinical picture, preoperative evaluation and criteria for operation. *Semin Pediatr Surg* 2008;17(3):181–93. <https://doi.org/10.1053/j.sempedsurg.2008.03.002>.
569. Lee KH, Kim KW, Kim EY, et al. Proper compression landmark and depth for cardiopulmonary resuscitation in patients with pectus excavatum: a study using CT. *Emerg Med J* 2015;32(4):301–3. <https://doi.org/10.1136/emered-2013-202671>.
570. Russo V, Ranno M, Nigro G. Cardiopulmonary resuscitation in pectus excavatum patients: is it time to say more?. *Resuscitation* 2015;88:e5–6. <https://doi.org/10.1016/j.resuscitation.2014.07.023>.
571. Stearns JD, Twaibu J, Kwaku D, et al. Efficacy of standard chest compressions in patients with Nuss bars. *J Thorac Dis* 2020;12(8):4299–306. <https://doi.org/10.21037/jtd-20-702>.
572. Petrovic IS, Colombotto C, Urso F. Pectus excavatum and mechanical chest compression of a dangerous bond. *Am J Emerg Med* 2022;56:394.e5–7. <https://doi.org/10.1016/j.ajem.2022.03.016>.
573. Zoeller GK, Zallen GS, Glick PL. Cardiopulmonary resuscitation in patients with a Nuss bar—a case report and review of the literature. *J Pediatr Surg* 2005;40(11):1788–91. <https://doi.org/10.1016/j.jpedsurg.2005.07.036>.
574. Picton P, Walker D, White N, Deakin CD. Cardiopulmonary resuscitation following minimally invasive repair of pectus excavatum (Nuss technique). *Resuscitation* 2003;57(3):309–10. [https://doi.org/10.1016/S0300-9572\(03\)00041-8](https://doi.org/10.1016/S0300-9572(03)00041-8).
575. Collaboration. M-U, Felker A, Patel R, Kotnis R, Kenyon S, Knight M. Saving Lives, Improving Mothers' Care Compiled Report - Lessons learned to inform maternity care from the UK and Ireland Confidential Enquiries into Maternal Deaths and Morbidity 2020–22: National Perinatal Epidemiology Unit: University of Oxford, 2024.
576. Jeejeebhoy FM, Zelop CM, Lipman S, et al. Cardiac arrest in pregnancy: a scientific statement from the American heart association. *Circulation* 2015;132(18):1747–73. <https://doi.org/10.1161/CIR.0000000000000300>.
577. Association. OA. Quick Reference Handbook for Obstetric Emergencies: Obstetric Anaesthetists' Association. 2024.
578. Lipman S, Cohen S, Einav S, et al. The Society for Obstetric Anesthesia and Perinatology consensus statement on the

- management of cardiac arrest in pregnancy. *Anesth Analg* 2014;118(5):1003–16. <https://doi.org/10.1213/ANE.0000000000000171>.
579. ACOG Practice Bulletin No. 211: Critical Care in Pregnancy. *Obstet Gynecol* 2019;133(5):e303–e319. <https://doi.org/10.1097/AOG.0000000000003241>.
 580. Felker A, Knight M. MBRRACE-UK update: key messages from the UK and Ireland confidential enquiries into maternal death and morbidity 2024. *The Obstetrician & Gynaecologist* 2025;27(1):37–41. <https://doi.org/10.1111/tog.12965>.
 581. Chu J, Johnston TA, Geoghegan J, Royal College of O, Gynaecologists. Maternal Collapse in Pregnancy and the Puerperium: Green-top Guideline No. 56. *BJOG* 2020;127(5):e14–e52. <https://doi.org/10.1111/1471-0528.15995>.
 582. Regitz-Zagrosek V, Roos-Hesselink JW, Bauersachs J, et al. 2018 ESC Guidelines for the management of cardiovascular diseases during pregnancy. *Eur Heart J* 2018;39(34):3165–241. <https://doi.org/10.1093/eurheartj/ehy340>.
 583. Obstetric Anaesthetists' Association (UK). 2024.
 584. Zelop CM, Einav S, Mhyre JM, et al. Characteristics and outcomes of maternal cardiac arrest: a descriptive analysis of Get with the guidelines data. *Resuscitation* 2018;132:17–20. <https://doi.org/10.1016/j.resuscitation.2018.08.029>.
 585. Say L, Chou D, Gemmill A, et al. Global causes of maternal death: a WHO systematic analysis. *Lancet Glob Health* 2014;2(6):e323–33. [https://doi.org/10.1016/S2214-109X\(14\)70227-X](https://doi.org/10.1016/S2214-109X(14)70227-X).
 586. Beckett VA, Knight M, Sharpe P. The CAPS Study: incidence, management and outcomes of cardiac arrest in pregnancy in the UK: a prospective, descriptive study. *BJOG* 2017;124(9):1374–81. <https://doi.org/10.1111/1471-0528.14521>.
 587. Lucas DN, Kursumovic E, Cook TM, et al. Cardiac arrest in obstetric patients receiving anaesthetic care: results from the 7th national audit project of the royal college of anaesthetists. *Anaesthesia* 2024;79(5):514–23. <https://doi.org/10.1111/anae.16204>.
 588. Hasegawa J, Sekizawa A, Tanaka H, et al. Current status of pregnancy-related mortality in Japan: a report from the maternal death exploratory committee in Japan. *BMJ Open* 2016;6(3):e010304. <https://doi.org/10.1136/bmjopen-2015-010304>.
 589. Shields LE, Wiesner S, Klein C, Pelletreau B, Hedriana HL. Use of maternal early warning trigger tool reduces maternal morbidity. *Am J Obstet Gynecol* 2016;214(4):527.e1–6. <https://doi.org/10.1016/j.ajog.2016.01.154>.
 590. Umar A, Ameh CA, Muriithi F, Mathai M. Early warning systems in obstetrics: a systematic literature review. *PLoS One* 2019;14(5):e0217864. <https://doi.org/10.1371/journal.pone.0217864>.
 591. Quinn AC, Meek T, Waldmann C. Obstetric early warning systems to prevent bad outcome. *Curr Opin Anaesthesiol* 2016;29(3):268–72. <https://doi.org/10.1097/ACO.0000000000000338>.
 592. Chimwaza Y, Hunt A, Oliveira-Ciabatini L, et al. Early warning systems for identifying severe maternal outcomes: findings from the WHO global maternal sepsis study. *EClinicalMedicine* 2025;79:102981. <https://doi.org/10.1016/j.eclinm.2024.102981>.
 593. Krawczyk P, Dabrowska D, Guasch E, et al. Preparedness for severe maternal morbidity in European hospitals: the MaCriCare study. *Anaesth Crit Care Pain Med* 2024;43(3):101355. <https://doi.org/10.1016/j.accpm.2024.101355>.
 594. Chesnutt AN. Physiology of normal pregnancy. *Crit Care Clin* 2004;20(4):609–15. <https://doi.org/10.1016/j.ccc.2004.06.001>.
 595. Humphries A, Mirjalili SA, Tarr GP, Thompson JMD, Stone P. The effect of supine positioning on maternal hemodynamics during late pregnancy. *J Matern Fetal Neonatal Med* 2019;32(23):3923–30. <https://doi.org/10.1080/14767058.2018.1478958>.
 596. Lee AJ, Landau R. Aortocaval compression syndrome: time to revisit certain dogmas. *Anesth Analg* 2017;125(6):1975–85. <https://doi.org/10.1213/ANE.0000000000002313>.
 597. Fujita N, Higuchi H, Sakuma S, Takagi S, Latif M, Ozaki M. Effect of right-lateral versus left-lateral tilt position on compression of the inferior vena cava in pregnant women determined by magnetic resonance imaging. *Anesth Analg* 2019;128(6):1217–22. <https://doi.org/10.1213/ANE.0000000000004166>.
 598. Higuchi H, Takagi S, Zhang K, Furui I, Ozaki M. Effect of lateral tilt angle on the volume of the abdominal aorta and inferior vena cava in pregnant and nonpregnant women determined by magnetic resonance imaging. *Anesthesiology* 2015;122(2):286–93. <https://doi.org/10.1097/ALN.0000000000000553>.
 599. Ip JK, Campbell JP, Bushby D, Yentis SM. Cardiopulmonary resuscitation in the pregnant patient: a manikin-based evaluation of methods for producing lateral tilt. *Anaesthesia* 2013;68(7):694–9. <https://doi.org/10.1111/anae.12181>.
 600. Dohi S, Ichizuka K, Matsuoka R, Seo K, Nagatsuka M, Sekizawa A. Coronary perfusion pressure and compression quality in maternal cardiopulmonary resuscitation in supine and left-lateral tilt positions: a prospective, crossover study using mannequins and swine models. *Eur J Obstet Gynecol Reprod Biol* 2017;216:98–103. <https://doi.org/10.1016/j.ejogrb.2017.07.019>.
 601. Enomoto N, Yamashita T, Furuta M, et al. Effect of maternal positioning during cardiopulmonary resuscitation: a systematic review and meta-analyses. *BMC Pregnancy Childbirth* 2022;22(1):159. <https://doi.org/10.1186/s12884-021-04334-y>.
 602. Jeejeebhoy FM, Morrison LJ. Maternal cardiac arrest: a practical and comprehensive review. *Emerg Med Int* 2013;2013:274814. <https://doi.org/10.1155/2013/274814>.
 603. Kundra P, Khanna S, Habeebullah S, Ravishankar M. Manual displacement of the uterus during caesarean section. *Anaesthesia* 2007;62(5):460–5. <https://doi.org/10.1111/j.1365-2044.2007.05025.x>.
 604. Einav S, Kaufman N, Sela HY. Maternal cardiac arrest and perimortem caesarean delivery: evidence or expert-based?. *Resuscitation* 2012;83(10):1191–200. <https://doi.org/10.1016/j.resuscitation.2012.05.005>.
 605. Benson MD, Padovano A, Bourjeily G, Zhou Y. Maternal collapse: challenging the four-minute rule. *EBioMedicine* 2016;6:253–7. <https://doi.org/10.1016/j.ebiom.2016.02.042>.
 606. Chu JJ, Hinshaw K, Paterson-Brown S, et al. Perimortem caesarean section – why, when and how. *The Obstetrician & Gynaecologist* 2018;20(3):151–8. <https://doi.org/10.1111/tog.12493>.
 607. Siassakos D, Fox R, Crofts JF, Hunt LP, Winter C, Draycott TJ. The management of a simulated emergency: better teamwork, better performance. *Resuscitation* 2011;82(2):203–6. <https://doi.org/10.1016/j.resuscitation.2010.10.029>.
 608. Leech C, Nutbeam T, Chu J, et al. Maternal and neonatal outcomes following resuscitative hysterotomy for out of hospital cardiac arrest: a systematic review. *Resuscitation* 2025;207:110479. <https://doi.org/10.1016/j.resuscitation.2024.110479>.
 609. Wu Y, Luo J, Chen T, et al. Successful ECMO-assisted open chest cardiopulmonary resuscitation in a postpartum patient with delayed amniotic fluid embolism. *Eur J Med Res* 2022;27(1):19. <https://doi.org/10.1186/s40001-021-00628-1>.
 610. Burton A, Ratwatte S, Zalberg D, Morgan M, Narayan R, Cordina R. Cardiac arrest in pregnancy with successful stabilization and delivery on veno-arterial extracorporeal membrane oxygenation: a case report. *Eur Heart J Case Rep* 2024;8(11):ytac551. <https://doi.org/10.1093/ehjcr/ytac551>.
 611. Ramanathan K, Tan CS, Rycus P, et al. Extracorporeal membrane oxygenation in pregnancy: an analysis of the extracorporeal life support organization registry. *Crit Care Med* 2020;48(5):696–703. <https://doi.org/10.1097/CCM.0000000000004269>.
 612. Ducas RA, Elliott JE, Melnyk SF, et al. Cardiovascular magnetic resonance in pregnancy: insights from the cardiac hemodynamic imaging and remodeling in pregnancy (CHIRP) study. *J Cardiovasc Magn Reson* 2014;16(1):1. <https://doi.org/10.1186/1532-429X-16-1>.
 613. Delgado C, Dawson K, Schwaegler B, Zachariah R, Einav S, Bollag L. Hand placement during chest compressions in parturients: a pilot

- study to identify the location of the left ventricle using transthoracic echocardiography. *Int J Obstet Anesth* 2020;43:31–5. <https://doi.org/10.1016/j.ijoa.2020.05.003>.
614. Sheraton M, Columbus J, Surani S, Chopra R, Kashyap R. Effectiveness of mechanical chest compression devices over manual cardiopulmonary resuscitation: a systematic review with meta-analysis and trial sequential analysis. *West J Emerg Med* 2021;22(4):810–9. <https://doi.org/10.5811/westjem.2021.3.50932>.
 615. Myatra SN, Divatia JV, Brewster DJ. The physiologically difficult airway: an emerging concept. *Curr Opin Anaesthesiol* 2022;35(2):115–21. <https://doi.org/10.1097/ACO.0000000000001102>.
 616. Mushambi MC, Athanassoglou V, Kinsella SM. Anticipated difficult airway during obstetric general anaesthesia: narrative literature review and management recommendations. *Anaesthesia* 2020;75(7):945–61. <https://doi.org/10.1111/anae.15007>.
 617. Mushambi MC, Kinsella SM, Popat M, et al. Obstetric anaesthetists' association and difficult airway society guidelines for the management of difficult and failed tracheal intubation in obstetrics. *Anaesthesia* 2015;70(11):1286–306. <https://doi.org/10.1111/anae.13260>.
 618. Nanson J, Elcock D, Williams M, Deakin CD. Do physiological changes in pregnancy change defibrillation energy requirements?. *Br J Anaesth* 2001;87(2):237–9. <https://doi.org/10.1093/bja/87.2.237>.
 619. Nakamura E, Takahashi S, Matsunaga S, et al. Intravenous infusion route in maternal resuscitation: a scoping review. *BMC Emerg Med* 2021;21(1):151. <https://doi.org/10.1186/s12873-021-00546-9>.
 620. Excellence. NifHaC. Hypertension in pregnancy: diagnosis and management NICE guideline NG133. London: National Institute for Health and Care Excellence, 2019.
 621. Ahearn GS, Hadjiliadis D, Govert JA, Tapson VF. Massive pulmonary embolism during pregnancy successfully treated with recombinant tissue plasminogen activator: a case report and review of treatment options. *Arch Intern Med* 2002;162(11):1221–7. <https://doi.org/10.1001/archinte.162.11.1221>.
 622. Rigouzzo A, Froissant PA, Louvet N. Changing hemostatic management in post-partum hemorrhage. *Am J Hematol* 2024;99(Suppl 1):S13–8. <https://doi.org/10.1002/ajh.27264>.
 623. Robinson D, Basso M, Chan C, Duckitt K, Lett R. Guideline No. 431: Postpartum Hemorrhage and Hemorrhagic Shock. *J Obstet Gynaecol Can* 2022;44(12):1293–1310 e1. <https://doi.org/10.1016/j.jogc.2022.10.002>.
 624. Escobar MF, Nassar AH, Theron G, et al. FIGO recommendations on the management of postpartum hemorrhage 2022. *Int J Gynaecol Obstet* 2022;157 Suppl 1 (Suppl 1):3–50. <https://doi.org/10.1002/ijgo.14116>.
 625. Eyeberu A, Getachew T, Amare G, et al. Use of tranexamic acid in decreasing blood loss during and after delivery among women in Africa: a systematic review and meta-analysis. *Arch Gynecol Obstet* 2023;308(3):709–25. <https://doi.org/10.1007/s00404-022-06845-1>.
 626. Assis IC, Goveia CS, Miranda DB, Ferreira RS, Riccio LGC. Analysis of the efficacy of prophylactic tranexamic acid in preventing postpartum bleeding: systematic review with meta-analysis of randomized clinical trials. *Braz J Anesthesiol* 2023;73(4):467–76. <https://doi.org/10.1016/j.bjane.2022.08.002>.
 627. Yang F, Wang H, Shen M. Effect of preoperative prophylactic intravenous tranexamic acid on perioperative blood loss control in patients undergoing cesarean delivery: a systematic review and meta-analysis. *BMC Pregnancy Childbirth* 2023;23(1):420. <https://doi.org/10.1186/s12884-023-05753-9>.
 628. Collaborators WT. Effect of early tranexamic acid administration on mortality, hysterectomy, and other morbidities in women with post-partum haemorrhage (WOMAN): an international, randomised, double-blind, placebo-controlled trial. *Lancet* 2017;389(10084):2105–16. [https://doi.org/10.1016/s0140-6736\(17\)30638-4](https://doi.org/10.1016/s0140-6736(17)30638-4).
 629. Rittenberger JC, Kelly E, Jang D, Greer K, Heffner A. Successful outcome utilizing hypothermia after cardiac arrest in pregnancy: a case report. *Crit Care Med* 2008;36(4):1354–6. <https://doi.org/10.1097/CCM.0b013e318169ee99>.
 630. Song KH, Lee BK, Jeung KW, Lee SM. Safely completed therapeutic hypothermia in postpartum cardiac arrest survivors. *Am J Emerg Med* 2015;33(6):861.e5–6. <https://doi.org/10.1016/j.ajem.2014.12.042>.
 631. Banerjee A, Cantellow S. Maternal critical care: part I. *BJA Educ* 2021;21(4):140–7. <https://doi.org/10.1016/j.bjae.2020.12.003>.
 632. Merien AER, van de Ven J, Mol BW, Houterman S, Oei SG. Multidisciplinary team training in a simulation setting for acute obstetric emergencies: a systematic review. *Obstet Gynecol* 2010;115(5):1021–31. <https://doi.org/10.1097/AOG.0b013e3181d9f4cd>.
 633. Leonardsen AL, Svendsen EJ, Heitmann GB, et al. Development and validation of a questionnaire to assess healthcare personnel competence in cardiac arrest and resuscitation in pregnancy. *PLoS One* 2020;15(5):e0232984. <https://doi.org/10.1371/journal.pone.0232984>.
 634. Merriel A, Ficquet J, Barnard K, et al. The effects of interactive training of healthcare providers on the management of life-threatening emergencies in hospital. *Cochrane Database Syst Rev* 2019;9(9):CD012177. <https://doi.org/10.1002/14651858.CD012177.pub2>.
 635. Chou WK, Ullah N, Arjomandi Rad A, et al. Simulation training for obstetric emergencies in low- and lower-middle income countries: a systematic review. *Eur J Obstet Gynecol Reprod Biol* 2022;276:74–81. <https://doi.org/10.1016/j.ejogrb.2022.07.003>.
 636. Medicine. TFOIC, Society. IC. Guidelines for the provision of intensive care services. 2.1 ed. London: The Faculty of Intensive Care Medicine and Intensive Care Society, 2022.
 637. Burns R, Dent K. In: Burns R, Dent K, editors. *Advanced Life Support Group*. Hoboken, NJ, USA: Wiley Blackwell; 2022.
 638. Winter CDT, Muchatuta N, Crofts J, editors. *PROMPT Course Manual*. Cambridge University Press; 2018.
 639. McDermott BP, Casa DJ, Ganio MS, et al. Acute whole-body cooling for exercise-induced hyperthermia: a systematic review. *J Athl Train* 2009;44(1):84–93. <https://doi.org/10.4085/1062-6050-44.1.84>.